

North Carolina Division of Health Benefits Physician Administered Drug Program Catalog

•Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.

•11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacturer participating in the Medicaid Drug Rebate Program (MDRP).

•The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.

•The HCPCS Code effective date represents the date the HCPCS code was established

•Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacturer as they are not classified as covered outpatient drugs.

•Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/nccli-medicaid/medicaid-nccli-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Immune Globulins	90291	Cytomegalovirus immune globulin (CMV-IgV), human, for intravenous use	50 mL	1/1/2000	Cytogam®	cytomegalovirus immune globulin intravenous, human	Indicated for the prophylaxis of cytomegalovirus disease associated with transplantation of kidney, lung, liver, pancreas, and heart. In transplants of these organs other than kidney from CMV seropositive donors into seronegative recipients, prophylactic CMV-IgIV should be considered in combination with ganciclovir.	25.2	N/A	N/A	N/A	Y	Y	3/2024: Rebating Labeler Required field updated to align with policy that submitted NDCs must come from rebating labelers. Update not due to a change in policy.	3/28/2024
Immune Globulins	90371	Hepatitis B Immune Globulin (HBIG), human, for intramuscular use	1 mL	1/1/2000	HyperHEP B® S/D, Nabi-HB®	hepatitis b immune globulin, (human)	Indicated for treatment of acute exposure to blood containing HBSAg, perinatal exposure or infants born to HBSAg-positive mothers, sexual exposure to HBSAg-positive persons and household exposure to persons with acute HBV infection in the following settings: • Acute Exposure to Blood Containing HBSAg: Following either parenteral exposure (needlestick, bite, sharps), direct mucous membrane contact (accidental splash), or oral ingestion (pipetting accident), involving HBSAg-positive materials such as blood, plasma, or serum. • Perinatal Exposure of Infants Born to HBSAg-positive Mothers: Infants born to mothers positive for HBSAg with or without HBeAg.	18	N/A	N/A	N/A	Y	Y	3/2024: Rebating Labeler Required field updated to align with policy that submitted NDCs must come from rebating labelers. Update not due to a change in policy.	3/28/2024
Immune Globulins	90375	Rabies Immune Globulin (Rig), human, for intramuscular and/or subcutaneous use	150 IU	1/1/2000	HyperRAB® S/D, HyperRAB®	rabies immune globulin, (human) treated with solvent/detergent, for infiltration and intramuscular administration	HyperRAB S/D: Rabies vaccine and HyperRAB S/D should be given to all persons suspected of exposure to rabies with one exception: persons who have been previously immunized with rabies vaccine and have a confirmed adequate rabies antibody titer should receive only vaccine. HyperRAB S/D should be administered as promptly as possible after exposure, but can be administered up to the eighth day after the first dose of vaccine is given.	20	N/A	N/A	N/A	Y	Y		4/8/2020
Immune Globulins	90376	Rabies Immune Globulin, heat-treated (Rig-HT), human, for intramuscular and/or	150 IU	1/1/2000	Imogam® Rabies – HT	rabies immune globulin (human) USP, heat treated	Indicated for individuals suspected of exposure to rabies, particularly severe exposure, with one exception: persons who have been previously immunized with rabies vaccine prepared from human diploid cells (HDCV) in a pre-exposure or post exposure treatment series should receive only vaccine. Persons who indicated for passive, transient post-exposure prophylaxis of rabies infection to persons of all ages when given immediately after contact with a rabid or possibly rabid animal. Kedrab should be administered concurrently with a full course of rabies vaccine.	1	N/A	N/A	N/A	Y	Y		9/21/2018
Immune Globulins	90377	Rabies immune globulin, heat- and solvent/detergent-treated (Rig-HT S/D), human, for intramuscular and/or subcutaneous use	150 IU	1/1/2000	Kedrab™	rabies immune globulin (human) solution for intramuscular injection	• Do not exceed the recommended dose of Kedrab because this can partially suppress active production of rabies.	20	N/A	N/A	N/A	Y	Y		9/21/2022
Vaccines	90380	Respiratory syncytial virus, monoclonal antibody, seasonal dose; 0.5 mL dosage, for intramuscular use	0.5 mL	7/17/2023	Beyfortus™	nirsevimab-alip injection, for intramuscular use (0.5 mL dosage)	Indicated for the prevention of RSV lower respiratory tract disease in: • Neonates and infants born during or entering their first RSV season. • Children up to 24 months of age who remain vulnerable to severe RSV disease through their second RSV season.	1	N/A	24 months	N/A	Y	N		9/28/2023
Vaccines	90381	Respiratory syncytial virus, monoclonal antibody, seasonal dose; 1 mL dosage, for intramuscular use	1 mL	7/17/2023	Beyfortus™	nirsevimab-alip injection, for intramuscular use (1 mL dosage)	Indicated for the prevention of RSV lower respiratory tract disease in: • Neonates and infants born during or entering their first RSV season. • Children up to 24 months of age who remain vulnerable to severe RSV disease through their second RSV season.	2	N/A	24 months	N/A	Y	N		9/28/2023
Immune Globulins	90389	Tetanus Immune Globulin (Tig), human, for intramuscular use	250 units (1 mL)	1/1/2000	HyperTET® S/D	tetanus immune globulin (human)	Indicated for prophylaxis against tetanus following injury in patients whose immunization is incomplete or uncertain. It is also indicated, although evidence of effectiveness is limited, in the regimen of treatment of active cases of tetanus.	2	N/A	N/A	N/A	Y	Y		6/4/2019
Immune Globulins	90396	Varicella-zoster Immune Globulin (VZIG), human, for intramuscular use (Code Price is per 1 vial = 125 units)	125 units (1 vial)	1/1/2000	Varizig®	varicella zoster immune globulin (human) for intramuscular administration only	Indicated for post exposure prophylaxis in high risk individuals. High risk groups include: • immunocompromised children and adults, • newborns of mothers with varicella shortly before or after delivery, • premature infants, • infants less than one year of age, • adults without evidence of immunity, • pregnant women.	10	N/A	N/A	N/A	Y	Y		7/3/2018
Vaccines	90585	Bacillus Calmette-Guérin Vaccine (BCG) for tuberculosis, live, for percutaneous use.	50 mg	1/1/2000	BCG Vaccine	bacillus Calmette-Guérin vaccine (BCG) for tuberculosis, live, for percutaneous use.	Indicated for the prevention of tuberculosis (TB) in people not previously infected with Mycobacterium tuberculosis, who are at high risk for exposure.	1	N/A	N/A	N/A	Y	N		7/2/2018
Vaccines	90589	Chikungunya virus vaccine, live attenuated, for intramuscular use	0.5 mL (1 dose)	1/1/2024	Ixchiq	Chikungunya vaccine, live solution for intramuscular injection	Chikungunya Vaccine is indicated for the prevention of disease caused by chikungunya virus (CHIKV) in individuals 18 years of age and older who are at increased risk of exposure to CHIKV.	1	18 years	N/A	N/A	Y	N		3/22/2024
Vaccines	90611	Smallpox and monkeypox vaccine, attenuated vaccinia virus, live, non-replicating, preservative free, 0.5 mL dosage, suspension, for subcutaneous use	0.5 mL	7/26/2022	Jynneos™	smallpox and monkeypox vaccine, live, non-replicating suspension for subcutaneous and intradermal injection	FDA-Approved Indications: Indicated for prevention of smallpox and monkeypox disease in adults 18 years of age and older determined to be at high risk for smallpox or monkeypox infection. Emergency Use Authorization: The US Food and Drug Administration (FDA) has issued an Emergency Use Authorization (EUA) for the emergency use of Jynneos for: • active immunization by subcutaneous injection for prevention of monkeypox disease in individuals less than 18 years of age determined to be at high risk for monkeypox infection, and • active immunization by intradermal injection for prevention of monkeypox disease in individuals 18 years of age and older determined to be at high risk for monkeypox infection. Justification for Emergency Use of Jynneos During the Monkeypox Public Health Emergency There is currently an outbreak of monkeypox disease caused by monkeypox virus, an orthopoxvirus related to variola (the virus that causes smallpox disease). Following a 3-17 day incubation period, individuals infected with monkeypox virus develop painful lesions that progress sequentially through	2	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	N	Indication Specific Age Restrictions: FDA-Approved Indications: 18 years of age and older Emergency Use Authorization: N/A	5/31/2024

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

•Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.

•11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacture participating in the Medicaid Drug Rebate Program (MDRP).

•The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.

•The HCPCS Code effective date represents the date the HCPCS code was established

•Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacture as they are not classified as covered outpatient drugs.

•Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/nccli-medicaid/medicaid-nccli-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Vaccines	90619	Meningococcal conjugate vaccine, serogroups A, C, W, Y, quadrivalent, tetanus toxoid carrier (MenACWY-TT), for intramuscular use	0.5 mL	7/1/2009	MenQuadfi™	meningococcal (Groups A, C, Y, W) conjugate vaccine, solution for intramuscular injection	Indicated for active immunization for the prevention of invasive meningococcal disease caused by <i>Neisseria meningitidis</i> serogroups A, C, W, and Y. MenQuadfi vaccine is approved for use in individuals 2 years of age and older. MenQuadfi does not prevent <i>N. meningitidis</i> serogroup B disease.	1	2 years	N/A	N/A	Y	N		8/5/2021
Vaccines	90620	Meningococcal recombinant protein and outer membrane vesicle vaccine, serogroup B	0.5 mL	7/1/2017	Bexsero®	meningococcal group b vaccine suspension for intramuscular injection	Indicated for active immunization to prevent invasive disease caused by <i>Neisseria meningitidis</i> serogroup B. Bexsero is approved for use in individuals 10 through 25 years of age.	2	10 years	25 years	N/A	Y	N	12/2023: Maximum age restriction updated to align with FDA-approved and ACIP-	1/26/2024
Vaccines	90621	Meningococcal recombinant lipoprotein vaccine, serogroup B (MenB-FHbp), 2 or 3 dose schedule, for intramuscular use	0.5 mL	7/1/2017	Trumenba®	meningococcal group b vaccine suspension for intramuscular injection	Indicated for active immunization to prevent invasive disease caused by <i>Neisseria meningitidis</i> serogroup B. Trumenba is approved for use in individuals 10 through 25 years of age.	2	10 years	25 years	N/A	Y	N	12/2023: Maximum age restriction updated to align with FDA-approved and ACIP-recommended maximum age effective 10/1/2023.	1/26/2024
Vaccines	90623	Meningococcal pentavalent vaccine, conjugated Men A, C, W, Y- tetanus toxoid carrier, and Men B-FHbp, for intramuscular use	0.5 mL	7/1/2023	Penbraya™	meningococcal groups A, B, C, W, and Y vaccine, suspension for intramuscular injection	Meningococcal groups A, B, C, W, and Y vaccine, suspension for intramuscular injection is indicated for active immunization to prevent invasive disease caused by <i>Neisseria meningitidis</i> serogroups A, B, C, W, and Y. Penbraya is approved for use in individuals 10 through 25 years of age.	1	10 years	25 years	N/A	Y	N		1/26/2024
Vaccines	90625	Cholera vaccine, live, adult dosage, 1 dose schedule, for oral use	1 adult dosage (100 mL)	1/1/2016	Vaxchora®	cholera vaccine, live, oral suspension for oral administration	Indicated for active immunization against disease caused by <i>Vibrio cholerae</i> serogroup O1. Vaxchora is approved for use in persons 2 through 64 years of age traveling to cholera-affected areas. Limitations of Use: • The effectiveness of Vaxchora has not been established in persons living in cholera-affected areas. • The effectiveness of Vaxchora has not been established in persons who have pre-existing immunity due to previous exposure to <i>V. cholerae</i> or receipt of a cholera vaccine. • Vaxchora has not been shown to protect against disease caused by <i>V. cholerae</i> serogroup O139 or other	1	2 years	64 years	N/A	Y	N		10/27/2023
Vaccines	90626	Tick-borne encephalitis virus vaccine, inactivated; 0.25 mL dosage, for intramuscular use	0.25 mL	7/1/2021	TicoVac™	tick-borne encephalitis vaccine, suspension for intramuscular injection (0.25 mL dose)	Tick-borne encephalitis vaccine is indicated for active immunization to prevent tick-borne encephalitis (TBE). It is approved for use in individuals 1 year of age and older.	1	1 year	15 years	N/A	Y	Y	1/2024: Coverage effective 11/10/2023	1/4/2024
Vaccines	90627	Tick-borne encephalitis virus vaccine, inactivated; 0.5 mL dosage, for intramuscular use	0.5 mL	7/1/2021	TicoVac™	tick-borne encephalitis vaccine, suspension for intramuscular injection (0.5 mL dose)	Tick-borne encephalitis vaccine is indicated for active immunization to prevent tick-borne encephalitis (TBE). It is approved for use in individuals 1 year of age and older.	2	16 years	N/A	N/A	Y	Y	1/2024: Coverage effective 11/10/2023	1/4/2024
Vaccines	90632	Hepatitis A vaccine (Hep A), adult dosage, for intramuscular use	1 mL	1/1/2000	Havrix®, Vaqta®	hepatitis a vaccine, adult dosage, suspension for intramuscular injection	Indicated for active immunization against disease caused by hepatitis A virus (HAV). Approved for use in persons 12 months of age and older. Primary immunization should be administered at least 2 weeks prior to expected exposure to HAV.	1	19 years	N/A	N/A	Y	N		7/3/2018
Vaccines	90633	Hepatitis A vaccine (Hep A), pediatric/adolescent dosage - 2-dose schedule, for	0.5 mL	1/1/2000	Havrix®, Vaqta®	hepatitis a vaccine, pediatric/adolescent dosage - 2 dose schedule, for	Indicated for active immunization against disease caused by hepatitis A virus (HAV). Approved for use in persons 12 months of age and older. Primary immunization should be administered at least 2 weeks prior to expected exposure to HAV.	1	12 months	18 years	N/A	Y	N		7/3/2018
Vaccines	90636	Hepatitis A and Hepatitis B Vaccine (HepA-HepB), adult dosage, for intramuscular use	1 mL	1/1/2000	Twinrix®	hepatitis a & hepatitis b (recombinant) vaccine suspension for intramuscular injection	Indicated for active immunization against disease caused by hepatitis A virus and infection by all known subtypes of hepatitis B virus. Twinrix is approved for use in persons 18 years of age and older.	3	18 years	N/A	N/A	Y	N		9/12/2018
Vaccines	90647	Haemophilus influenzae type b vaccine (Hib), PRP-OMP conjugate, 3-dose schedule, for intramuscular use	0.5 mL	1/1/2000	PedvaxHib®	haemophilus b conjugate vaccine (meningococcal protein conjugate)	For routine vaccination against invasive disease caused by haemophilus influenzae type B in infants and children 2 – 71 months of age.	1	2 months	71 months	N/A	Y	N		7/2/2018
Vaccines	90648	Haemophilus influenzae b vaccine (Hib), PRP-T conjugate, 4-dose schedule, for intramuscular use	0.5 mL	1/1/2000	ActHIB®	haemophilus b conjugate vaccine (tetanus toxoid conjugate) solution for intramuscular injection	Indicated for the prevention of invasive disease caused by Haemophilus influenzae type b. ActHib vaccine is approved for use as a four dose series in infants and children 2 months through 5 years of age.	1	2 months	5 years	N/A	Y	N		7/3/2018

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

•Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.

•11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacture participating in the Medicaid Drug Rebate Program (MDRP).

•The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.

•The HCPCS Code effective date represents the date the HCPCS code was established

•Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacture as they are not classified as covered outpatient drugs.

•Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/ncci-medicaid/medicaid-ncci-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Vaccines	90651	Human Papillomavirus vaccine types 6, 11, 16, 18, 31, 33, 45, 52, 58, nonavalent (9VHPV), 2 or 3 dose schedule, for intramuscular use	0.5 mL	7/1/2017	Gardasil® 9	human papillomavirus 9-valent vaccine, recombinant suspension for intramuscular injection	Indicated in girls and women 9 through 45 years of age for the prevention of the following diseases: • Cervical, vulvar, vaginal, and anal cancer caused by HPV types 16, 18, 31, 33, 45, 52, and 58 • Genital warts (condyloma acuminata) caused by HPV types 6 and 11. The following precancerous or dysplastic lesions caused by HPV types 6, 11, 16, 18, 31, 33, 45, 52, and 58: • Cervical intraepithelial neoplasia (CIN) grade 2/3 and cervical adenocarcinoma in situ (AIS). • Cervical intraepithelial neoplasia (CIN) grade 1.	1	9 years	45 years	N/A	Y	N		7/28/2020
Vaccines	90662	Influenza virus vaccine (IV), split virus, preservative free, enhanced immunogenicity via increased antigen content, for intramuscular use	0.7 mL	1/1/2008	Fluzone® High-Dose Quadrivalent	influenza vaccine suspension for intramuscular injection	Indicated for active immunization for the prevention of influenza disease caused by influenza A subtype viruses and type B contained in the vaccine for use in persons 65 years of age and older.	1	65 years	N/A	N/A	Y	N		7/26/2023
Vaccines	90670	Pneumococcal conjugate vaccine, 13 valent (PCV13), for intramuscular use	0.5 mL	7/1/2009	Prenvar 13®	pneumococcal 13-valent conjugate vaccine (diphtheria CRM197 protein) suspension for intramuscular injection	In children 6 weeks through 5 years of age (prior to the 6th birthday), Prenvar 13 is indicated for: • Active immunization for the prevention of invasive disease caused by <i>Streptococcus pneumoniae</i> serotypes 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F and 23F. -Active immunization for the prevention of otitis media caused by <i>S. pneumoniae</i> serotypes 4, 6B, 9V, 14, 18C, 19F, and 23F. No otitis media efficacy data are available for serotypes 1, 3, 5, 6A, 7F, and 19A. In children 6 years through 17 years of age (prior to the 18th birthday), Prenvar 13 is indicated for: • Active immunization for the prevention of invasive disease caused by <i>S. pneumoniae</i> serotypes 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F and 23F. In adults 18 years of age and older, Prenvar 13 is indicated for: • Active immunization for the prevention of pneumonia and invasive disease caused by <i>S. pneumoniae</i> serotypes 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F and 23F.	1	6 weeks	N/A	N/A	Y	N		7/3/2018
Vaccines	90671	Pneumococcal conjugate vaccine, 15 valent (PCV15), for intramuscular use	0.5 mL (1 dose)	7/1/2021	Vaxneuvance™	pneumococcal 15-valent conjugate vaccine suspension for intramuscular injection	Indicated for active immunization for the prevention of invasive disease caused by <i>Streptococcus pneumoniae</i> serotypes 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F, 22F, 23F, and 33F in individuals 6 weeks of age and older. ACIP recommends use of PCV15 as an option for pneumococcal conjugate vaccination of persons aged <19 years, according to currently recommended PCV13 dosing and schedules.	1	6 weeks	N/A	N/A	Y	N	ACIP recommends for 6 weeks of age and older	10/20/2022
Vaccines	90672	Influenza virus vaccine, quadrivalent live (LAIV4), for intranasal use	0.2 mL	1/1/2013	FluMist® Quadrivalent	influenza virus vaccine, quadrivalent live, intranasal	Indicated for the active immunization of persons 2 – 49 years of age for the prevention of influenza disease caused by influenza A subtype viruses and type B viruses contained in the vaccine.	2	2 years	49 years	N/A	Y	N		9/21/2018
Vaccines	90674	Influenza virus vaccine, quadrivalent (cIIIV4), derived from cell cultures, subunit, preservative and antibiotic free, 0.5 mL dosage, for intramuscular use	0.5 mL	7/1/2016	Fluceelvax® Quadrivalent	influenza virus vaccine, suspension for intramuscular injection, preservative-free	Indicated for active immunization for the prevention of influenza disease caused by influenza virus subtypes A and type B contained in the vaccine.	2	6 months	N/A	N/A	Y	N		11/17/2021
Vaccines	90675	Rabies vaccine, for intramuscular use	1 mL	1/1/2000	Imovax® Rabies (Human Diploid-Cell Vaccine) and RabAvert® (Purified Chick Embryo Cell Culture)	rabies vaccine, for intramuscular use	Indicated for pre-exposure and post-exposure prophylaxis against rabies in all age groups.	5	N/A	N/A	N/A	Y	N		7/3/2018
Vaccines	90677	Pneumococcal conjugate vaccine, 20 valent (PCV20), for intramuscular use	0.5 mL	7/1/2021	Prenvar 20™	pneumococcal 20-valent conjugate vaccine, suspension for intramuscular injection	Prenvar 20 is a vaccine indicated for active immunization for the prevention of: • pneumonia caused by <i>S. pneumoniae</i> serotypes 1, 3, 4, 5, 6A, 6B, 7F, 8, 9V, 10A, 11A, 12F, 14, 15B, 18C, 19A, 19F, 22F, 23F, and 33F in individuals 18 years of age and older. • invasive disease caused by <i>Streptococcus pneumoniae</i> serotypes 1, 3, 4, 5, 6A, 6B, 7F, 8, 9V, 10A, 11A, 12F, 14, 15B, 18C, 19A, 19F, 22F, 23F, and 33F in individuals 6 weeks of age and older. • otitis media caused by <i>S. pneumoniae</i> serotypes 4, 6B, 9V, 14, 18C, 19F, and 23F in individuals 6 weeks through 5 years of age.	1	See Comments	N/A	N/A	Y	N	ACIP recommends for 2 months of age and older	9/28/2023

North Carolina Division of Health Benefits Physician Administered Drug Program Catalog

- Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.
- 11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacture participating in the Medicaid Drug Rebate Program (MDRP).
- The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.
- The HCPCS Code effective date represents the date the HCPCS code was established
- Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacture as they are not classified as covered outpatient drugs.
- Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/nccli-medicaid/medicaid-nccli-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Vaccines	90678	Respiratory syncytial virus vaccine, pref, subunit, bivalent, for intramuscular use	0.5 mL	1/1/2023	Abrysvo™	respiratory syncytial virus vaccine solution for intramuscular injection	Indicated for: - active immunization for the prevention of lower respiratory tract disease (LRTD) caused by respiratory syncytial virus (RSV) in individuals 60 years of age and older. - active immunization of pregnant individuals at 32 through 36 weeks gestational age for the prevention of lower respiratory tract disease (LRTD) and severe LRTD caused by respiratory syncytial virus (RSV) in infants from birth through 6 months of age.	1	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	N	Indication specific age restrictions: • Active immunization for the prevention of LRTD caused by RSV: 60 years of age and older • Active immunization of	1/26/2024
Vaccines	90679	Respiratory syncytial virus vaccine, pref, recombinant, subunit, adjuvanted, for intramuscular use	0.5 mL	5/3/2023	Arevvy	respiratory syncytial virus vaccine, adjuvanted suspension for intramuscular injection	Indicated for active immunization for the prevention of lower respiratory tract disease (LRTD) caused by respiratory syncytial virus in individuals 60 years of age and older.	1	60 years	N/A	N/A	Y	N		9/13/2023
Vaccines	90680	Rotavirus vaccine, pentavalent (RV5), 3 dose schedule, live, for oral use	2 mL	7/1/2005	RotaTeq®	rotavirus vaccine, live, oral, pentavalent	Indicated for the prevention of rotavirus gastroenteritis in infants and children caused by types G1, G2, G3, G4, and G9 when administered as a 3-dose series to infants between the ages of 6 to 32 weeks.	2	6 weeks	8 months	N/A	Y	N	ACIP recommends for 6 weeks of age to 8 months of age	3/30/2023
Vaccines	90681	Rotavirus vaccine, human, attenuated (RV1), 2 dose schedule, live, for oral use	1 mL	1/1/2008	Rotarix	rotavirus vaccine, live, oral	Indicated for the prevention of rotavirus gastroenteritis caused by G1 and non-G1 types (G3, G4, and G9). Rotarix is approved for use in infants 6 weeks to 24 weeks of age.	2	6 weeks	8 months	N/A	Y	N	ACIP recommends for 6 weeks of age to 8 months of age	3/30/2023
Vaccines	90682	Influenza virus vaccine, quadrivalent (RIV4), derived from recombinant DNA, hemagglutinin (HA) protein	1 dose (0.5 mL)	1/1/2017	Flublok® Quadrivalent	influenza vaccine, sterile solution for intramuscular injection	Indicated for active immunization against disease caused by influenza A subtype viruses and type B viruses contained in the vaccine.	1	18 years	N/A	N/A	Y	N		7/26/2023
Vaccines	90686	Influenza virus vaccine, quadrivalent (IV4), split virus, preservative free, 0.5 mL dosage, for intramuscular use	0.5 mL	1/1/2013	Afluria® Quadrivalent, Fluarix® Quadrivalent, FluLaval® Quadrivalent, Fluzone®	influenza vaccine suspension for intramuscular injection, preservative-free, 0.5 mL	Indicated for active immunization against influenza disease caused by influenza A subtype viruses and type B viruses contained in the vaccine.	2	Product Specific Age Restrictions (see comments)	N/A	N/A	Y	N	Product Specific Age Restrictions: Afluria Quad: 3 years and up Fluarix Quad, FluLaval Quad and Fluzone Quad: 6 months and up	8/10/2021
Vaccines	90687	Influenza virus vaccine, quadrivalent (IV4), split virus, 0.25 mL, for intramuscular use	0.25 mL	1/1/2013	Afluria® Quadrivalent, Fluzone® Quadrivalent	influenza virus vaccine, quadrivalent (IV4), split virus, 0.25 mL dosage, for intramuscular use	Indicated for active immunization for the prevention of influenza disease caused by influenza A subtype viruses and type B viruses contained in the vaccine.	2	6 months	35 months	N/A	Y	N		8/5/2020
Vaccines	90688	Influenza virus vaccine, quadrivalent (IV4), split virus, 0.5 mL dosage, for intramuscular use	0.5 mL	1/1/2013	Afluria® Quadrivalent, Fluzone® Quadrivalent	influenza vaccine suspension for intramuscular injection, 0.5 mL	Indicated for active immunization for the prevention of influenza disease caused by influenza A subtype viruses and type B viruses contained in the vaccine.	2	Product Specific Age Restrictions (see comments)	N/A	N/A	Y	N	Product Specific Age Restrictions: Afluria Quad: 3 years and up Fluzone Quad: 6 months and up	3/28/2023
Vaccines	90690	Typhoid vaccine, live, oral	4 capsules	1/1/2000	Vivotif®	typhoid vaccine live oral Ty21a	Indicated for immunization of adults and children greater than 6 years of age against disease caused by Salmonella typhi. Routine typhoid vaccination is not recommended in the United States of America. Selective immunization against typhoid fever is recommended for the following groups: 1) travelers to areas in which there is a recognized risk of exposure to S. typhi, 2) persons with intimate exposure (e.g. household contact) to a S. typhi carrier, and 3) microbiology laboratorians who work frequently with S. typhi.	1	6 years	N/A	N/A	Y	N		10/27/2023
Vaccines	90691	Typhoid vaccine, Vi capsular polysaccharide (ViCPS), for intramuscular use	0.5 mL	1/1/2000	Typhim Vi®	typhoid vi polysaccharide vaccine	Indicated for active immunization for the prevention of typhoid fever caused by S typhi and is approved for use in persons two years of age or older. Immunization with Typhim Vi vaccine should occur at least two weeks prior to expected exposure to S typhi. Typhim Vi vaccine is not indicated for routine immunization of individuals in the United States (US). Selective immunization against typhoid fever is recommended under the following circumstances: 1) travelers to areas where a recognized risk of exposure to typhoid exists, particularly ones who will have	1	2 years	N/A	N/A	Y	N		10/27/2023
Vaccines	90694	Influenza virus vaccine, quadrivalent (aIV4), inactivated, adjuvanted, preservative free, 0.5 mL dosage, for intramuscular use	0.5 mL	1/1/2020	Fluad® Quadrivalent	influenza vaccine, adjuvanted injectable emulsion for intramuscular use	Indicated for active immunization against influenza disease caused by influenza virus subtypes A and types B contained in the vaccine for use in persons 65 years of age and older.	1	65 years	N/A	N/A	Y	N		8/5/2020

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

•Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.

•11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacturer participating in the Medicaid Drug Rebate Program (MDRP).

•The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.

•The HCPCS Code effective date represents the date the HCPCS code was established

•Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacturer as they are not classified as covered outpatient drugs.

•Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/nccli-medicaid/medicaid-nccli-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Vaccines	90696	Diphtheria, tetanus toxoids, acellular pertussis vaccine and inactivated poliovirus vaccine, (DTaP-IPV), when	0.5 mL	1/1/2008	Kinrix [®] , Quadacel [™]	diphtheria and tetanus toxoids, acellular pertussis adsorbed and inactivated poliovirus vaccine,	• Kinrix: A single dose of Kinrix is indicated for active immunization against diphtheria, tetanus, pertussis, and poliomyelitis as the fifth dose in the diphtheria, tetanus, and acellular pertussis (DTaP) vaccine series and the fourth dose in the inactivated poliovirus vaccine (IPV) series in children 4 through 6 years of age whose previous DTaP vaccine doses have been with INFANRIX and/or PEDIARIX for the first three doses	1	4 years	6 years	N/A	Y	N		7/2/2018
Vaccines	90697	Diphtheria, tetanus toxoids, acellular pertussis vaccine, inactivated poliovirus vaccine, Haemophilus influenzae type b PRP-OMP	0.5 mL	1/1/2015	Vaxelis [™]	diphtheria and tetanus toxoids and acellular pertussis, inactivated poliovirus, haemophilus b conjugate and hepatitis B	Indicated for active immunization to prevent diphtheria, tetanus, pertussis, poliomyelitis, hepatitis B, and invasive disease due to Haemophilus influenzae type b. Vaxelis is approved for use as a 3-dose series in children from 6 weeks through 4 years of age (prior to the 5th birthday).	1	6 weeks	4 years	N/A	Y	N		12/20/2022
Vaccines	90698	Diphtheria, tetanus toxoids, acellular pertussis vaccine, Haemophilus influenzae type b, and inactivated poliovirus vaccine, (DTaP-IPV / Hib), for intramuscular use	0.5 mL	1/1/2004	Pentacel [®]	diphtheria and tetanus toxoids and acellular pertussis adsorbed, inactivated poliovirus and haemophilus b conjugate (tetanus toxoid conjugate)	Indicated for active immunization against diphtheria, tetanus, pertussis, poliomyelitis, and invasive disease due to Haemophilus influenzae type b. Pentacel vaccine is approved for use as a four dose series in children 6 weeks through 4 years of age (prior to fifth birthday).	1	6 weeks	4 years	N/A	Y	N		7/2/2018
Vaccines	90700	Diphtheria, tetanus toxoids, and acellular pertussis vaccine (DTaP), when administered to individuals younger than seven years, for intramuscular use	0.5 mL	1/1/2004	Daptacel [®] , Infanrix [®]	diphtheria, tetanus toxoids, and acellular pertussis vaccine adsorbed suspension for intramuscular injection	Indicated for active immunization against diphtheria, tetanus and pertussis as a five dose series in infants and children 6 weeks through 6 years of age (prior to 7th birthday).	1	6 weeks	6 years	N/A	Y	N		7/2/2018
Vaccines	90702	Diphtheria and tetanus toxoids adsorbed (DT) when administered to individuals younger than 7 years, for intramuscular use.	0.5 mL	1/1/2000	Diphtheria and Tetanus Toxoids, Adsorbed	diphtheria and tetanus toxoids (DT), adsorbed, for use in individuals younger than seven years, for intramuscular use.	Indicated for active immunization against diphtheria and tetanus. Diphtheria and Tetanus Toxoids Adsorbed is approved for use in children from 6 weeks through 6 years of age (prior to 7th birthday).	1	6 weeks	6 years	N/A	Y	N		7/2/2018
Vaccines	90707	Measles, mumps and rubella virus vaccine (MMR), live, for subcutaneous use	0.5 mL	1/1/2000	M-M-R [®] II	measles, mumps, and rubella virus vaccine live suspension for intramuscular or subcutaneous injection	Indicated for simultaneous vaccination against measles, mumps, and rubella in individuals 12 months of age or older.	1	12 months	N/A	N/A	Y	N	10/2023: HCPCS Effective Date updated from 1/1/2004 to 1/1/2000.	10/27/2023
Vaccines	90707	Measles, mumps and rubella virus vaccine (MMR), live, for subcutaneous use	0.5 mL	1/1/2000	Priorix	measles, mumps, and rubella vaccine, live, suspension for subcutaneous injection	Indicated for active immunization for the prevention of measles, mumps, and rubella in individuals 12 months of age and older.	2	12 months	N/A	N/A	Y	N		8/16/2022
Vaccines	90710	Measles, mumps, rubella, and varicella vaccine (MMRV), live, for subcutaneous use	0.5 mL	1/1/2000	ProQuad [®]	measles, mumps, rubella and varicella virus vaccine live suspension for intramuscular or subcutaneous injection	Indicated for active immunization for the prevention of measles, mumps, rubella, and varicella in children 12 months through 12 years of age.	1	12 months	12 years	N/A	Y	N		3/16/2023
Vaccines	90713	Poliovirus vaccine, Inactivated (IPV), for subcutaneous or intramuscular use	0.5 mL	7/1/2005	IPOL [®]	poliovirus vaccine, inactivated	Indicated for active immunization of infants (as young as 6 weeks of age), children and adults for the prevention of poliomyelitis caused by poliovirus types 1, 2, and 3.	2	6 weeks	N/A	N/A	Y	N		9/21/2018
Vaccines	90714	Tetanus and diphtheria toxoids adsorbed (Td), preservative free, when administered to individuals 7 years or older, for	0.5 mL	7/1/2005	Tenivac [®]	tetanus and diphtheria toxoids, adsorbed, suspension for intramuscular injection	Indicated for active immunization for the prevention of tetanus and diphtheria in persons 7 years of age and older.	2	7 years	N/A	N/A	Y	N		7/3/2018
Vaccines	90715	Tetanus, diphtheria toxoids and acellular pertussis vaccine (Tdap), when administered to individuals 7 years or older, for intramuscular use	0.5 mL	7/1/2005	Adacel [®] , Boostrix [®]	tetanus toxoid, reduced diphtheria toxoid and acellular pertussis vaccine adsorbed, suspension for intramuscular injection	Adacel: Indicated for: • active booster immunization against tetanus, diphtheria and pertussis. Adacel is approved for use in persons 10 through 64 years of age. • immunization during the third trimester of pregnancy to prevent pertussis in infants younger than 2 months of age. Boostrix: Indicated for: • active booster immunization against tetanus, diphtheria, and pertussis in individuals aged 10 years and older. • immunization during the third trimester of pregnancy to prevent pertussis in infants younger than 2	1	Min age restriction updated at the request of the State: 7 years	Product Specific Age Restrictions (see comments)	N/A	Y	N	Product specific maximum age restrictions: • Adacel: 64 years • Boostrix: N/A	2/23/2023
Vaccines	90716	Varicella virus vaccine (VAR), Live, for subcutaneous use	0.5 mL	1/1/2000	Varivax [®]	varicella virus vaccine live suspension for intramuscular or subcutaneous injection	Indicated for active immunization for the prevention of varicella in individuals 12 months of age and older.	2	12 months	N/A	N/A	Y	N		3/16/2023

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

- Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.
- 11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacture participating in the Medicaid Drug Rebate Program (MDRP).
- The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.
- The HCPCS Code effective date represents the date the HCPCS code was established
- Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacture as they are not classified as covered outpatient drugs.
- Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/nccli-medicaid/medicaid-nccli-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Vaccines	90717	Yellow fever vaccine, live, for subcutaneous use	0.5 mL	1/1/2000	YF-Vax*	yellow fever vaccine, for subcutaneous use	Indicated for active immunization for the prevention of yellow fever in persons 9 months of age and older in the following categories: • Persons Living in or Traveling to Endemic Areas: While the actual risk for contracting yellow fever during travel is probably low, variability of itineraries, behaviors and seasonal incidence of disease make it difficult to predict the actual risk for a given individual living in or traveling to a known endemic or epidemic area. Greater risk is associated with living in or traveling to areas of South America and Africa where yellow fever infection is officially reported at the time of travel and with traveling outside the urban areas of countries that do not officially report the disease but that lie in a yellow fever endemic zone. • Persons Traveling Internationally Through Countries with Yellow Fever: Some countries require an	1	9 months	N/A	N/A	Y	N		10/27/2023
Vaccines	90723	Diphtheria, tetanus toxoids, acellular pertussis vaccine, hepatitis B, and inactivated poliovirus vaccine.- (DTaP-HepB-IPV) for intramuscular use	0.5 mL	1/1/2001	Pediarix*	diphtheria and tetanus toxoids and acellular pertussis adsorbed, hepatitis b (recombinant) and inactivated poliovirus vaccine, suspension for intramuscular injection	Indicated for active immunization against diphtheria, tetanus, pertussis, infection caused by all known subtypes of hepatitis B virus, and poliomyelitis. Pediarix is approved for use as a three-dose series in infants born of hepatitis B surface antigen (HBsAg)-negative mothers. Pediarix may be given as early as 6 weeks of age through 6 years of age (prior to the 7th birthday).	1	6 weeks	6 years	N/A	Y	N		7/2/2018
Vaccines	90732	Pneumococcal polysaccharide vaccine, 23-valent (PPSV23), adult or immunosuppressed patient dosage, for use in individuals 2 years or older, for subcutaneous or intramuscular use	0.5 mL	1/1/2002	Pneumovax* 23	pneumococcal vaccine polyvalent sterile, liquid vaccine for intramuscular or subcutaneous injection	• Indicated for active immunization for the prevention of pneumococcal disease caused by the 23 serotypes contained in the vaccine (1, 2, 3, 4, 5, 6B, 7F, 8, 9N, 9V, 10A, 11A, 12F, 14, 15B, 17F, 18C, 19F, 19A, 20, 22F, 23F, and 33F). •Pneumovax 23 is approved for use in persons 50 years of age or older and persons aged greater than or equal to 2 years who are at increased risk for pneumococcal disease.	1	2 years	N/A	N/A	Y	N		7/3/2018
Vaccines	90734	Meningococcal conjugate vaccine, serogroups A, C, W, Y, quadrivalent, diphtheria toxoid carrier (MenACWY-D) or CRM197 carrier (MenACWY-CRM), for intramuscular use	0.5 mL	1/1/2017	Menactra*, Menveo	meningococcal (groups a, c, y, and w-135) polysaccharide diphtheria toxoid conjugate vaccine solution for intramuscular injection	Menactra: Indicated for active immunization to prevent invasive meningococcal disease caused by Neisseria meningitidis serogroups A, C, Y and W-135. Menactra is approved for use in individuals 9 months through 55 years of age. Menactra does not prevent N meningitidis serogroup B disease. Menveo: Indicated for active immunization to prevent invasive meningococcal disease caused by Neisseria meningitidis serogroups A, C, Y, and W-135 in individuals 2 months through 55 years of age. Menveo does	1	Product Specific Age Restrictions (see comments)	55 years	N/A	Y	N	Product specific age restrictions: • Menactra: 9 months through 55 years of age • Menveo: 2 months through 55 years of age	1/26/2024
Vaccines	90738	Japanese encephalitis virus vaccine, inactivated, for intramuscular use	0.5 mL	7/1/2008	Ixiaro*	Japanese encephalitis vaccine, inactivated, adsorbed suspension for intramuscular injection	Indicated for active immunization for the prevention of disease caused by Japanese encephalitis virus (JEV). Ixiaro is approved for use in individuals 2 months of age and older.	2	2 months	N/A	N/A	Y	N		10/27/2023
Vaccines	90739	Hepatitis B vaccine (HepB), CpG-adjuvanted, adult dosage, 2 dose or 4 dose schedule, for intramuscular use	0.5 mL	1/1/2013	Hepelisav-B*	hepatitis b vaccine (recombinant), adjuvanted solution for intramuscular injection	Indicated for prevention of infection caused by all known subtypes of hepatitis B virus in adults 18 years of age and older.	2	18 years	N/A	N/A	Y	N		6/6/2022
Vaccines	90740	Hepatitis B vaccine (HepB), dialysis or immunosuppressed patient dosage, 3-dose schedule, for intramuscular use	40 mcg	1/1/2001	Recombivax HB* Dialysis Formulation	hepatitis b vaccine, dialysis patient dosage (3 dose schedule), for intramuscular use	Recombivax HB Dialysis Formulation is approved for use in adult predialysis and dialysis patients 18 years of age and older for prevention of infection caused by all known subtypes of hepatitis B virus.	2	18 years	N/A	N/A	Y	N		10/31/2018
Vaccines	90743	Hepatitis B vaccine (HepB), adolescent, 2-dose schedule, for intramuscular use	1 mL	1/1/2001	Recombivax HB*	hepatitis B vaccine (recombinant) suspension for intramuscular injection (2 dose schedule)	Indicated for prevention of infection caused by all known subtypes of hepatitis B virus. Recombivax HB is approved for use in individuals of all ages. Recombivax HB Dialysis Formulation is approved for use in predialysis and dialysis patients 18 years of age and older.	1	11 years	15 years	N/A	Y	N		9/28/2021
Vaccines	90744	Hepatitis B vaccine (HepB), pediatric/adolescent dosage, 3-dose schedule, for intramuscular use	0.5 mL	1/1/2000	Engerix B* Pediatric, Recombivax HB* Pediatric	hepatitis b vaccine, pediatric/adolescent dosage (3 dose schedule), for intramuscular use	Hepatitis B vaccination is appropriate for people expected to receive human alpha-1 proteinase inhibitor that is produced from heat-treated, pooled human plasma that may contain the causative agents of hepatitis and other viral diseases.	2	N/A	19 years	N/A	Y	N		10/31/2018
Vaccines	90746	Hepatitis B vaccine (HepB), adult dosage, 3 dose schedule, for intramuscular use	1 mL	1/1/2000	Engerix B*, Recombivax HB*	hepatitis b vaccine (recombinant) suspension for intramuscular injection for adult use, 3 dose schedule	Indicated for immunization against infection caused by all known subtypes of hepatitis B virus.	1	20 years	N/A	N/A	Y	N		9/21/2018
Vaccines	90747	Hepatitis B vaccine (HepB), dialysis or immunosuppressed patient dosage, 4-dose schedule, for intramuscular use	40 mcg	1/1/2000	Engerix B*	hepatitis b vaccine, dialysis or immunosuppressed patient dosage (4 dose schedule), for intramuscular use	This schedule is designed for certain populations (e.g. dialysis patients, neonates born of hepatitis B-infected mothers, others who have or might have been recently exposed to the virus, certain travelers to high-risk areas) for immunization against infection caused by all known subtypes of hepatitis B virus.	2	N/A	N/A	N/A	Y	N		10/31/2018

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

- Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.
- 11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacturer participating in the Medicaid Drug Rebate Program (MDRP).
- The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.
- The HCPCS Code effective date represents the date the HCPCS code was established
- Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacturer as they are not classified as covered outpatient drugs.
- Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/nccli-medicaid/medicaid-nccli-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Vaccines	90750	Zoster (shingles) vaccine, (HZV), recombinant, sub-unit, adjuvanted, for intramuscular injection	0.5 mL	1/1/2017	Shingrix	zoster vaccine recombinant, adjuvanted, suspension for intramuscular injection	Indicated for prevention of herpes zoster (HZ) (shingles) in adults aged 50 years and older. Indicated for prevention of herpes zoster (HZ) (shingles) in adults aged 18 years and older who are or will be at increased risk of HZ due to immunodeficiency or immunosuppression caused by known disease or therapy. Limitations of Use: • Shingrix is not indicated for prevention of primary varicella infection (chickenpox).	2	19 years	N/A	N/A	Y	N	ACIP recommends for ≥ 19 years of age in immunodeficient or immunosuppressed adults	11/4/2021
Vaccines	90756	Influenza virus vaccine, quadrivalent (ccIV4), derived from cell cultures, subunit, antibiotic free, 0.5 mL dosage, for intramuscular use	0.5 mL	7/1/2017	Flucevax® Quadrivalent	influenza virus vaccine, suspension for intramuscular injection	Indicated for active immunization for the prevention of influenza disease caused by influenza virus subtypes A and type B contained in the vaccine.	2	6 months	N/A	N/A	Y	N		11/17/2021
Vaccines	90759	Hepatitis B vaccine (HepB), 3-antigen (S, Pre-S1, Pre-S2), 10 mcg dosage, 3 dose schedule, for intramuscular use	10 mcg	1/1/2022	PreHevbrio™	hepatitis b vaccine (recombinant) injectable suspension, for intramuscular use	Indicated for prevention of infection caused by all known subtypes of hepatitis B virus in adults 18 years of age and older.	2	18 years	N/A	N/A	Y	N		3/30/2022
Vaccines	91304	Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) (coronavirus disease [COVID-19]) vaccine, recombinant spike protein nanoparticle, saponin-based adjuvant, preservative free, 5	0.5 mL (5 mcg)	7/13/2022	N/A	Novavax COVID-19 Vaccine, Adjuvanted suspension for injection, for intramuscular use (2023-2024 Formula)	Emergency Use Authorization: The U.S. Food and Drug Administration (FDA) has issued an Emergency Use Authorization (EUA) for the emergency use of Novavax COVID-19 Vaccine, Adjuvanted (2023-2024 Formula) for active immunization to prevent coronavirus disease 2019 (COVID-19) caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) in individuals 12 years of age and older.	2	12 years	N/A	N/A	Y	N	9/2023: Aligned procedure code effective date with CMS effective date.	10/26/2023
Vaccines	91318	Severe acute respiratory syndrome coronavirus 2 (SARSCoV-2) (coronavirus	0.3 mL (3 mcg)	9/11/2023	N/A	Pfizer-BioNTech COVID-19 Vaccine suspension for injection, for intramuscular	The U.S. Food and Drug Administration (FDA) has issued an Emergency Use Authorization (EUA) for the emergency use of Pfizer-BioNTech COVID-19 Vaccine (2023-2024 Formula) for active immunization to prevent coronavirus disease 2019 (COVID-19) caused by severe acute respiratory syndrome coronavirus 2	2	6 months	4 years	N/A	Y	N		9/18/2023
Vaccines	91319	Severe acute respiratory syndrome coronavirus 2 (SARSCoV-2) (coronavirus disease [COVID-19]) vaccine, mRNA-LNP, spike protein, 10 mcg/0.3 mL dosage, trisucrose formulation, for intramuscular use	0.3 mL (10 mcg)	9/11/2023	N/A	Pfizer-BioNTech COVID-19 Vaccine suspension for injection, for intramuscular use - 5 years through 11 years of age (2023-2024 Formula)	The U.S. Food and Drug Administration (FDA) has issued an Emergency Use Authorization (EUA) for the emergency use of Pfizer-BioNTech COVID-19 Vaccine (2023-2024 Formula) for active immunization to prevent coronavirus disease 2019 (COVID-19) caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) in individuals 5 years through 11 years of age.	1	5 years	11 years	N/A	Y	N		9/18/2023
Vaccines	91320	Severe acute respiratory syndrome coronavirus 2 (SARSCoV-2) (coronavirus disease [COVID-19]) vaccine,	0.3 mL	9/11/2023	Comirnaty®	Pfizer-BioNTech COVID-19 Vaccine, mRNA suspension for injection, for intramuscular use - 12 years	Indicated for active immunization to prevent coronavirus disease 2019 (COVID-19) caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) in individuals 12 years of age and older.	1	12 years	N/A	N/A	Y	N		9/18/2023
Vaccines	91321	Severe acute respiratory syndrome coronavirus 2 (SARSCoV-2) (coronavirus disease [COVID-19]) vaccine, mRNA-	0.25 mL	9/11/2023	N/A	Moderna COVID-19 Vaccine Suspension for injection, for intramuscular use - 6 months through 11 years of age	The U.S. Food and Drug Administration (FDA) has issued an Emergency Use Authorization (EUA) for the emergency use of Moderna COVID-19 Vaccine (2023-2024 Formula) for active immunization to prevent coronavirus disease 2019 (COVID-19) caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) in individuals 6 months through 11 years of age.	1	6 months	11 years	N/A	Y	N		9/18/2023
Vaccines	91322	Severe acute respiratory syndrome coronavirus 2 (SARSCoV-2) (coronavirus disease [COVID-19]) vaccine, mRNA-LNP, 50 mcg/0.5 mL dosage, for intramuscular use	0.5 mL	9/11/2023	Spikevax™	Moderna COVID-19 Vaccine, mRNA Suspension for injection, for intramuscular use - 12 years of age and older (2023-2024 Formula)	Indicated for active immunization to prevent coronavirus disease 2019 (COVID-19) caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) in individuals 12 years of age and older.	1	12 years	N/A	N/A	Y	N		9/18/2023

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

•Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.

•11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacture participating in the Medicaid Drug Rebate Program (MDRP).

•The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.

•The HCPCS Code effective date represents the date the HCPCS code was established

•Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacture as they are not classified as covered outpatient drugs.

•Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/ncci-medicaid/medicaid-ncci-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Drugs	J0121	Injection, omadacycline, 1 mg	1 mg	10/1/2019	Nuzyra™	omadacycline for injection, for intravenous use	Indicated for the treatment of adult patients with the following infections caused by susceptible microorganisms: • Community-acquired bacterial pneumonia (CABP) • Acute bacterial skin and skin structure infections (ABSSSI) To reduce the development of drug-resistant bacteria and maintain the effectiveness of Nuzyra and other	1,500	18 years	N/A	N/A	Y	Y		9/27/2019
Drugs	J0122	Injection, eravacycline, 1 mg	1 mg	10/1/2019	Xerava™	eravacycline for injection, for intravenous use	Indicated for the treatment of complicated intra-abdominal infections in patients 18 years of age and older. Limitations of Use: Xerava is not indicated for the treatment of complicated urinary tract infections (cUTI).	7,000	18 years	N/A	N/A	Y	Y		9/27/2019
Biologics	J0129	Injection, abatacept, 10 mg	10 mg	1/1/2007	Orencia®	abatacept injection, for intravenous use	Treatment of: • Adult Rheumatoid Arthritis (RA): moderately to severely active RA in adults. Orencia may be used as monotherapy or concomitantly with DMARDs other than TNF antagonists. • Juvenile Idiopathic Arthritis: moderately to severely active polyarticular juvenile idiopathic arthritis in patients 2 years of age and older. Orencia may be used as monotherapy or concomitantly with methotrexate. • Active Psoriatic Arthritis (PsA) in adults. Indicated for prophylaxis of: • Acute graft versus host disease (aGVHD): in combination with a calcineurin inhibitor and methotrexate, in adults and pediatric patients 2 years of age and older undergoing hematopoietic stem cell transplantation (HSCT) from a matched or 1 allele-mismatched unrelated donor. Important Limitations of Use: • Should not be given concomitantly with TNF antagonists.	400	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	Indication specific age restrictions: • RA and PsA: 18 years of age and older • JIA and aGVHD: 2 years of age and older	1/14/2022
Drugs	J0133	Injection, acyclovir, 5 mg	5 mg	1/1/2006	N/A	acyclovir sodium, for injection, for intravenous infusion	Indicated for: • Herpes simplex infections in immunocompromised patients • Initial episodes of herpes genitalis • Herpes simplex encephalitis • Neonatal herpes simplex virus infection • Varicella-zoster infections in immunocompromised patients	8,400	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	Indication specific age restrictions: • Herpes Simplex Infections: Mucosal and Cutaneous Herpes Simplex (HSV-1 and HSV-2) Infections in Immunocompromised	5/14/2019
Drugs	J0153	Injection, adenosine, 1 mg, (not to be used to report any adenosine phosphate	1 mg	1/1/2015	Adenocard®, Adenoscan®	adenosine injection, for intravenous use	Adenoscan: Adjunct to thallium-201 myocardial perfusion scintigraphy in patients unable to exercise adequately.	118	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	Product specific age restrictions: Adenoscan: 18 years of age	5/6/2019
Drugs	J0171	Injection, adrenalin, epinephrine, 0.1 mg	0.1 mg	1/1/2011	Adrenalin®	epinephrine injection, for intramuscular or subcutaneous use	Indicated for emergency treatment of allergic reactions (Type 1), including anaphylaxis	N/A	N/A	N/A	N/A	Y	Y		10/26/2018
Biologics	J0177	Injection, aflibercept hd, 1 mg	1 mg	4/1/2024	Eylea® HD	aflibercept injection, for intravitreal use	Indicated for the treatment of patients with: • Neovascular (Wet) Age-Related Macular Degeneration (nAMD) • Diabetic Macular Edema (DME) • Diabetic Retinopathy (DR)	32	18 years	N/A	N/A	Y	Y		4/12/2024
Biologics	J0178	Injection, aflibercept, 1 mg	1 mg	1/1/2013	Eylea®	aflibercept injection for intravitreal injection	Indicated for: • Neovascular (Wet) Age-Related Macular Degeneration (AMD) • Macular Edema Following Retinal Vein Occlusion (RVO)	8	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	AMD, RVO, DME, DR: 18 years of age and older ROP: N/A	3/16/2023
Biologics	J0179	Injection, brolicizumab-dbl, 1 mg	1 mg	1/1/2020	Beovu®	brolicizumab-dbl injection, for intravitreal injection	Indicated for the treatment of: - Neovascular (Wet) Age-Related Macular Degeneration (AMD) - Diabetic Macular Edema (DME)	24	18 years	N/A	N/A	Y	Y		6/9/2022
Drugs	J0180	Injection, agalsidase beta, 1 mg	1 mg	1/1/2005	Fabryzyme®	agalsidase beta injection, powder, lyophilized for solution for intravenous use	Indicated for treatment of adult and pediatric patients 2 years of age and older with confirmed Fabry disease.	420	2 years	N/A	N/A	Y	Y		4/26/2021

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

•Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.

•11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacture participating in the Medicaid Drug Rebate Program (MDRP).

•The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.

•The HCPCS Code effective date represents the date the HCPCS code was established

•Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacture as they are not classified as covered outpatient drugs.

•Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/nccli-medicaid/medicaid-nccli-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Drugs	J0184	Injection, amisulpride, 1 mg	1 mg	1/1/2024	Barhemsys®	amisulpride injection, for intravenous use	Indicated in adults for: • Prevention of postoperative nausea and vomiting (PONV), either alone or in combination with an antiemetic of a different class. • Treatment of PONV in patients who have received antiemetic prophylaxis with an agent of a different class or have not received prophylaxis.	50	18 years	N/A	N/A	Y	Y		12/22/2023
Drugs	J0185	Injection, aprepitant, 1 mg	1 mg	1/1/2019	Cirvanti™	aprepitant injectable emulsion, for intravenous use	Indicated in adults, in combination with other antiemetic agents, for the prevention of: • acute and delayed nausea and vomiting associated with initial and repeat courses of highly emetogenic	650	18 years	N/A	N/A	Y	Y	9/2023: Max monthly units updated from 390 units to 650	9/28/2023
Biologics	J0202	Injection, alemtuzumab, 1 mg	1 mg	1/1/2016	Lemtrada®	alemtuzumab injection, for intravenous use	Indicated for the treatment of patients with relapsing forms of multiple sclerosis (MS).	60	17 years	N/A	N/A	Y	Y		7/2/2018
Drugs	J0207	Injection, amifostine, 500 mg	500 mg	1/1/2000	Ethyl®	amifostine for injection	Indicated to: • Reduce the incidence of moderate to severe xerostomia in patients undergoing postoperative radiation treatment of head and neck cancer. • Reduce the cumulative renal toxicity associated with repeated administration of cisplatin in patients with advanced ovarian cancer, where the radiation port includes a substantial portion of the parotid glands.	155	18 years	N/A	N/A	Y	Y		9/25/2018
Drugs	J0208	Injection, sodium thiosulfate (pedmark), 100 mg	100 mg	4/1/2023	Pedmark®	sodium thiosulfate injection, for intravenous use	Indicated to reduce the risk of ototoxicity associated with cisplatin in pediatric patients 1 month of age and older with localized, non-metastatic solid tumors.	5,000	1 month	18 years	N/A	Y	Y		3/22/2024
Drugs	J0210	Injection, methyldopate HCl, up to 250mg	250 mg	1/1/2000	N/A	methyldopate hydrochloride injection	Indicated for hypertension, when parenteral medication is indicated. The treatment of hypertensive crises may be initiated with methyldopate HCl injection.	496	N/A	N/A	N/A	Y	Y		10/26/2018
Biologics	J0217	Injection, velmanase alfa-tycv, 1 mg	1 mg	1/1/2024	Lamzede®	velmanase alfa-tycv for injection, for intravenous use	Indicated for the treatment of non-central nervous system manifestations of alpha-mannosidosis in adult and pediatric patients.	700	N/A	N/A	N/A	Y	Y		12/21/2023
Biologics	J0218	Injection, olipudase alfa-rpcp, 1 mg	1 mg	4/1/2023	Xenpozyme™	olipudase alfa-rpcp for injection, for intravenous use	Indicated for treatment of non-central nervous system manifestations of acid sphingomyelinase deficiency (ASMD) in adult and pediatric patients.	1,260	N/A	N/A	N/A	Y	Y		3/16/2023
Biologics	J0219	Injection, avalglucosidase alfa-ngpt, 4 mg	4 mg	4/1/2022	Nexviazyme™	avalglucosidase alfa-ngpt for injection, for intravenous use	Indicated for the treatment of patients 1 year of age and older with late-onset Pompe disease (lysosomal acid alpha-glucosidase [GAA] deficiency).	2,100	1 year	N/A	N/A	Y	Y		3/17/2022
Biologics	J0221	Injection, alglucosidase alfa, (Lumizyme), 10 mg	10 mg	1/1/2012	Lumizyme®	alglucosidase alfa for injection, for intravenous use	A hydrolytic lysosomal glycogen-specific enzyme indicated for patients with Pompe disease (GAA deficiency).	900	N/A	N/A	N/A	Y	Y		6/4/2019
Drugs	J0222	Injection, Patisiran, 0.1 mg	0.1 mg	10/1/2019	Onpatro™	patisiran lipid complex injection, for intravenous use	Indicated for the treatment of the polyneuropathy of hereditary transthyretin-mediated amyloidosis in adults.	600	18 years	N/A	N/A	Y	Y		9/27/2019
Drugs	J0223	Injection, givosiran, 0.5 mg	0.5 mg	7/1/2020	Givlaari™	givosiran injection, for subcutaneous use	Indicated for the treatment of adults with acute hepatic porphyria (AHP).	1,512	18 years	N/A	N/A	Y	Y		6/17/2020
Drugs	J0224	Injection, lumasiran, 0.5 mg	0.5 mg	7/1/2021	Oxlumo™	lumasiran injection, for subcutaneous use	Indicated for the treatment of primary hyperoxaluria type 1 (PH1) to lower urinary and plasma oxalate levels in pediatric and adult patients.	1,890	N/A	N/A	N/A	Y	Y		11/30/2022
Drugs	J0225	Injection, vutrisiran, 1 mg	1 mg	1/1/2023	Amvuttra™	vutrisiran injection, for subcutaneous use	Indicated for the treatment of the polyneuropathy of hereditary transthyretin-mediated amyloidosis in adults.	25	18 years	N/A	N/A	Y	Y		12/6/2022
Drugs	J0248	Injection, remdesivir, 1 mg	1 mg	12/23/2021	Veklury®	remdesivir injection, for intravenous use	Indicated for the treatment of coronavirus disease 2019 (COVID-19) in adults and pediatric patients (birth to less than 18 years of age weighing at least 1.5 kg) who are: • Hospitalized, or • Not hospitalized and have mild-to-moderate COVID-19, and are at high risk for progression to severe	400		Pediatric patients from birth to less than 28 days of age weighing at least 1.5	N/A	Y	Y		3/22/2024
Biologics	J0256	Injection, alpha 1-proteinase inhibitor, human, 10 mg, not otherwise specified	10 mg	1/1/2000	Aralast NP®, Prolastin-C®, Zemaira®	alpha 1-proteinase inhibitor (human) for intravenous use	Indicated for chronic augmentation therapy in adults with clinically evident emphysema due to severe congenital deficiency of Alpha1-Pi (alpha1-antitrypsin deficiency).	5,000	18 years	N/A	N/A	Y	Y		6/6/2019

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

•Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.

•11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacture participating in the Medicaid Drug Rebate Program (MDRP).

•The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.

•The HCPCS Code effective date represents the date the HCPCS code was established

•Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacture as they are not classified as covered outpatient drugs.

•Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/nccli-medicaid/medicaid-nccli-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Biologics	J0257	Injection, alpha-1 proteinase inhibitor (human), (Glassia), 10 mg	10 mg	1/1/2012	Glassia™	alpha 1-proteinase inhibitor (human) injection solution, for intravenous use	Indicated for chronic augmentation and maintenance therapy in adults with clinically evident emphysema due to severe hereditary deficiency of Alpha1-Pi (alpha1-antitrypsin deficiency). Glassia increases antigenic and functional (anti-neutrophil elastase capacity, ANEC) serum levels and antigenic lung epithelial lining fluid levels of alpha1-Pi. Limitations of Use: • The effect of augmentation therapy with any Alpha1-Pi, including Glassia, on pulmonary exacerbations and on the progression of emphysema in alpha1-antitrypsin deficiency has not been conclusively demonstrated in randomized, controlled clinical trials. • Clinical data demonstrating the long-term effects of chronic augmentation and maintenance therapy of individuals with Glassia are not available. • Glassia is not indicated as therapy for lung disease in patients in whom severe Alpha1-Pi deficiency has	4,200	18 years	N/A	N/A	Y	Y		9/25/2018
Drugs	J0278	Injection, amikacin sulfate, 100 mg	100 mg	1/1/2006	N/A	amikacin sulfate injection, solution	Indicated in the short-term treatment of serious infections due to susceptible strains of Gram-negative bacteria, including Pseudomonas species, Escherichia coli, species of indole-positive and indole-negative Proteus, Providencia species, Klebsiella-Enterobacter-Serratia species, and Acinetobacter (Mima-Herellea) species.	150	N/A	N/A	N/A	Y	Y		4/10/2019
Drugs	J0280	Injection, aminophylline, up to 250mg	up to 250 mg	1/1/2000	N/A	aminophylline injection	Indicated as an adjunct to inhaled beta-2 selective agonists and systemically administered corticosteroids for the treatment of acute exacerbations of the symptoms and reversible airflow obstruction associated with asthma and other chronic lung diseases, e.g., emphysema and chronic bronchitis.	217	N/A	N/A	N/A	Y	Y		9/25/2018
Drugs	J0285	Injection, amphotericin B, 50 mg	50 mg	1/1/2000	N/A	amphotericin B for injection	Amphotericin B for injection is specifically intended to treat potentially life-threatening fungal infections: aspergillosis, cryptococcosis (torulosis), North American blastomycosis, systemic candidiasis,	93	N/A	N/A	N/A	Y	Y		9/25/2018
Drugs	J0287	Injection, amphotericin B lipid complex, 10 mg	10 mg	1/1/2003	Abelcet®	amphotericin B lipid complex injection	Indicated for the treatment of invasive fungal infections in patients who are refractory to or intolerant of conventional amphotericin B therapy.	2,170	N/A	N/A	N/A	Y	Y		5/6/2019
Drugs	J0289	Injection, amphotericin B liposome, 10 mg	10 mg	1/1/2003	AmBisome®	amphotericin B liposome for injection	Indicated for: • Empirical therapy for presumed fungal infection in febrile, neutropenic patients • Treatment of patients with Aspergillus species, Candida species, and/or Cryptococcus species infections refractory to amphotericin B desoxycholate, or in patients where renal impairment or unacceptable toxicity precludes the use of amphotericin B desoxycholate • Treatment of Cryptococcal Meningitis in HIV-infected patients • Treatment of visceral leishmaniasis. In immunocompromised patients with visceral leishmaniasis treated with AmBisome, relapse rates were high following initial clearance of parasites.	2,604	1 month	N/A	N/A	Y	Y		4/10/2019
Drugs	J0290	Injection, ampicillin sodium, 500 mg	500 mg	1/1/2000	N/A	ampicillin sodium for injection, for intravenous or intramuscular use	Indicated in the treatment of infections caused by susceptible strains of the designated organisms in the following conditions: • Respiratory Tract Infections caused by Streptococcus pneumoniae, Staphylococcus aureus (penicillinase and nonpenicillinase-producing), H. influenzae, and Group A beta-hemolytic streptococci. • Bacterial Meningitis caused by E. coli, Group B streptococci, and other Gram-negative bacteria (Listeria monocytogenes, N. meningitidis). The addition of an aminoglycoside with ampicillin may increase its effectiveness against Gram-negative bacteria.	1,736	N/A	N/A	N/A	Y	Y		4/10/2019
Drugs	J0291	Injection, plazomicin, 5 mg	5 mg	10/1/2019	Zemdri™	plazomicin injection, for intravenous use	• Indicated for the treatment of patients 18 years of age or older with complicated urinary tract infections (CUTI) including pyelonephritis. • As only limited clinical safety and efficacy data are available, reserve Zemdri for use in patients who have limited or no alternative treatment options. • To reduce the development of drug-resistant bacteria and maintain effectiveness of Zemdri and other antibacterial drugs, Zemdri should be used only to treat infections that are proven or strongly suspected to be caused by susceptible microorganisms.	2,940	18 years	N/A	N/A	Y	Y		10/3/2019
Drugs	J0295	Injection, ampicillin sodium/sulbactam sodium, per 1.5 gm	per 1.5 gm	1/1/2000	Unasyn®	ampicillin sodium and sulbactam sodium injection, powder, for solution	Indicated for the treatment of infection due to susceptible strains of the designated microorganisms in the conditions listed below: • Skin and skin structure infections caused by beta-lactamase producing strains of Staphylococcus aureus, Escherichia coli, Klebsiella spp. (including K. pneumoniae), Proteus mirabilis, Bacteroides fragilis, Enterobacter spp., and Acinetobacter calcoaceticus.	168	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	Indication specific: • Skin and skin structure infections: 1 year of age and older • Intra-abdominal infections:	6/7/2019

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

•Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.

•11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacturer participating in the Medicaid Drug Rebate Program (MDRP).

•The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.

•The HCPCS Code effective date represents the date the HCPCS code was established

•Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacture as they are not classified as covered outpatient drugs.

•Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/ncci-medicaid/medicaid-ncci-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Drugs	J0300	Injection, amobarbital, up to 125mg	up to 125 mg	1/1/2000	Amytal®	amobarbital sodium for injection	Indicated for use as a: • Sedative	112	6 years	N/A	N/A	Y	Y		4/10/2019
Drugs	J0330	Injection, succinylcholine chloride, up to 20mg	up to 20 mg	1/1/2000	Anectine® , Quelicin™	succinylcholine chloride injection	Indicated as an adjunct to general anesthesia, to facilitate tracheal intubation, and to provide skeletal muscle relaxation during surgery or mechanical ventilation.	8	N/A	N/A	N/A	Y	Y		9/21/2018
Drugs	J0349	Injection, rezafungin, 1 mg	1 mg	10/1/2023	Rezzayo™	rezafungin for injection, for intravenous use	Indicated in patients 18 years of age or older who have limited or no alternative options for the treatment of candidemia and invasive candidiasis.	1,000	18 years	N/A	N/A	Y	Y		9/28/2023
Drugs	J0360	Injection, hydralazine HCl, up to 20mg	up to 20 mg	1/1/2000	N/A	hydralazine hydrochloride injection	Indicated for severe essential hypertension when the drug cannot be given orally or when there is an urgent need to lower blood pressure.	75	N/A	N/A	N/A	Y	Y		6/4/2019
Drugs	J0401	Injection, aripiprazole, extended release, 1 mg	1 mg	1/1/2014	Abilify Maintena®	aripiprazole extended-release injectable suspension, for intramuscular use	Indicated for the treatment of schizophrenia in adults. Indicated for maintenance monotherapy treatment of bipolar I disorder in adults.	800	18 years	N/A	N/A	Y	Y		5/20/2019
Drugs	J0402	Injection, aripiprazole (abilify asimtufil), 1 mg	1 mg	1/1/2024	Abilify Asimtufil®	aripiprazole extended-release injectable suspension, for	Indicated: • for the treatment of schizophrenia in adults	960	18 years	N/A	N/A	Y	Y		12/21/2023
Drugs	J0456	Injection, azithromycin, 500 mg	500 mg	1/1/2000	Zithromax®	azithromycin for intravenous infusion	Indicated for mild to moderate infections caused by designated, susceptible bacteria in community-acquired pneumonia in adults and pelvic inflammatory disease.	10	16 years	N/A	N/A	Y	Y		9/25/2018
Drugs	J0461	Injection, atropine sulfate, 0.01 mg	0.01 mg	1/1/2010	N/A	atropine sulfate injection for intravenous, intramuscular, subcutaneous, intraosseous, or endotracheal use	Indicated for temporary blockade of severe or life threatening muscarinic effects.	27,900	N/A	N/A	N/A	Y	Y		10/4/2018
Drugs	J0470	Injection, dimercaprol, per 100mg	per 100 mg	1/1/2000	BAL in oil™	dimercaprol injection	Indicated in the treatment of: • Arsenic, gold and mercury poisoning. • Acute lead poisoning when used concomitantly with Edetate Calcium Disodium Injection. Dimercaprol is effective for use in acute poisoning by mercury salts if therapy is begun within one or two hours following ingestion. It is not very effective for chronic mercury poisoning. Dimercaprol is of questionable value in poisoning by other heavy metals such as antimony and bismuth. It should not be used in iron, cadmium, or selenium poisoning because the resulting dimercaprol-metal complexes are more toxic than the metal alone, especially to the kidneys.	252	N/A	N/A	N/A	Y	Y		6/7/2019
Drugs	J0475	Injection, baclofen, 10 mg	10 mg	1/1/2000	Gablofen®, Lioresal® Intrathecal	baclofen injection	Indicated for use in the management of severe spasticity of cerebral or spinal origin in adult and pediatric patients age 4 years and above. • Baclofen intrathecal should be reserved for patients unresponsive to oral baclofen therapy, or those who experience intolerable central nervous system side effects at effective doses.	8	4 years	N/A	N/A	Y	Y	5/2023: NC Suggested Max Monthly Units updated to align with NCTracks, which has been set to 8 units/month since	9/13/2023
Drugs	J0476	Injection, baclofen, 50 mcg, for intrathecal trial	50 mcg	1/1/2000	Gablofen®, Lioresal® Intrathecal	baclofen injection, for intrathecal trial	Management of severe spasticity caused by spinal cord lesions or multiple sclerosis. Baclofen also is used intrathecally in patients with spasticity of cerebral origin, including those with cerebral palsy and acquired brain injury. Baclofen injection is designated an orphan drug by the FDA for the management of spasticity in patients with cerebral palsy.	5	N/A	N/A	N/A	Y	Y		5/21/2019
Biologicals	J0485	Injection, belatacept, 1 mg	1 mg	1/1/2013	Nulojix®	belatacept for injection, for intravenous use	Prophylaxis of organ rejection in adult patients receiving a kidney transplant. Use in combination with basiliximab induction, mycophenolate mofetil, and corticosteroids. Limitations of Use: • Use only in patients who are EBV seropositive. • Use has not been established for the prophylaxis of organ rejection in transplanted organs other than the kidney.	6,000	18 years	N/A	N/A	Y	Y		6/6/2019
Biologicals	J0490	Injection, belimumab, 10 mg	10 mg	1/1/2012	Benlysta®	belimumab injection, for intravenous use	Indicated for the treatment of patients aged 5 years and older with active, autoantibody-positive, systemic lupus erythematosus who are receiving standard therapy. Indicated for the treatment of patients aged 5 years and older with active lupus nephritis who are receiving standard therapy.	420	5 years	N/A	N/A	Y	Y		8/16/2022
Biologicals	J0491	Injection, anifrolumab-fnia, 1 mg	1 mg	4/1/2022	Saphnelo™	anifrolumab-fnia injection, for intravenous use	Indicated for the treatment of adult patients with moderate to severe systemic lupus erythematosus (SLE), who are receiving standard therapy.	600	18 years	N/A	N/A	Y	Y		3/21/2022

North Carolina Division of Health Benefits Physician Administered Drug Program Catalog

- Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.
- 11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacture participating in the Medicaid Drug Rebate Program (MDRP).
- The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.
- The HCPCS Code effective date represents the date the HCPCS code was established
- Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacture as they are not classified as covered outpatient drugs.
- Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/ncci-medicaid/medicaid-ncci-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Drugs	J0500	Injection, dicyclomine HCl, up to 20mg	up to 20 mg	1/1/2000	Bentyl®	dicyclomine hydrochloride injection for intramuscular use	Indicated for the treatment of functional bowel/irritable bowel syndrome.	8	18 years	N/A	N/A	Y	Y		4/10/2019
Drugs	J0515	Injection, benzotropine mesylate, per 1 mg	1 mg	1/1/2000	Cogentin®	benztropine mesylate injection	Indicated: - for use as an adjunct in the therapy of all forms of parkinsonism. - for use in the control of extrapyramidal disorders (except tardive dyskinesia) due to neuroleptic drugs (e.g., phenothiazines).	248	3 years	N/A	N/A	Y	Y		11/17/2021
Drugs	J0558	Injection, penicillin G benzathine and penicillin G procaine, 100,000 units	100,000 units	1/1/2011	Bicillin® C-R	penicillin G benzathine and penicillin G procaine injectable suspension	Indicated for the treatment of moderately severe infections due to penicillin G-susceptible microorganisms that are susceptible to serum levels common to this particular dosage form. Therapy should be guided by bacteriological studies (including susceptibility testing) and by clinical response. Bicillin C-R is indicated in the treatment of the following in adults and pediatric patients: • Moderately severe to severe infections of the upper-respiratory tract, scarlet fever, erysipelas, and skin and soft-tissue infections due to susceptible streptococci. NOTE: Streptococci in Groups A, C, G, H, L, and M are very sensitive to penicillin G. Other groups, including Group D (enterococci), are resistant. Penicillin	96	N/A	N/A	N/A	Y	Y		8/24/2018
Drugs	J0561	Injection, penicillin G benzathine, 100,000 units	100,000 units	1/1/2011	Bicillin® L-A	penicillin G benzathine injectable suspension	Indicated for the treatment of infections due to penicillin G-sensitive microorganisms that are susceptible to the low and very prolonged serum levels common to this particular dosage form. Therapy should be guided by bacteriological studies (including sensitivity tests) and by clinical response. The following infections will usually respond to adequate dosage of intramuscular penicillin G benzathine: mild to moderate upper respiratory infections due to susceptible streptococci, venereal infections (syphilis, yaws, bejel, and pinta) and prophylaxis of rheumatic fever and chorea.	96	N/A	N/A	N/A	Y	Y		8/24/2018
Biologicals	J0565	Injection, bezlotroxumab, 10 mg	10 mg	1/1/2018	Zinplava™	bezlotroxumab injection, for intravenous use	Indicated to reduce recurrence of <i>Clostridioides difficile</i> infection (CDI) in adult and pediatric patients 1 year of age or older who are receiving antibacterial drug treatment for CDI and are high risk for CDI recurrence.	140	1 year	N/A	N/A	Y	Y		6/19/2023
Biologicals	J0567	Injection, cerliponase alfa, 1 mg	1 mg	1/1/2019	Brineura®	cerliponase alfa injection, for intraventricular use	Indicated to slow the loss of ambulation in symptomatic pediatric patients 3 years of age and older with late infantile neuronal ceroid lipofuscinosis type 2 (CLN2), also known as tripeptidyl peptidase 1 (TPP1) deficiency.	900	3 years	N/A	N/A	Y	Y		7/2/2018
Drugs	J0570	Buprenorphine implant, 74.2 mg	74.2 mg = 1 implant	1/1/2017	Probuphine®	buprenorphine implant for subdermal administration (CIII)	Indicated for the maintenance treatment of opioid dependence in patients who have achieved and sustained prolonged clinical stability on low-to-moderate doses of a transmucosal buprenorphine-containing product (i.e., doses of no more than 8 mg per day of Subutex® or Suboxone® sublingual tablet or generic equivalent). Probuphine should be used as part of a complete treatment program to include counseling and	4	16 years	N/A	N/A	Y	Y		9/27/2018
Drugs	J0577	Injection, buprenorphine extended-release (brixadi), less than or equal to 7 days of therapy	1 syringe	4/1/2024	Brixadi™	buprenorphine extended-release injection for subcutaneous use CIII (weekly)	Indicated for the treatment of moderate to severe opioid use disorder in patients who have initiated treatment with a single dose of a transmucosal buprenorphine product or who are already being treated with buprenorphine. Brixadi should be used as part of a complete treatment plan that includes counseling and psychosocial support.	5	18 years	N/A	N/A	Y	Y		3/22/2024
Drugs	J0578	Injection, buprenorphine extended-release (brixadi), greater than 7 days and up to 28 days of therapy	1 syringe	4/1/2024	Brixadi™	buprenorphine extended-release injection for subcutaneous use CIII (monthly)	Indicated for the treatment of moderate to severe opioid use disorder in patients who have initiated treatment with a single dose of a transmucosal buprenorphine product or who are already being treated with buprenorphine. Brixadi should be used as part of a complete treatment plan that includes counseling and psychosocial support.	2	18 years	N/A	N/A	Y	Y		3/22/2024

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

•Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.

•11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacture participating in the Medicaid Drug Rebate Program (MDRP).

•The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.

•The HCPCS Code effective date represents the date the HCPCS code was established

•Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacture as they are not classified as covered outpatient drugs.

•Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/nccli-medicaid/medicaid-nccli-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Biologics	J0584	Injection, bursosumab-twza 1 mg	1 mg	1/1/2019	Crysvita*	bursosumab-twza injection, for subcutaneous use	Indicated for: <ul style="list-style-type: none"> • The treatment of X-linked hypophosphatemia (XLH) in adult and pediatric patients 6 months of age and older. • The treatment of FGF23-related hypophosphatemia in tumor-induced osteomalacia (TIO) associated with phosphaturic mesenchymal tumors that cannot be curatively resected or localized in adult and pediatric patients 2 years of age and older. 	540	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	Indication specific age restrictions: <ul style="list-style-type: none"> • XLH: 6 months of age and older • TIO: 2 years of age and older 	7/28/2020
Biologics	J0585	Injection, onabotulinumtoxinA, 1 unit	1 unit	1/1/2000	Botox*	onabotulinumtoxinA for injection, for intramuscular, intradetrusor, or intradermal use	Indicated for: <ul style="list-style-type: none"> • Treatment of overactive bladder (OAB) with symptoms of urge urinary incontinence, urgency, and frequency, in adults who have an inadequate response to or are intolerant of an anticholinergic medication • Treatment of urinary incontinence due to detrusor overactivity associated with a neurologic condition (e.g., spinal cord injury (SCI), multiple sclerosis (MS)) in adults who have an inadequate response to or are intolerant of an anticholinergic medication • Treatment of neurogenic detrusor overactivity (NDO) in pediatric patients 5 years of age and older who have an inadequate response to or are intolerant of anticholinergic medication. • Prophylaxis of headaches in adult patients with chronic migraine (≥15 days per month with headache lasting 4 hours a day or longer) 	600 in 90 day interval	N/A	N/A	N/A	Y	Y	1/2023: NC Suggested Max Monthly Units updated to align with NCTracks, which has been set to 600 units in 90 days since 1/1/2019. 9/2023: NC Suggested Max Monthly Units updated from 3 month interval to 90 day interval to align with NCTracks. 11/2023: Edited 1/2023 and	11/3/2023
Biologics	J0586	implant, 1 microgram	5 units	1/1/2010	Dysport*	abobotulinumtoxinA for injection, for intramuscular use	<ul style="list-style-type: none"> • Treatment of adults with cervical dystonia. • The temporary improvement in the appearance of moderate to severe glabellar lines associated with procerus and corrugator muscle activity in adult patients <65 years of age. • Treatment of spasticity in patients 2 years of age and older. 	300	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	Indication specific recommendations. <ul style="list-style-type: none"> • Cervical Dystonia: 18 years of age and older 	8/25/2020
Biologics	J0587	Injection, rimabotulinumtoxinB, 100 units	100 units	1/1/2002	Myobloc*	rimabotulinumtoxin B injection	Indicated for: <ul style="list-style-type: none"> - Treatment of adult patients with cervical dystonia to reduce the severity of abnormal head position and neck pain associated with cervical dystonia. - Treatment of chronic sialorrhea in adults. 	100	18 years	N/A	N/A	Y	Y		9/27/2019
Biologics	J0588	Injection, incobotulinumtoxinA, 1 unit	1 unit	1/1/2012	Xeomin*	incobotulinumtoxinA for injection, for intramuscular or intraglandular use	Indicated for the treatment or improvement of: <ul style="list-style-type: none"> • Chronic sialorrhea in patients 2 years of age and older • Upper limb spasticity in adults • Upper limb spasticity in pediatric patients 2 to 17 years of age, excluding spasticity caused by cerebral palsy 	600 in a 12-week interval	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	Indication specific age restrictions: <ul style="list-style-type: none"> Cervical dystonia and blepharospasm: 18 years of age and older 	9/13/2023
Drugs	J0594	Injection, busulfan, 1 mg	1 mg	1/1/2007	Busulfex*	busulfan injection for intravenous use	Indicated for use in combination with cyclophosphamide as a conditioning regimen prior to allogeneic hematopoietic progenitor cell transplantation for chronic myelogenous leukemia (CML).	1,312	N/A	N/A	N/A	Y	Y	<ul style="list-style-type: none"> • Upper Limb Spasticity: Safety and effectiveness in pediatric patients below the age of 2 years have not been established. 	9/27/2018
Drugs	J0595	Injection, butorphanol tartrate, 1mg	1 mg	1/1/2004	N/A	butorphanol tartrate injection	Indicated: <ul style="list-style-type: none"> • As a preoperative or pre-anesthetic medication • As a supplement to balanced anesthesia • For the relief of pain during labor, and • For the management of pain severe enough to require an opioid analgesic and for which alternative 	992	18 years	N/A	N/A	Y	Y	<ul style="list-style-type: none"> • Lower Limb Spasticity: Safety and effectiveness in pediatric patients below the age of 2 years have not been established. 	9/27/2018
Biologics	J0596	Injection, c-1 esterase inhibitor (recombinant), Ruconest, 10 units	10 units	1/1/2016	Ruconest*	c1 esterase inhibitor (recombinant) for intravenous use, lyophilized	Indicated for treatment of acute attacks in adult and adolescent patients with hereditary angioedema (HAE).	3,360	N/A	N/A	N/A	Y	Y		4/10/2019

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

- Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.
- 11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacturer participating in the Medicaid Drug Rebate Program (MDRP).
- The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.
- The HCPCS Code effective date represents the date the HCPCS code was established
- Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacture as they are not classified as covered outpatient drugs.
- Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/ncci-medicaid/medicaid-ncci-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Biologicals	J0597	Injection, C-1 esterase inhibitor (human), Berinert, 10 units	10 units	1/1/2011	Berinert*	c1 esterase inhibitor (human) for intravenous use	Treatment of acute abdominal, facial, or laryngeal hereditary angioedema (HAE) attacks in adult and pediatric patients.	1,120	N/A	N/A	N/A	Y	Y		4/10/2019
Biologicals	J0598	Injection, C1 esterase inhibitor (human), Cinryze, 10 units	10 units	1/1/2010	Cinryze*	c1 esterase inhibitor (human) for intravenous use	Indicated for routine prophylaxis against angioedema attacks in adults, adolescents and pediatric patients (6 years of age and older) with hereditary angioedema (HAE).	2,750	6 years	N/A	N/A	Y	Y		7/26/2018
Drugs	J0600	Injection, edetate calcium disodium, up to 1000 mg	up to 1000 mg	1/1/2000	Calcium Disodium Versanate	edetate calcium disodium injection for intravenous or intramuscular use	Indicated for the reduction of blood levels and depot stores of lead in lead poisoning (acute and chronic) and lead encephalopathy in both pediatric populations and adults.	15	N/A	N/A	N/A	Y	Y		10/10/2018
Drugs	J0606	Injection, etelcalcetide, 0.1 mg	0.1 mg	1/1/2018	Parsabiv™	etelcalcetide injection, for intravenous use	Indicated for secondary hyperparathyroidism (HPT) in adult patients with chronic kidney disease (CKD) on hemodialysis. Limitations of Use: Parsabiv has not been studied in adult patients with parathyroid carcinoma, primary hyperparathyroidism or with CKD who are not on hemodialysis and is not recommended for use in these populations.	2,250	18 years	N/A	N/A	Y	Y		6/4/2019
Drugs	J0612	Injection, calcium gluconate, not otherwise specified, 10 mg	10 mg	4/1/2023	N/A	calcium gluconate injection, for intravenous use	Indicated for pediatric and adult patients for the treatment of acute symptomatic hypocalcemia. Limitations of Use: The safety of calcium gluconate injection for long term use has not been established.	124,000	N/A	N/A	N/A	Y	Y		3/22/2024
Drugs	J0613	Injection, calcium gluconate (wg critical care), not therapeutically equivalent to	10 mg	4/1/2023	N/A	calcium gluconate injection, for intravenous use (WG Critical Care)	Calcium Gluconate in Sodium Chloride Injection is a form of calcium indicated for pediatric and adult patients for the treatment of acute symptomatic hypocalcemia.	24,800	N/A	N/A	N/A	Y	Y		3/22/2024
Drugs	J0636	Injection, calcitriol, 0.1 mcg	0.1 mcg	1/1/2003	N/A	calcitriol injection	Indicated in the management of hypocalcemia in patients undergoing chronic renal dialysis. It has been shown to significantly reduce elevated parathyroid hormone levels. Reduction of PTH has been shown to result in an improvement in renal osteodystrophy.	560	13 years	N/A	N/A	Y	Y		9/27/2018
Biologicals	J0638	Injection, canakinumab, 1 mg	1 mg	1/1/2011	Ilaris*	canakinumab injection, for subcutaneous use	Indicated for the treatment of: - Periodic Fever Syndromes: • Cryopyrin-Associated Periodic Syndromes (CAPS), in adults and children 4 years of age and older including: Familial Cold Autoinflammatory Syndrome (FCAS) and Muckle-Wells Syndrome (MWS). • Tumor Necrosis Factor Receptor Associated Periodic Syndrome (TRAPS) in adult and pediatric patients. • Hyperimmunoglobulin D Syndrome (HIDS)/Mevalonate Kinase Deficiency (MKD) in adult and pediatric patients. • Familial Mediterranean Fever (FMF) in adult and pediatric patients. - Active Still's Disease:	600	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	Indication specific age restrictions: • SJA, AQSD, TRAPS, HIDS/MKD, and FMF: 2 years of age and older • CAPS (FCAS and MWS): 4 years of age and older • Gout flares: 18 years of age and older	9/28/2023
Drugs	J0640	Injection, leucovorin calcium, per 50 mg	50 mg	1/1/2000	N/A	leucovorin calcium for injection for intravenous or intramuscular use	Indicated: • After high dose methotrexate therapy in osteosarcoma. • To diminish the toxicity and counteract the effects of impaired methotrexate elimination and of inadvertent overdosages of folic acid antagonists. • In the treatment of megaloblastic anemias due to folic acid deficiency when oral therapy is not feasible. • For use in combination with 5-fluorouracil to prolong survival in the palliative treatment of patients with advanced colorectal cancer. Leucovorin should not be mixed in the same infusion as 5-fluorouracil because a precipitate may form.	80	N/A	N/A	N/A	Y	Y		7/2/2018
Drugs	J0641	Injection, levoleucovorin, not otherwise specified, 0.5 mg	0.5 mg	1/1/2009	Fusilev*	levoleucovorin injection solution for intravenous use	Indicated for: • Rescue after high-dose methotrexate therapy in osteosarcoma. • Diminishing the toxicity and counteracting the effects of impaired methotrexate elimination and of inadvertent overdosage of folic acid antagonists. • Use in combination chemotherapy with 5-fluorouracil in the palliative treatment of patients with advanced metastatic colorectal cancer. Limitations of Use: Fusilev is not approved for pernicious anemia and megaloblastic anemias. Improper use may cause a hematologic remission while neurologic manifestations continue to progress.	10,000	N/A	N/A	N/A	Y	Y		10/3/2019

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

•Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.

•11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacturer participating in the Medicaid Drug Rebate Program (MDRP).

•The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.

•The HCPCS Code effective date represents the date the HCPCS code was established

•Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacture as they are not classified as covered outpatient drugs.

•Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/ncci-medicaid/medicaid-ncci-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Drugs	J0642	Injection, levoleucovorin (khapzory), 0.5 mg	0.5 mg	10/1/2019	Khapzory™	levoleucovorin for injection, for intravenous use	Indicated for: • Rescue after high-dose methotrexate therapy in patients with osteosarcoma. • Diminishing the toxicity associated with overdosage of folic acid antagonists or impaired methotrexate elimination. • Treatment of patients with metastatic colorectal cancer in combination with fluorouracil. Limitations of Use: Khapzory is not indicated for the treatment of pernicious anemia and megaloblastic anemia secondary to lack of vitamin B12 because of the risk of progression of neurologic manifestations despite hematologic remission.	4,800	N/A	N/A	N/A	Y	Y		10/3/2019
Drugs	J0665	Injection, bupivacaine, not otherwise specified, 0.5 mg	0.5 mg	7/1/2023	Marcaïne™, Sensorcaine®	bupivacaine hydrochloride injection, for infiltration, perineural, caudal, epidural,	Bupivacaine hydrochloride injection: • Indicated in adults for the production of local or regional anesthesia or analgesia for surgery, dental and oral surgery procedures, diagnostic and therapeutic procedures, and for obstetrical procedures. For each	4,000	Formulation-specific age restrictions (see comments)	N/A	N/A	Y	Y	Formulation-specific age restrictions: • Bupivacaine hydrochloride	10/26/2023
Drugs	J0670	Injection, mepivacaine hydrochloride, per 10 mL	10 mL	1/1/2000	Carbocaine™, Polocaine®, Polocaine® MPF	mepivacaine hydrochloride injection	Carbocaine, Polocaine and Polocaine MPF: Indicated for production of local or regional analgesia and anesthesia by local infiltration, peripheral nerve block techniques, and central neural techniques including epidural and caudal blocks.	50	N/A	N/A	N/A	Y	Y		4/10/2019
Drugs	J0688	Injection, cefazolin sodium (hikma), not therapeutically equivalent to J0690, 500 mg	500 mg	1/1/2024	N/A	cefazolin for injection, for intravenous use (Hikma)	Cefazolin for injection is a cephalosporin antibacterial indicated for: • Treatment of the following infections caused by susceptible isolates of the designated microorganisms in adult and pediatric patients 1 month of age and older for whom appropriate dosing with this formulation can be achieved: o Respiratory tract infections o Urinary tract infections o Skin and skin structure infections o Biliary tract infections o Bone and joint infections o Genital infections o Septicemia o Endocarditis • Perioperative prophylaxis in adult patients To reduce the development of drug-resistant bacteria and maintain the effectiveness of cefazolin for injection and other antibacterial drugs, cefazolin for injection should be used only to treat or prevent infections that are proven or strongly suspected to be caused by bacteria.	372	1 month	N/A	N/A	Y	Y		1/11/2024
Drugs	J0689	Injection, cefazolin sodium (baxter), not therapeutically equivalent to J0690, 500 mg	500 mg	1/1/2023	N/A	cefazolin injection, for intravenous use (Baxter)	Indicated for: • Treatment of the following infections caused by susceptible isolates of the designated microorganisms in adult and pediatric patients for whom appropriate dosing with this formulation can be achieved: o Respiratory tract infections o Urinary tract infections o Skin and skin structure infections o Biliary tract infections o Bone and joint infections o Genital infections o Septicemia o Endocarditis • Perioperative prophylaxis in adults and pediatric patients aged 10 to 17 years old for whom appropriate dosing with this formulation can be achieved. To reduce the development of drug-resistant bacteria and maintain the effectiveness of cefazolin injection and other antibacterial drugs, cefazolin injection should be used only to treat or prevent infections that are proven or strongly suspected to be caused by bacteria.	124	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	Indication specific age restrictions: • Treatment of infections caused by susceptible isolates of the designated microorganisms: 1 month and older • Perioperative prophylaxis: 10 years of age and older	3/22/2024

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

- Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.
- 11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacture participating in the Medicaid Drug Rebate Program (MDRP).
- The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.
- The HCPCS Code effective date represents the date the HCPCS code was established
- Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacture as they are not classified as covered outpatient drugs.
- Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/ncci-medicaid/medicaid-ncci-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Drugs	J0690	Injection, cefazolin sodium, 500 mg	500 mg	1/1/2000	N/A	cefazolin sodium for injection	Indicated for the treatment of the following serious infections when due to susceptible organisms: • Respiratory Tract Infections: Due to <i>S. pneumoniae</i> , <i>Klebsiella</i> species, <i>H. influenzae</i> , <i>S. aureus</i> (penicillin-sensitive and penicillin-resistant), and group A beta-hemolytic streptococci. Injectable benzathine penicillin is considered the drug of choice in treatment and prevention of streptococcal infections, including the prophylaxis of rheumatic fever. Cefazolin is effective in the eradication of streptococci from the nasopharynx; however, data establishing the efficacy of cefazolin in the subsequent prevention of rheumatic fever are not available at present. • Urinary Tract Infections: Due to <i>E. coli</i> , <i>P. mirabilis</i> , <i>Klebsiella</i> species, and some strains of enterobacter	372	1 month	N/A	N/A	Y	Y		1/26/2024
Drugs	J0691	Injection, lefamulin, 1 mg	1 mg	7/1/2020	Xenleta™	lefamulin injection, for intravenous use	Indicated for the treatment of adults with community-acquired bacterial pneumonia (CABP) caused by the following susceptible microorganisms: <i>Streptococcus pneumoniae</i> , <i>Staphylococcus aureus</i> (methicillin-susceptible isolates), <i>Haemophilus influenzae</i> , <i>Legionella pneumophila</i> , <i>Mycoplasma pneumoniae</i> , and <i>Chlamydia pneumoniae</i> . To reduce the development of drug resistant bacteria and maintain the effectiveness of Xenleta and other antibacterial drugs, Xenleta should be used only to treat or prevent infections that are proven or strongly suspected to be caused by bacteria.	2,100	18 years	N/A	N/A	Y	Y		6/17/2020
Drugs	J0692	Injection, cefepime HCl, 500 mg	500 mg	1/1/2002	Maxipime™	cefepime hydrochloride injection for intravenous or intramuscular use	Indicated for the treatment of the following infections caused by susceptible strains of the designated microorganisms: • Moderate to severe pneumonia • Empiric therapy for febrile neutropenic patients • Uncomplicated and complicated urinary tract infections (including pyelonephritis) • Uncomplicated skin and skin structure infections • Complicated intra-abdominal infections (used in combination with metronidazole) in adults	120	2 months	N/A	N/A	Y	Y		8/5/2021
Drugs	J0694	Injection, cefoxitin sodium, 1 gram	1 g	1/1/2000	N/A	cefoxitin for injection	Indicated for the treatment of serious infections caused by susceptible strains of the designated microorganisms in the diseases listed below. • Lower respiratory tract infections: including pneumonia and lung abscess, caused by <i>Streptococcus pneumoniae</i> , other streptococci (excluding enterococci, e.g., <i>Enterococcus faecalis</i> [formerly <i>Streptococcus faecalis</i>]), <i>Staphylococcus aureus</i> (including penicillinase-producing strains), <i>Escherichia coli</i> , <i>Klebsiella</i> species, <i>Haemophilus influenzae</i> , and <i>Bacteroides</i> species. • Urinary tract infections caused by <i>Escherichia coli</i> , <i>Klebsiella</i> species, <i>Proteus mirabilis</i> , <i>Morganella morganii</i> , <i>Proteus vulgaris</i> and <i>Providencia</i> species (including <i>P. rettgeri</i>). • Intra-abdominal infections, including peritonitis and intra-abdominal abscess, caused by <i>Escherichia coli</i> , <i>Klebsiella</i> species, <i>Bacteroides</i> species including <i>Bacteroides fragilis</i> , and <i>Clostridium</i> species. • Gynecological infections: including endometritis, pelvic cellulitis, and pelvic inflammatory disease caused by <i>Escherichia coli</i> , <i>Neisseria gonorrhoeae</i> (including penicillinase-producing strains), <i>Bacteroides</i> species including <i>B. fragilis</i> , <i>Clostridium</i> species, <i>Peptococcus niger</i> , <i>Peptostreptococcus</i> species, and <i>Streptococcus agalactiae</i> . Cefoxitin, like cephalosporins, has no activity against <i>Chlamydia trachomatis</i> . Therefore, when cefoxitin is used in the treatment of patients with pelvic inflammatory	372	3 months	N/A	N/A	Y	Y		9/27/2018

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

•Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.

•11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacture participating in the Medicaid Drug Rebate Program (MDRP).

•The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.

•The HCPCS Code effective date represents the date the HCPCS code was established

•Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacture as they are not classified as covered outpatient drugs.

•Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/ncci-medicaid/medicaid-ncci-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Drugs	J0695	Injection, ceftolozane 50 mg and tazobactam 25 mg	75 mg	1/1/2016	Zerbaxa*	ceftolozane and tazobactam for injection, for intravenous use	<p>Indicated in patients 18 years or older for the treatment of the following infections caused by designated susceptible microorganisms:</p> <ul style="list-style-type: none"> • Complicated intra-abdominal infections (cIAI), used in combination with metronidazole. • Complicated urinary tract infections (cUTI), including pyelonephritis. • Hospital-acquired Bacterial Pneumonia and Ventilator-associated Bacterial Pneumonia (HABP/VABP) <p>Indicated in pediatric patients (birth to less than 18 years old) for the treatment of the following infections caused by designated susceptible microorganisms:</p> <ul style="list-style-type: none"> • Complicated Intra-abdominal Infections (cIAI), used in combination with metronidazole • Complicated Urinary Tract Infections (cUTI), including pyelonephritis <p>To reduce the development of drug-resistant bacteria and maintain the effectiveness of Zerbaxa and other antibacterial drugs, Zerbaxa should be used only to treat or prevent infections that are proven or strongly suspected to be caused by bacteria.</p>	1,680	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	cIAI and cUTI: N/A HABP/VABP: 18 years of age and older	5/9/2022
Drugs	J0696	Injection, ceftriaxone sodium, per 250 mg	250 mg	1/1/2000	Rocephin*	ceftriaxone sodium injection	<p>Indicated for the treatment of the following infections when caused by susceptible organisms:</p> <ul style="list-style-type: none"> • Lower Respiratory Tract Infections: Caused by Streptococcus pneumoniae, Staphylococcus aureus, Haemophilus influenzae, Haemophilus parainfluenzae, Klebsiella pneumoniae, Escherichia coli, Enterobacter aerogenes, Proteus mirabilis or Serratia marcescens. • Acute Bacterial Otitis Media: Caused by Streptococcus pneumoniae, Haemophilus influenzae (including beta-lactamase producing strains) or Moraxella catarrhalis (including beta-lactamase producing strains). • Skin and Skin-Structure Infections: Caused by Staphylococcus aureus, Staphylococcus epidermidis, Streptococcus pyogenes, Viridans group streptococci, Escherichia coli, Enterobacter cloacae, Klebsiella oxytoca, Klebsiella pneumoniae, Proteus mirabilis, Morganella morganii, Pseudomonas aeruginosa, Serratia marcescens, Acinetobacter calcoaceticus, Bacteroides fragilis or Peptostreptococcus species. • Urinary Tract Infections: Caused by Escherichia coli, Proteus mirabilis, Proteus vulgaris, Morganella morganii or Klebsiella pneumoniae. • Uncomplicated Gonorrhoea (cervical/urethral and rectal): Caused by Neisseria gonorrhoeae, including both penicillinase- and non-penicillinase-producing strains, and pharyngeal gonorrhoea caused by non-penicillinase-producing strains of Neisseria gonorrhoeae. • Pelvic Inflammatory Disease: Caused by Neisseria gonorrhoeae. Ceftriaxone sodium, like other cephalosporins, has no activity against Chlamydia trachomatis. Therefore, when cephalosporins are used in the treatment of patients with pelvic inflammatory disease and Chlamydia trachomatis is one of the suspected pathogens, appropriate antichlamydial coverage should be added. • Bacterial Septicemia: Caused by Staphylococcus aureus, Streptococcus pneumoniae, Escherichia coli, Haemophilus influenzae or Klebsiella pneumoniae. 	496	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	See package insert for specific neonate contraindication.	10/4/2018
Drugs	J0697	Injection, sterile cefuroxime sodium, per 750 mg	750 mg	1/1/2000	Zinacef*	cefuroxime for injection	<p>Indicated for the treatment of patients with infections caused by susceptible strains of the designated organisms in the following diseases:</p> <ul style="list-style-type: none"> • Lower Respiratory Tract Infections: including pneumonia, caused by Streptococcus pneumoniae, Haemophilus influenzae (including ampicillin-resistant strains), Klebsiella spp., Staphylococcus aureus (penicillinase- and non-penicillinase-producing strains), Streptococcus pyogenes, and Escherichia coli. • Urinary Tract Infections: caused by Escherichia coli and Klebsiella spp. • Skin and Skin-Structure Infections: caused by Staphylococcus aureus (penicillinase- and non-penicillinase-producing strains), Streptococcus pyogenes, Escherichia coli, Klebsiella spp., and Enterobacter spp. • Septicemia: caused by Staphylococcus aureus (penicillinase- and non-penicillinase-producing strains), Streptococcus pneumoniae, Escherichia coli, Haemophilus influenzae (including ampicillin-resistant strains), and Klebsiella spp. • Meningitis: caused by Streptococcus pneumoniae, Haemophilus influenzae (including ampicillin-resistant strains), Neisseria meningitidis, and Staphylococcus aureus (penicillinase- and non-penicillinase-producing strains). • Gonorrhoeae: Uncomplicated and disseminated gonococcal infections due to Neisseria gonorrhoeae (penicillinase- and non-penicillinase-producing strains) in both males and females. • Bone and Joint Infections: caused by Staphylococcus aureus (penicillinase- and non-penicillinase-producing strains). 	372	3 months	N/A	N/A	Y	Y		10/4/2018

North Carolina Division of Health Benefits Physician Administered Drug Program Catalog

- Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.
- 11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacture participating in the Medicaid Drug Rebate Program (MDRP).
- The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.
- The HCPCS Code effective date represents the date the HCPCS code was established
- Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacture as they are not classified as covered outpatient drugs.
- Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/ncci-medicaid/medicaid-ncci-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Drugs	J0698	Cefotaxime sodium, per gram	1 g	1/1/2000	Claforan®	cefotaxime for injection	Indicated for the treatment of patients with serious infections caused by susceptible strains of the designated microorganisms in the diseases listed below. <ul style="list-style-type: none"> • Lower respiratory tract infections: including pneumonia, caused by Streptococcus pneumoniae (formerly Diplococcus pneumoniae), Streptococcus pyogenes* (Group A streptococci) and other streptococci (excluding enterococci, e.g., Enterococcus faecalis), Staphylococcus aureus (penicillinase and non-penicillinase producing), Escherichia coli, Klebsiella species, Haemophilus influenzae (including ampicillin resistant strains), Haemophilus parainfluenzae, Proteus mirabilis, Serratia marcescens*, Enterobacter species, indole positive Proteus and Pseudomonas species (including P. aeruginosa). • Genitourinary infections: Urinary tract infections caused by Enterococcus species, Staphylococcus epidermidis, Staphylococcus aureus*, (penicillinase and non-penicillinase producing), Citrobacter species, Enterobacter species, Escherichia coli, Klebsiella species, Proteus mirabilis, Proteus vulgaris*, Providencia stuartii, Morganella morganii*, Providencia rettgeri*, Serratia marcescens and Pseudomonas species (including P. aeruginosa). Also, uncomplicated gonorrhoea (cervical/urethral and rectal) caused by Neisseria gonorrhoeae, including penicillinase producing strains. 	372	N/A	N/A	N/A	Y	Y		5/20/2019
Drugs	J0699	Injection, cefiderocol, 10 mg	10 mg	10/1/2021	Fetroja®	cefiderocol for injection, for intravenous use	Indicated in patients 18 years of age or older for the treatment of complicated urinary tract infections (cUTI), including pyelonephritis caused by the following susceptible Gram-negative microorganisms: Escherichia coli, Klebsiella pneumoniae, Proteus mirabilis, Pseudomonas aeruginosa and Enterobacter cloacae complex. Indicated in patients 18 years of age or older for the treatment of hospital-acquired bacterial pneumonia and ventilator-associated bacterial pneumonia, caused by the following susceptible Gram-negative microorganisms: Acinetobacter baumannii complex, Escherichia coli, Enterobacter cloacae complex, Klebsiella pneumoniae, Pseudomonas aeruginosa, and Serratia marcescens.	11,200	18 years	N/A	N/A	Y	Y		9/29/2021
Drugs	J0701	Injection, cefepime hydrochloride (baxter), not therapeutically equivalent to maxipime, 500 mg	500 mg	1/1/2023	N/A	cefepime injection for intravenous use (Baxter)	Indicated in the treatment of the following infections caused by susceptible isolates of the designated microorganisms: pneumonia; empiric therapy for febrile neutropenic patients; uncomplicated and complicated urinary tract infections; uncomplicated skin and skin structure infections; and complicated intra-abdominal infections (used in combination with metronidazole). To reduce the development of drug-resistant bacteria and maintain the effectiveness of Cefepime Injection and other antibacterial drugs, Cefepime Injection should be used only to treat or prevent infections that are proven or strongly suspected to be caused by bacteria.	120	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	Indication specific age restrictions: <ul style="list-style-type: none"> • Complicated intra-abdominal infections: 17 years of age and older • All other indications: 2 months of age and older 	12/19/2022
Drugs	J0702	Injection, betamethasone acetate 3 mg and betamethasone sodium phosphate 3 mg	1 mL	1/1/2000	Celestone® Soluspan®	betamethasone sodium phosphate and betamethasone acetate injectable suspension	When oral therapy is not feasible, the intramuscular use of Celestone Soluspan is indicated as follows: <ul style="list-style-type: none"> • Allergic States: Control of severe or incapacitating allergic conditions intractable to adequate trials of conventional treatment in asthma, atopic dermatitis, contact dermatitis, drug hypersensitivity reactions, perennial or seasonal allergic rhinitis, serum sickness, transfusion reactions. • Dermatologic Diseases: Bullous dermatitis herpetiformis, exfoliative erythroderma, mycosis fungoides, pemphigus, severe erythema multiforme (Stevens-Johnson syndrome). • Endocrine Disorders: Congenital adrenal hyperplasia, hypercalcemia associated with cancer, nonsuppurative thyroiditis. Hydrocortisone or cortisone is the drug of choice in primary or secondary adrenocortical insufficiency. Synthetic analogs may be used in conjunction with mineralocorticoids where applicable; in infancy mineralocorticoid supplementation is of particular importance. • Gastrointestinal Diseases: To tide the patient over a critical period of the disease in regional enteritis and ulcerative colitis. 	155	N/A	N/A	N/A	Y	Y		9/25/2018
Drugs	J0703	Injection, cefepime hydrochloride (b braun), not therapeutically equivalent to maxipime, 500 mg	500 mg	1/1/2023	N/A	cefepime for injection and dextrose injection for intravenous use (B. Braun)	Indicated in the treatment of the following infections caused by susceptible strains of the designated microorganisms: <ul style="list-style-type: none"> • Pneumonia • Empiric therapy for febrile neutropenic patients • Uncomplicated and complicated urinary tract infections • Uncomplicated skin and skin structure infections • Complicated intra-abdominal infections (used in combination with metronidazole) 	120	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	Indication specific age restrictions: <ul style="list-style-type: none"> • Complicated intra-abdominal infections: 17 years of age and older • All other indications: 2 months of age and older 	12/12/2022

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

•Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.

•11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacturer participating in the Medicaid Drug Rebate Program (MDRP).

•The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.

•The HCPCS Code effective date represents the date the HCPCS code was established

•Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacture as they are not classified as covered outpatient drugs.

•Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/ncci-medicaid/medicaid-nci-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Drugs	J0712	Injection, ceftazidime-avibactam, 10 mg	10 mg	1/1/2012	Teflaro*	ceftazidime fosamil for injection, for intravenous use	Indicated for the treatment of the following infection caused by designated susceptible bacteria: • Community-acquired bacterial pneumonia (CABP) in adult and pediatric patients 2 months of age and older • Acute bacterial skin and skin structure infections (ABSSSI) in adult and pediatric patients (at least 34 weeks gestational age and 12 days postnatal age)	1,680	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	Indication specific: CABP: 2 months of age and older ABSSSI: 34 weeks gestational age and 12 days postnatal age and older	10/28/2019
Drugs	J0713	Injection, ceftazidime, per 500 mg	per 500 mg	1/1/2000	Tazicef*	ceftazidime for injection, for intravenous or intramuscular use	Indicated for the treatment of patients with infections caused by susceptible strains of the designated organisms in the following diseases: • Lower Respiratory Tract Infections: including pneumonia, caused by Pseudomonas aeruginosa and other Pseudomonas spp.; Haemophilus influenzae, including ampicillin-resistant strains; Klebsiella spp.; Enterobacter spp.; Proteus mirabilis; Escherichia coli; Serratia spp.; Citrobacter spp.; Streptococcus pneumoniae; and Staphylococcus aureus (methicillin-susceptible strains). • Skin and Skin-Structure Infections: caused by Pseudomonas aeruginosa; Klebsiella spp.; Escherichia coli; Indicated for the treatment of the following infections caused by designated susceptible Gram-negative microorganisms in adult and pediatric patients (at least 31 weeks gestational age): • Complicated Intra-abdominal Infections (cIAI), used in combination with metronidazole • Complicated Urinary Tract Infections (cUTI), including Pyelonephritis • Hospital-acquired Bacterial Pneumonia and Ventilator-associated Bacterial Pneumonia (HABP/VABP)	372	N/A	N/A	N/A	Y	Y		5/21/2019
Drugs	J0714	Injection, ceftazidime and avibactam, 0.5 g/0.125 g	0.625 g	1/1/2016	Avycaz*	ceftazidime and avibactam for injection, for intravenous use	Indicated for the treatment of the following infections caused by designated susceptible Gram-negative microorganisms in adult and pediatric patients (at least 31 weeks gestational age): • Complicated Intra-abdominal Infections (cIAI), used in combination with metronidazole • Complicated Urinary Tract Infections (cUTI), including Pyelonephritis • Hospital-acquired Bacterial Pneumonia and Ventilator-associated Bacterial Pneumonia (HABP/VABP)	168	31 weeks gestational age	N/A	N/A	Y	Y		2/27/2024
Biologics	J0716	Injection, centruroides immune F(ab) ₂ , up to 120 milligrams	up to 120 mg (1 vial)	1/1/2013	Anascorp*	centruroides (scorpion) immune F(ab) ₂ (equine) injection lyophilized for solution, for intravenous use only	Antivenom indicated for treatment of clinical signs of scorpion envenomation.	N/A	N/A	N/A	N/A	Y	Y		4/10/2019
Biologics	J0717	Injection, certolizumab pegol, 1 mg	1 mg	1/1/2014	Cimzia*	certolizumab pegol for injection, for subcutaneous use	Indicated for: • Reducing signs and symptoms of Crohn's disease and maintaining clinical response in adult patients with moderately to severely active disease who have had an inadequate response to conventional therapy. • Treatment of adults with moderately to severely active rheumatoid arthritis. • Treatment of adult patients with active psoriatic arthritis. • Treatment of adults with active ankylosing spondylitis. • Treatment of adults with moderate-to-severe plaque psoriasis who are candidates for systemic therapy or phototherapy. • Treatment of adults with active non-radiographic axial spondyloarthritis who have objective signs of inflammation.	1,200	18 years	N/A	N/A	Y	Y		5/1/2019

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

•Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.

•11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacture participating in the Medicaid Drug Rebate Program (MDRP).

•The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.

•The HCPCS Code effective date represents the date the HCPCS code was established

•Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacture as they are not classified as covered outpatient drugs.

•Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/ncci-medicare/medicaid-ncci-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Drugs	J0720	Injection, chloramphenicol sodium succinate, up to 1 g	up to 1 g	1/1/2000	N/A	chloramphenicol sodium succinate for injection, for intravenous administration	Indicated for: • Acute infections caused by Salmonella typhi. In treatment of typhoid fever some authorities recommend that chloramphenicol be administered at therapeutic levels for 8 to 10 days after the patient has become afebrile to lessen the possibility of relapse. It is not recommended for the routine treatment of the typhoid carrier state. • Serious infections caused by susceptible strains in accordance with the concepts expressed in the package insert: Indicated for: • Prepubertal cryptorchidism not due to anatomic obstruction. In general, HCG is thought to induce testicular descent in situations when descent would have occurred at puberty. HCG thus may help to predict whether or not orchiopexy will be needed in the future. Although, in some cases, descent following	217	N/A	N/A	N/A	Y	Y		10/4/2018
Biologicals	J0725	Injection, chorionic gonadotropin, per 1,000 USP units	1,000 USP units	1/1/2000	Novarel®, Pregnyl®	chorionic gonadotropin for injection	Indicated for: • Prepubertal cryptorchidism not due to anatomic obstruction. In general, HCG is thought to induce testicular descent in situations when descent would have occurred at puberty. HCG thus may help to predict whether or not orchiopexy will be needed in the future. Although, in some cases, descent following	60	4 years	N/A	N/A	Y	Y		6/19/2023
Drugs	J0735	Injection, clonidine hydrochloride, 1 mg	1 mg	1/1/2000	Duraclon®	clonidine hydrochloride injection solution	Indicated in combination with opiates for the treatment of severe pain in cancer patients that is not adequately relieved by opioid analgesics alone. Epidural clonidine is more likely to be effective in patients with neuropathic pain than somatic or visceral pain.	See Comments	N/A	N/A	N/A	Y	Y	Maximum daily and monthly doses are individualized and patient specific.	10/4/2018
Drugs	J0739	Injection, cabotegravir, 1 mg, FDA approved prescription,	1 mg	1/1/2000	Apretude	cabotegravir extended-release injectable suspension,	Indicated in at-risk adults and adolescents weighing at least 35 kg for PrEP to reduce the risk of sexually acquired HIV-1 infection.	1,200	12 years	N/A	N/A	Y	Y		1/4/2024
Drugs	J0740	Injection, cidofovir, 375 mg	375 mg	1/1/2000	Vistide®	cidofovir injection for intravenous infusion	Indicated for the treatment of cytomegalovirus (CMV) retinitis in patients with acquired immunodeficiency syndrome (AIDS).	6	18 years	N/A	N/A	Y	Y		9/27/2018
Drugs	J0741	Injection, cabotegravir and rilpivirine, 2mg/3mg	2 mg/3 mg	10/1/2021	Cabenuva™	cabotegravir extended-release injectable suspension; rilpivirine extended-release injectable suspension, co-packaged for intramuscular use	Indicated as a complete regimen for the treatment of HIV-1 infection in adults and adolescents 12 years of age and older and weighing at least 35 kg to replace the current antiretroviral regimen in those who are virologically suppressed (HIV-1 RNA less than 50 copies per mL) on a stable antiretroviral regimen with no history of treatment failure and with no known or suspected resistance to either cabotegravir or rilpivirine.	600	12 years	N/A	N/A	Y	Y		4/21/2022
Drugs	J0742	Injection, imipenem 4 mg, cilastatin 4 mg and relebactam 2 mg	10 mg	7/1/2020	Recarbrio™	imipenem, cilastatin, and relebactam for injection, for intravenous use	Indicated in patients 18 years of age and older who have limited or no alternative treatment options, for the treatment of the following infections caused by susceptible gram-negative bacteria: • Complicated urinary tract infections, including pyelonephritis (cUTI) • Complicated intra-abdominal infections (cIAI) • Hospital-acquired bacterial pneumonia and ventilator-associated bacterial pneumonia (HABP/VABP) To reduce the development of drug-resistant bacteria and maintain the effectiveness of Recarbrio and other antibacterial drugs, Recarbrio should be used only to treat or prevent infections that are proven or strongly suspected to be caused by bacteria.	7,000	18 years	N/A	N/A	Y	Y		7/28/2020
Drugs	J0743	Injection, cilastatin sodium; imipenem, per 250 mg	250 mg	1/1/2000	Primaxin®	imipenem and cilastatin for injection, for intravenous use	Indicated for the treatment of the following serious infections caused by designated susceptible bacteria: • Lower respiratory tract infections • Urinary tract infections • Intra-abdominal infections • Gynecologic infections • Bacterial septicemia • Bone and joint infections • Skin and skin structure infections • Endocarditis Limitations of Use: • Not indicated in patients with meningitis because safety and efficacy have not been established. • Not recommended in pediatric patients with CNS infections because of the risk of seizures. • Not recommended in pediatric patients weighing less than 30 kg with impaired renal function.	496	N/A	N/A	N/A	Y	Y		9/27/2018
Drugs	J0744	Injection, ciprofloxacin for intravenous infusion, 200 mg	200 mg	1/1/2002	Cipro IV®	ciprofloxacin injection for intravenous use	Indicated in adults (≥ 18 years of age) with the following infections caused by designated, susceptible bacteria and in pediatric patients where indicated: • Skin and skin structure infections • Bone and joint infections • Complicated intra-abdominal infections	186	N/A	N/A	N/A	Y	Y		4/9/2019

North Carolina Division of Health Benefits Physician Administered Drug Program Catalog

•Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.

•11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacturer participating in the Medicaid Drug Rebate Program (MDRP).

•The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.

•The HCPCS Code effective date represents the date the HCPCS code was established

•Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacturer as they are not classified as covered outpatient drugs.

•Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/nci-medicaid/medicaid-nci-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Drugs	J0770	Injection, colistimethate sodium, up to 150 mg	up to 150 mg	1/1/2000	Coly-Mycin® M	colistimethate for injection	Indicated for the treatment of acute or chronic infections due to sensitive strains of certain gram-negative bacilli. Particularly indicated when the infection is caused by sensitive strains of <i>P. aeruginosa</i> . Clinically effective in treatment of infections due to the following gram-negative organisms: <i>Enterobacter</i>	124	N/A	N/A	N/A	Y	Y		6/4/2019
Biologicals	J0775	Injection, collagenase, clostridium histolyticum, 0.01 mg	0.01 mg	1/1/2011	Xiaflex®	collagenase clostridium histolyticum	• Treatment of adult patients with Dupuytren's contracture with a palpable cord. • Treatment of adult men with Peyronie's disease with a palpable plaque and curvature deformity of at least 30 degrees at the start of therapy.	360	18 years	N/A	N/A	Y	Y		6/6/2019
Drugs	J0780	Injection, prochlorperazine, up to 10 mg	up to 10 mg	1/1/2000	N/A	prochlorperazine edisylate injection	Indicated to control severe nausea and vomiting and for the treatment of schizophrenia. Prochlorperazine has not been shown effective in the management of behavioral complications in patients with mental retardation.	124	2 years	N/A	N/A	Y	Y		8/24/2018
Biologicals	J0791	Injection, crizanlizumab-tmca, 5 mg	5 mg	7/1/2020	Adakveo®	crizanlizumab-tmca injection, for intravenous use	Indicated to reduce the frequency of vasoocclusive crises in adults and pediatric patients aged 16 years and older with sickle cell disease.	280	16 years	N/A	N/A	Y	Y		6/17/2020
Drugs	J0801	Injection, corticotropin (acthar gel), up to 40 units	up to 40 units	10/1/2023	H.P. Acthar® Gel	repository corticotropin injection, gel for intramuscular or subcutaneous use	• Indicated as monotherapy for the treatment of infantile spasms in infants and children under 2 years of age. • Indicated for the treatment of exacerbations of multiple sclerosis in adults. • May be used for the following disorders and diseases: rheumatic, collagen, dermatologic, allergic states, ophthalmic, respiratory, and edematous state.	63	N/A	N/A	N/A	Y	Y		9/28/2023
Drugs	J0802	Injection, corticotropin (ani), up to 40 units	up to 40 units	10/1/2023	Purified Cortrophin® Gel	repository corticotropin injection USP	Indicated in the following disorders: 1. Rheumatic disorders: As adjunctive therapy for short-term administration (to tide the patient over an	63	N/A	N/A	N/A	Y	Y		9/28/2023
Drugs	J0834	Injection, cosyntropin, 0.25 mg	0.25 mg	1/1/2010	Cortrosyn™	cosyntropin injection for diagnostic use	Intended for use as a diagnostic agent in the screening of patients presumed to have adrenocortical insufficiency.	3	N/A	N/A	N/A	Y	Y		2/4/2019
Biologicals	J0840	Injection, crotalidae polyvalent immune fab (Ovine), up to 1 gram	up to 1 g (1 vial)	1/1/2012	CroFab®	crotalidae polyvalent immune fab (ovine) lyophilized powder for solution for intravenous injection	Indicated for the management of adult and pediatric patients with North American crotalid envenomation. The term crotalid is used to describe the Crotalinae subfamily (formerly known as Crotalidae) of venomous snakes which includes rattlesnakes, copperheads and cottonmouths/water moccasins.	N/A	N/A	N/A	N/A	Y	N		1/4/2019

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

•Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.

•11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacturer participating in the Medicaid Drug Rebate Program (MDRP).

•The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.

•The HCPCS Code effective date represents the date the HCPCS code was established

•Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacturer as they are not classified as covered outpatient drugs.

•Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/ncci-medicaid/medicaid-ncci-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Biologics	J0841	Injection, crotalidae immune f(ab)2 (equine), 120 mg	120 mg	1/1/2019	Anavip*	crotalidae immune f(ab)2 (equine), lyophilized powder for solution for injection for intravenous use	Indicated for the management of adult and pediatric patients with North American rattlesnake envenomation.	N/A	N/A	N/A	N/A	Y	Y		12/28/2018
Drugs	J0873	Injection, daptomycin (xellia) not therapeutically equivalent to j0878, 1 mg	1 mg	1/1/2024	N/A	daptomycin for injection, for intravenous use (Xellia)	Daptomycin for injection is a lipopeptide antibacterial indicated for the treatment of: • Complicated skin and skin structure infections (cSSSI) in adult and pediatric patients (1 to 17 years of age) and, • Staphylococcus aureus bloodstream infections (bacteremia), in adult patients including those with right-sided infective endocarditis.	26,350	1 year	N/A	N/A	Y	Y		12/22/2023
Drugs	J0874	Injection, daptomycin (baxter), not therapeutically equivalent to j0878, 1 mg	1 mg	10/1/2023	N/A	daptomycin in sodium chloride injection, for intravenous use (Baxter)	Indicated for the treatment of: • Complicated skin and skin structure infections (cSSSI) in adult and pediatric patients (1 to 17 years of age) for whom appropriate dosing can be achieved and, • Staphylococcus aureus bloodstream infections (bacteremia), in adult patients for whom appropriate dosing can be achieved, including those with right-sided infective endocarditis, • Staphylococcus aureus bloodstream infections (bacteremia) in pediatric patients (1 to 17 years of age) for whom appropriate dosing can be achieved. Limitations of Use: • Daptomycin in Sodium Chloride Injection is not indicated for the treatment of pneumonia. • Daptomycin in Sodium Chloride Injection is not indicated for the treatment of left-sided infective endocarditis due to <i>S. aureus</i> . • Daptomycin in Sodium Chloride Injection is not recommended in pediatric patients younger than one year of age due to the risk of potential effects on muscular, neuromuscular, and/or nervous systems (either peripheral and/or central) observed in neonatal dogs. To reduce the development of drug-resistant bacteria and maintain the effectiveness of Daptomycin in Sodium Chloride Injection and other antibacterial drugs, Daptomycin in Sodium Chloride Injection should	31,000	1 year	N/A	N/A	Y	Y		9/28/2023
Drugs	J0875	Injection, dalbavancin, 5 mg	5 mg	1/1/2016	Dalvance*	dalbavancin for injection, for intravenous use	Indicated for the treatment of: - adult patients with acute bacterial skin and skin structure infections (ABSSSI) caused by designated susceptible strains of Gram-positive microorganisms. - pediatric patients with acute bacterial skin and skin structure infections (ABSSSI) caused by designated susceptible strains of Gram-positive microorganisms.	300	N/A	N/A	N/A	Y	Y		8/25/2021
Drugs	J0877	Injection, daptomycin (hospira), not therapeutically equivalent to j0878, 1 mg	1 mg	1/1/2023	N/A	daptomycin for injection, for intravenous use (Hospira)	Indicated for the treatment of: • Complicated skin and skin structure infections (cSSSI) in adult patients • Staphylococcus aureus bloodstream infections (bacteremia), in adult patients including those with right-sided infective endocarditis	26,350	18 years	N/A	N/A	Y	Y		12/12/2022

North Carolina Division of Health Benefits Physician Administered Drug Program Catalog

- Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.
- 11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacture participating in the Medicaid Drug Rebate Program (MDRP).
- The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.
- The HCPCS Code effective date represents the date the HCPCS code was established
- Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacture as they are not classified as covered outpatient drugs.
- Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/ncci-medicare/medicaid-ncci-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Drugs	J0878	Injection, daptomycin, 1 mg	1 mg	1/1/2005	Cubicin®	daptomycin injection, for intravenous use	Indicated for the treatment of: - Complicated skin and skin structure infections (cSSSI) in adult and pediatric patients (1 to 17 years of age). - Staphylococcus aureus bloodstream infections (bacteremia), in adult patients including those with right-sided infective endocarditis. - Indicated for the treatment of Staphylococcus aureus bloodstream infections (bacteremia) in pediatric patients (1 to 17 years of age). Limitations of Use: - Cubicin is not indicated for the treatment of pneumonia. - Cubicin is not indicated for the treatment of left-sided infective endocarditis due to S. aureus. - Cubicin is not recommended in pediatric patients younger than one year of age due to the risk of potential effects on muscular, neuromuscular, and/or nervous systems (either peripheral and/or central) observed in neonatal dogs.	26,040	1 year	N/A	N/A	Y	Y		10/4/2018
Drugs	J0879	Injection, difelikefalin, 0.1 microgram, (for esrd on dialysis)	0.1 mcg	4/1/2002	Korsuva™	difelikefalin injection, for intravenous use	Indicated for the treatment of moderate-to-severe pruritus associated with chronic kidney disease (CKD-aP) in adults undergoing hemodialysis (HD). Limitation of Use: Korsuva has not been studied in patients on peritoneal dialysis and is not recommended for use in this population.	19,500	18 years	N/A	N/A	Y	Y		4/21/2022
Biologics	J0881	Injection, darbepoetin alfa, 1 microgram (non-ESRD use)	1 mcg	1/1/2006	Aranesp®	darbepoetin alfa injection, for intravenous or subcutaneous use (non-ESRD use)	Indicated for the treatment of anemia due to: • Chronic Kidney Disease (CKD) in patients on dialysis and patient not on dialysis. • The effects of concomitant myelosuppressive chemotherapy, and upon initiation, there is a minimum of two additional months of planned chemotherapy. Limitations of Use: Aranesp has not been shown to improve quality of life, fatigue, or patient well-being. Aranesp is not indicated for use: • In patients with cancer receiving hormonal agents, biologic products, or radiotherapy, unless also receiving concomitant myelosuppressive chemotherapy. • In patients with cancer receiving myelosuppressive chemotherapy when the anticipated outcome is cure. • In patients with cancer receiving myelosuppressive chemotherapy in whom the anemia can be managed by transfusion. • As a substitute for RBC transfusions in patients who require immediate correction of anemia.	1,575	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	Indication specific age restrictions: • CKD: None • Cancer: 18 years of age and older	4/10/2019
Biologics	J0882	Injection, darbepoetin alfa, 1 microgram (for ESRD on dialysis)	1 mcg	1/1/2006	Aranesp®	darbepoetin alfa injection, for intravenous or subcutaneous use (ESRD use on dialysis)	Indicated for the treatment of anemia due to: • Chronic Kidney Disease (CKD) in patients on dialysis and patients not on dialysis. • The effects of concomitant myelosuppressive chemotherapy, and upon initiation, there is a minimum of two additional months of planned chemotherapy. Limitations of Use: Aranesp has not been shown to improve quality of life, fatigue, or patient well-being. Aranesp is not indicated for use: • In patients with cancer receiving hormonal agents, biologic products, or radiotherapy, unless also receiving concomitant myelosuppressive	315	N/A	N/A	N/A	Y	Y		4/10/2019
Biologics	J0885	Injection, epoetin alfa, (for non-ESRD use), 1000 units	1,000 units	1/1/2006	Epogen®, Procrit®	epoetin alfa for injection, for intravenous or subcutaneous use (for non ESRD use)	•Indicated for treatment of anemia due to - Chronic Kidney Disease (CKD) in patients on dialysis and not on dialysis. - Zidovudine in patients with HIV-infection. - The effects of concomitant myelosuppressive chemotherapy, and upon initiation, there is a minimum of two additional months of planned chemotherapy. • Reduction of allogeneic RBC transfusions in patients undergoing elective, noncardiac, nonvascular surgery. Limitations of Use: Epoetin alfa has not been shown to improve quality of life, fatigue, or patient well-being.	630	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	Indication specific age restrictions: • CKD not on dialysis: 1 month of age and older • Anemia due to concomitant myelosuppressive chemotherapy: 5 years of age and older • Zidovudine-treated, anemia, patients with HIV infection: 8 months and older	1/12/2022

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

•Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.

•11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacture participating in the Medicaid Drug Rebate Program (MDRP).

•The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.

•The HCPCS Code effective date represents the date the HCPCS code was established

•Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacture as they are not classified as covered outpatient drugs.

•Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/ncci-medicare/medicaid-ncci-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Biologics	J0887	Injection, epoetin beta, 1 microgram, (for ESRD on dialysis)	1 mcg	1/1/2015	Mircera*	methoxy polyethylene glycol-epoetin beta injection, for intravenous or subcutaneous use (for ESRD on dialysis)	Indicated for the treatment of anemia associated with chronic kidney disease (CKD) in: • adult patients on dialysis and adult patients not on dialysis. • pediatric patients 5 to 17 years of age on hemodialysis who are converting from another ESA after their hemoglobin level was stabilized with an ESA. Limitations of Use: Indicated for the treatment of anemia associated with chronic kidney disease (CKD) in: • Adult patients on dialysis and adult patients not on dialysis. • Pediatric patients 5 to 17 years of age on hemodialysis who are converting from another ESA after their hemoglobin level was stabilized with an ESA.	720	5 years	N/A	N/A	Y	Y		10/10/2018
Biologics	J0888	Injection, epoetin beta, 1 microgram, (for non-ESRD use)	1 mcg	1/1/2015	Mircera*	methoxy polyethylene glycol-epoetin beta injection, for intravenous or subcutaneous use (for non-ESRD use)	Indicated for the treatment of anemia associated with chronic kidney disease (CKD) in: • Adult patients on dialysis and adult patients not on dialysis. • Pediatric patients 5 to 17 years of age on hemodialysis who are converting from another ESA after their hemoglobin level was stabilized with an ESA. Limitations of Use: Indicated for treatment of adult patients with myelodysplastic syndromes (MDS) including previously treated and untreated, de novo and secondary MDS of all French-American-British subtypes (refractory	720	18 years	N/A	N/A	Y	Y		9/14/2021
Drugs	J0893	Injection, decitabine (sun pharma), not therapeutically	1 mg	1/1/2023	N/A	decitabine for injection, for intravenous use (Sun	Indicated for treatment of adult patients with myelodysplastic syndromes (MDS) including previously treated and untreated, de novo and secondary MDS of all French-American-British subtypes (refractory	450	18 years	N/A	N/A	Y	Y		12/6/2022
Drugs	J0894	Injection, decitabine, 1 mg	1 mg	1/1/2007	N/A	decitabine for injection, for intravenous infusion	Indicated for treatment of patients with myelodysplastic syndromes (MDS) including previously treated and untreated, de novo and secondary MDS of all French-American-British subtypes (refractory anemia, refractory anemia with ringed sideroblasts, refractory anemia with excess blasts, refractory anemia with excess blasts in transformation, and chronic myelomonocytic leukemia) and intermediate-1, intermediate-2, and high-risk International Prognostic Scoring System groups.	450	18 years	N/A	N/A	Y	Y		10/4/2018
Drugs	J0895	Injection, deferoxamine mesylate, 500 mg	500 mg	1/1/2000	Desferal*	deferoxamine mesylate for injection	Indicated for the treatment of acute iron intoxication and of chronic iron overload due to transfusion-dependent anemias.	372	3 years	N/A	N/A	Y	Y		10/4/2018
Biologics	J0896	Injection, luspatercept-aamt, 0.25 mg	0.25 mg	7/1/2020	Reblozyl*	luspatercept-aamt for injection, for subcutaneous	Indicated for the treatment of: • anemia in adult patients with beta thalassemia who require regular red blood cell (RBC) transfusions.	2,000	18 years	N/A	N/A	Y	Y		9/28/2023
Biologics	J0897	Injection, denosumab, 1 mg (Xgeva, Prolia)	1 mg	1/1/2012	Prolia*, Xgeva*	denosumab injection, for subcutaneous use	 Prolia Indicated for: • The treatment in postmenopausal women with osteoporosis at high risk for fracture • The treatment to increase bone mass in men with osteoporosis at high risk for fracture • The treatment to increase bone mass in men at high risk for fracture receiving androgen deprivation therapy for nonmetastatic prostate cancer • The treatment to increase bone mass in women at high risk for fracture receiving adjuvant aromatase inhibitor therapy for breast cancer. • The treatment of glucocorticoid-induced osteoporosis in men and women at high risk for fracture. Xgeva Indicated for: • The prevention of skeletal-related events in patients with multiple myeloma and in patients with bone metastases from solid tumors • The treatment of adults and skeletally mature adolescents with giant cell tumor of bone that is unresectable or where surgical resection is likely to result in severe morbidity • The treatment of hypercalcemia of malignancy refractory to bisphosphonate therapy	360	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	Product/indication specific age restrictions: • Prolia: 18 years of age and older • Xgeva: Indication specific. o Giant cell tumor of bone: Only use in skeletally mature adolescents. o All other indications: 18 years of age and older	10/31/2018

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

- Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.
- 11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacturer participating in the Medicaid Drug Rebate Program (MDRP).
- The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.
- The HCPCS Code effective date represents the date the HCPCS code was established
- Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacture as they are not classified as covered outpatient drugs.
- Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/ncci-medicaid/medicaid-ncci-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Drugs	J1000	Injection, depo-estradiol cypionate, up to 5 mg	up to 5 mg	1/1/2000	Depo®-Estradiol	estradiol cypionate injection	Indicated in the treatment of hypoestrogenism caused by hypogonadism and moderate to severe vasomotor symptoms associated with the menopause.	2	18 years	N/A	Females Only	Y	Y		10/4/2018
Drugs	J1010	Injection, methylprednisolone acetate, 1 mg	1 mg	4/1/2024	Depo-Medrol®	methylprednisolone acetate injection, suspension, USP	Indicated as follows when the oral route is not feasible: Intramuscular Administration • Allergic States: Control of severe or incapacitating allergic conditions intractable to adequate trials of conventional treatment in asthma, atopic dermatitis, contact dermatitis, drug hypersensitivity reactions, serum sickness, transfusion reactions. • Dermatologic Diseases: Bullous dermatitis herpetiformis, exfoliative dermatitis, mycosis fungoides, pemphigus, severe erythema multiforme (Stevens-Johnson syndrome). • Endocrine Disorders: Primary or secondary adrenocortical insufficiency (hydrocortisone or cortisone is the drug of choice; synthetic analogs may be used in conjunction with mineralocorticoids where	800	N/A	N/A	N/A	Y	Y		3/22/2024
Drugs	J1050	Injection, medroxyprogesterone acetate, 1 mg	1 mg	1/1/2013	Depo-Provera®	medroxyprogesterone acetate, injectable suspension	Indicated for prevention of pregnancy in females and adjunctive therapy and palliative treatment of inoperable, recurrent, and metastatic endometrial or renal carcinoma.	5,000	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	Indication specific age restrictions: • Endometrial and renal carcinoma: 18 years and older	10/26/2018
Drugs	J1071	Injection, testosterone cypionate, 1 mg	1 mg	1/1/2015	Depo®-Testosterone	testosterone cypionate injection, USP	Indicated for replacement therapy in the male in conditions associated with symptoms of deficiency or absence of endogenous testosterone. 1. Primary hypogonadism (congenital or acquired)-testicular failure due to cryptorchidism, bilateral	1,200	12 years	N/A	Males Only	Y	Y		4/10/2019
Drugs	J1095	Injection, dexamethasone 9 percent, intraocular, 1 microgram	1 mcg	1/1/2019	Dexycu™	dexamethasone intraocular suspension 9%, for intraocular administration	Indicated for the treatment of postoperative inflammation.	1,034	18 years	N/A	N/A	Y	Y		3/26/2019
Drugs	J1096	Dexamethasone, lacrimal ophthalmic insert, 0.1 mg	0.1 mg	10/1/2019	Dextenza®	dexamethasone ophthalmic insert 0.4 mg, for intracanalicular use	Indicated for: • The treatment of ocular inflammation and pain following ophthalmic surgery. • The treatment of ocular itching associated with allergic conjunctivitis.	8	18 years	N/A	N/A	Y	Y		11/17/2021
Drugs	J1097	phenylephrine 10.16 mg/ml and ketorolac 2.88 mg/ml ophthalmic irrigation solution, 1 ml	1 mL	10/1/2019	Omidria®	phenylephrine and ketorolac intraocular solution, 1% /0.3%, for addition to ocular irrigating solution	Indicated for maintaining pupil size by preventing intraoperative miosis and reducing postoperative ocular pain.	8	N/A	N/A	N/A	Y	Y		9/27/2019
Drugs	J1100	Injection, dexamethasone sodium phosphate, 1 mg	1 mg	1/1/2000	N/A	dexamethasone sodium phosphate injection	Intravenous or Intramuscular Administration: When oral therapy is not feasible and the strength, dosage form, and route of administration of the drug reasonably lend the preparation to the treatment of the condition, those products labeled for intravenous or intramuscular use are indicated as follows:	310	N/A	N/A	N/A	Y	Y		10/4/2018
Drugs	J1105	Dexmedetomidine, oral, 1 mcg	1 mcg	1/1/2024	Igalmi™	dexmedetomidine sublingual film, for sublingual or buccal use	Indicated in adults for the acute treatment of agitation associated with schizophrenia or bipolar I or II disorder. Limitations of Use: The safety and effectiveness of Igalmi has not been established beyond 24 hours from the first dose.	1,800	18 years	N/A	N/A	Y	Y		12/22/2023

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

•Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.

•11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacturer participating in the Medicaid Drug Rebate Program (MDRP).

•The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.

•The HCPCS Code effective date represents the date the HCPCS code was established

•Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacturer as they are not classified as covered outpatient drugs.

•Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/nccli-medicaid/medicaid-nccli-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Drugs	J1110	Injection, dihydroergotamine mesylate, per 1 mg	1 mg	1/1/2000	DHE 45*	dihydroergotamine mesylate injection	Indicated for the acute treatment of migraine headaches with or without aura and the acute treatment of cluster headache episodes.	30	18 years	N/A	N/A	Y	Y		10/10/2018
Drugs	J1120	Injection, acetazolamide sodium, up to 500 mg	up to 500 mg	1/1/2000	Diamox®	acetazolamide sodium injection, powder, lyophilized, for solution	Indicated for the adjunctive treatment of: • Edema due to congestive heart failure • Drug-induced edema • Centrencephalic epilepsies (petit mal, unlocalized seizures)	62	18 years	N/A	N/A	Y	Y		10/31/2018
Drugs	J1160	Injection, digoxin, up to 0.5 mg	up to 0.5 mg	1/1/2000	Lanoxin®	digoxin injection, for intravenous or intramuscular use	Indicated for: • Treatment of mild to moderate heart failure in adults. • Increasing myocardial contractility in pediatric patients with heart failure. (Indication added to the portal 10/4/2018) • Control of resting ventricular rate in adults with chronic atrial fibrillation.	35	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	Indication specific age restrictions: • Mild to moderate heart failure and control of resting ventricular rate in chronic atrial fibrillation: 18 years of age and older • Increasing myocardial	10/10/2018
Drugs	J1165	Injection, phenytoin sodium, per 50 mg	per 50 mg	1/1/2000	N/A	phenytoin sodium injection, for intravenous or intramuscular use	Indicated for the treatment of generalized tonic-clonic status epilepticus and prevention and treatment of seizures occurring during neurosurgery. Intravenous phenytoin can also be substituted, as short-term use, for oral phenytoin. Parenteral phenytoin should be used only when oral phenytoin administration is not possible.	288	N/A	N/A	N/A	Y	Y		6/8/2019
Drugs	J1170	Injection, hydromorphone, up to 4 mg	up to 4 mg	1/1/2000	Dilaudid®	hydromorphone hydrochloride for intravenous, intramuscular, and subcutaneous use	Indicated for the management of pain severe enough to require an opioid analgesic and for which alternate treatments are inadequate. Limitations of Use: Because of the risks of addiction, abuse, and misuse with opioids, even at recommended doses, reserve hydromorphone injection for use in patients for whom alternative treatment options [e.g., nonopioid analgesics or opioid combination products]: • Have not been tolerated, or are not expected to be tolerated • Have not provided adequate analgesia, or are not expected to provide adequate analgesia	186	18 years	N/A	N/A	Y	Y		10/26/2018
Drugs	J1190	Injection, dexrazoxane hydrochloride, per 250 mg	250 mg	1/1/2000	Totect®, Zinecard®	dexrazoxane for injection	Zinecard: Indicated for reducing the incidence and severity of cardiomyopathy associated with doxorubicin administration in women with metastatic breast cancer who have received a cumulative doxorubicin dose of 300 mg/m ² and who will continue to receive doxorubicin therapy to maintain tumor control. Do not use with doxorubicin initiation. Totect: Indicated for the treatment of extravasation resulting from IV anthracycline chemotherapy. • Reducing the incidence and severity of cardiomyopathy associated with doxorubicin administration in women with metastatic breast cancer who have received a cumulative doxorubicin dose of 300 mg/m ² and who will continue to receive doxorubicin therapy to maintain tumor control. Do not use Totect with doxorubicin initiation.	20	18 years	N/A	Zinecard: Females Only Totect: Extravasation: N/A Cardiomyopathy: Females only	Y	Y		12/28/2020
Drugs	J1200	Injection, diphenhydramine HCl, up to 50 mg	50 mg	1/1/2000	N/A	diphenhydramine hydrochloride injection	Diphenhydramine in the injectable form is effective in adults and pediatric patients, other than premature infants and neonates, for the following conditions when diphenhydramine in the oral form is impractical: • Antihistaminic: For amelioration of allergic reactions to blood or plasma, in anaphylaxis as an adjunct to epinephrine and other standard measures after the acute symptoms have been controlled, and for other uncomplicated allergic conditions of the immediate type when oral therapy is impossible or contraindicated. • Motion Sickness: For active treatment of motion sickness. • Antiparkinsonism: For use in parkinsonism, when oral therapy is impossible or contraindicated, as follows: parkinsonism in the elderly who are unable to tolerate more potent agents; mild cases of parkinsonism in other age groups, and in other cases of parkinsonism in combination with centrally acting	248	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	Contraindicated in newborns or premature infants.	10/4/2018
Drugs	J1202	Miglustat, oral, 65 mg	65 mg	4/1/2024	Opfolda™	miglustat capsules, for oral use	Miglustat capsule is indicated, in combination with Pombiliti, for the treatment of adult patients with late-onset Pompe disease (lysosomal acid alpha-glucosidase [GAA] deficiency) weighing ≥40 kg and who are not improving on their current enzyme replacement therapy (ERT).	12	18 years	N/A	N/A	Y	Y		3/22/2024
Biologicals	J1203	Injection, cipaglucosidase alfa-atga, 5 mg	5 mg	4/1/2024	Pombiliti™	cipaglucosidase alfa-atga for injection, for intravenous use	Indicated, in combination with Opfolda, an enzyme stabilizer, for the treatment of adult patients with late-onset Pompe disease (lysosomal acid alpha-glucosidase [GAA] deficiency) weighing ≥40 kg and who are	1,701	18 years	N/A	N/A	Y	Y		3/22/2024

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

•Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.

•11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacturer participating in the Medicaid Drug Rebate Program (MDRP).

•The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.

•The HCPCS Code effective date represents the date the HCPCS code was established

•Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacturer as they are not classified as covered outpatient drugs.

•Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/nccli-medicaid/medicaid-nccli-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Drugs	J1205	Injection, chlorothiazide sodium, per 500 mg	500 mg	1/1/2000	N/A	chlorothiazide sodium for injection	Indicated as adjunctive therapy in edema associated with congestive heart failure, hepatic cirrhosis, and corticosteroid and estrogen therapy.	100	18 years	N/A	N/A	Y	Y		9/27/2018
Drugs	J1212	Injection, DMSO, dimethyl sulfoxide, 50%, 50 mL	50 mL	1/1/2000	RIMSO-50®	dimethyl sulfoxide (DMSO) irrigation	Indicated for symptomatic relief of patients with interstitial cystitis.	3	N/A	N/A	N/A	Y	Y		10/4/2018
Drugs	J1230	Injection, methadone HCl, up to 10 mg	up to 10 mg	1/1/2000	N/A	methadone hydrochloride injection	Indicated for: • The management of pain severe enough to require an opioid analgesic and for which alternative	93	18 years	N/A	N/A	Y	Y		10/26/2018
Drugs	J1240	Injection, dimenhydrinate, up to 50 mg	up to 50 mg	1/1/2000	N/A	dimenhydrinate injection	Indicated for prevention and treatment of nausea, vomiting and vertigo of motion sickness.	372	N/A	N/A	N/A	Y	Y		6/10/2019
Drugs	J1245	Injection, dipyridamole, per 10 mg	per 10 mg	1/1/2000	N/A	dipyridamole injection	As an alternative to exercise in thallium myocardial perfusion imaging for the evaluation of coronary artery disease in patients who cannot exercise adequately.	6	18 years	N/A	N/A	Y	Y		6/10/2019
Drugs	J1250	Injection, dobutamine hydrochloride, per 250 mg	250 mg	1/1/2000	N/A	dobutamine injection	Indicated: • When parenteral therapy is necessary for inotropic support in the short-term treatment of adults with	930	18 years	N/A	N/A	Y	Y		10/4/2018
Drugs	J1265	Injection, dopamine hydrochloride, 40 mg	40 mg	1/1/2006	N/A	dopamine hydrochloride	Indicated for the correction of hemodynamic imbalances present in the shock syndrome due to myocardial infarction, trauma, endotoxic septicemia, open-heart surgery, renal failure, and chronic cardiac	6,355	18 years	N/A	N/A	Y	Y		10/4/2018
Drugs	J1270	Injection, doxercalciferol, 1 mcg	1 mcg	1/1/2002	Hectorol®	doxercalciferol injection	Indicated for the treatment of secondary hyperparathyroidism in adult patients with chronic kidney disease on dialysis.	90	18 years	N/A	N/A	Y	Y		10/4/2018
Drugs	J1290	Injection, ecallantide, 1 mg	1 mg	1/1/2011	Kalbitor®	ecallantide injection for subcutaneous use	Indicated for treatment of acute attacks of hereditary angioedema in patients 12 years of age and older.	120	12 years	N/A	N/A	Y	Y		10/10/2018
Biologicals	J1300	Injection, eculizumab, 10 mg	10 mg	1/1/2008	Soliris®	eculizumab injection, for intravenous use	Indicated for: • Treatment of patients with paroxysmal nocturnal hemoglobinuria (PNH) to reduce hemolysis. • Treatment of patients with atypical hemolytic uremic syndrome (aHUS) to inhibit complement-mediated	480	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	Indication specific age restrictions: • PNH: 18 years of age and	7/26/2019
Drugs	J1301	Injection, edaravone, 1 mg	1 mg	1/1/2019	Radicava®	edaravone injection, for intravenous use	Indicated for the treatment of amyotrophic lateral sclerosis (ALS).	1,020	18 years	N/A	N/A	Y	Y		10/10/2018
Biologicals	J1302	Injection, sutimlimab-jome, 10 mg	10 mg	10/1/2022	Enjaymo™	sutimlimab-jome injection, for intravenous use	Indicated for the treatment of hemolysis in adults with cold agglutinin disease (CAD).	2,310	18 years	N/A	N/A	Y	Y		2/23/2023
Biologicals	J1303	Injection, ravulizumab-cwvz, 10 mg	10 mg	10/1/2019	Ultomiris™	ravulizumab-cwvz injection, for intravenous use	Indicated for: - the treatment of adult and pediatric patients one month of age and older with paroxysmal nocturnal hemoglobinuria (PNH). - the treatment of adults and pediatric patients one month of age and older with atypical hemolytic uremic syndrome (aHUS) to inhibit complement-mediated thrombotic microangiopathy (TMA). Limitations of Use: Ultomiris is not indicated for the treatment of patients with Shiga toxin E. coli related hemolytic uremic syndrome (STEC-HUS). - the treatment of adult patients with generalized myasthenia gravis (gMG) who are anti-acetylcholine receptor (AChR) antibody-positive.	660	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	PNH and aHUS: 1 month of age and older gMG and NMOSD: 18 years of age and older	5/3/2024
Drugs	J1304	Injection, tofersen, 1 mg	1 mg	1/1/2024	Qalsody™	tofersen injection, for intrathecal use	Indicated for the treatment of amyotrophic lateral sclerosis (ALS) in adults who have a mutation in the superoxide dismutase 1 (SOD1) gene.	300	18 years	N/A	N/A	Y	Y		12/22/2023
Biologicals	J1305	Injection, evinacumab-dgnb, 5mg	5 mg	10/1/2021	Evkeeza™	evinacumab-dgnb injection, for intravenous use	Indicated as an adjunct to other low-density lipoprotein-cholesterol (LDL-C) lowering therapies for the treatment of adult and pediatric patients, aged 5 years and older, with homozygous familial	894	5 years	N/A	N/A	Y	Y		4/25/2023

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

•Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.

•11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacturer participating in the Medicaid Drug Rebate Program (MDRP).

•The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.

•The HCPCS Code effective date represents the date the HCPCS code was established

•Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacturer as they are not classified as covered outpatient drugs.

•Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/ncci-medicaid/medicaid-ncci-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Drugs	J1306	Injection, inclisiran, 1 mg	1 mg	1/1/2000	Leqvio*	inclisiran injection, for subcutaneous use	Indicated as an adjunct to diet and maximally tolerated statin therapy for the treatment of adults with heterozygous familial hypercholesterolemia (HeFH) or clinical atherosclerotic cardiovascular disease (ASCVD), who require additional lowering of low-density lipoprotein cholesterol (LDL-C).	284	18 years	N/A	N/A	Y	Y		9/13/2023
Biologicals	J1322	Injection, elosulfase alfa, 1 mg	1 mg	1/1/2015	Vimizim*	elosulfase alfa injection, for intravenous use	Indicated for patients with Mucopolysaccharidosis type IVA (MPS IVA; Morquio A syndrome).	1,400	5 years	N/A	N/A	Y	Y		6/8/2019
Biologicals	J1323	Injection, elranatamab-bcmm, 1 mg	1 mg	4/1/2024	Elexfio™	elranatamab-bcmm injection, for subcutaneous use	Indicated for the treatment of adult patients with relapsed or refractory multiple myeloma who have received at least four prior lines of therapy including a proteasome inhibitor, an immunomodulatory agent, and an anti-CD38 monoclonal antibody.	380	18 years	N/A	N/A	Y	Y		4/12/2024
Drugs	J1325	Injection, epoprostenol, 0.5 mg	0.5 mg	1/1/2000	Folan®, Veletri®	epoprostenol for injection, for intravenous use	Indicated for the treatment of pulmonary arterial hypertension (PAH) (WHO Group I) to improve exercise capacity. Studies establishing effectiveness included predominantly (97%) patients with NYHA Functional Class III-IV symptoms and etiologies of idiopathic or heritable PAH (49%) or PAH associated with connective tissue diseases (51%).	248	18 years	N/A	N/A	Y	Y		6/4/2019
Drugs	J1335	Injection, ertapenem sodium, 500 mg	500 mg	1/1/2004	Invanz*	ertapenem injection for intravenous or intramuscular use	Indicated in adult patients and pediatric patients (3 months of age and older) for the treatment of the following moderate to severe infections caused by susceptible bacteria: • Complicated intra-abdominal infections. • Complicated skin and skin structure infections, including diabetic foot infections without osteomyelitis. • Community-acquired pneumonia. • Complicated urinary tract infections including pyelonephritis. • Acute pelvic infections including postpartum endometritis, septic abortion and post surgical	28	3 months	N/A	N/A	Y	Y		10/10/2018
Drugs	J1364	Injection, erythromycin lactobionate, per 500 mg	500 mg	1/1/2000	Erythrocin™	erythromycin lactobionate for injection	Indicated in the treatment of infections caused by susceptible strains of the designated organisms in the diseases listed below when oral administration is not possible or when the severity of the infection requires immediate high serum levels of erythromycin. Intravenous therapy should be replaced by oral	248	N/A	N/A	N/A	Y	Y		10/10/2018
Drugs	J1380	Injection, estradiol valerate, up to 10 mg	up to 10 mg	1/1/2000	Delestrogen®	estradiol valerate injection	Indicated in the treatment of: • Moderate-to-severe vasomotor symptoms associated with the menopause • Hypoestrogenism caused by hypogonadism, castration or primary ovarian failure	20	18 years	N/A	N/A	Y	Y		6/10/2019
Drugs	J1410	Injection, estrogens, conjugated, per 25 mg	25 mg	1/1/2000	Premarin® IV	conjugated estrogens for injection for intravenous and intramuscular use	Indicated in the treatment of abnormal uterine bleeding caused by hormonal imbalance in the absence of organic pathology. Indicated for short-term use only, to provide a rapid and temporary increase in estrogen levels.	62	N/A	N/A	Females Only	Y	Y		10/10/2018
Drugs	J1437	Injection, ferric derisomaltose, 10 mg	10 mg	10/1/2020	MonoFerric™	ferric derisomaltose injection, for intravenous use	Indicated for the treatment of iron deficiency anemia in adult patients: • who have intolerance to oral iron or have had unsatisfactory response to oral iron. • who have non-hemodialysis dependent chronic kidney disease.	100	18 years	N/A	N/A	Y	Y		12/28/2020

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

•Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.

•11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacture participating in the Medicaid Drug Rebate Program (MDRP).

•The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.

•The HCPCS Code effective date represents the date the HCPCS code was established

•Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacture as they are not classified as covered outpatient drugs.

•Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/nccli-medicare/medicaid-nccli-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Drugs	J1439	Injection, ferric carboxymaltose, 1 mg	1 mg	1/1/2015	Injectafer®	ferric carboxymaltose injection, for intravenous use	Indicated for the treatment of iron deficiency anemia (IDA) in adult patients: - Who have intolerance to oral iron or have had unsatisfactory response to oral iron. - Who have non-dialysis dependent chronic kidney disease. - With heart failure and New York Heart Association class II/III to improve exercise capacity. Indicated for the treatment of iron deficiency anemia in pediatric patients 1 year of age to 17 years of age who have either intolerance to oral iron or an unsatisfactory response to oral iron.	1,500	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	Indication specific age restrictions: • IDA in patients who have either intolerance to oral iron or an unsatisfactory response to oral iron: 1 year of age and older • IDA in patients who have non dialysis dependent chronic kidney disease, iron deficiency in adult patients with heart failure and New York Heart Association class II/III to improve exercise capacity: 18 years of age and older	6/19/2023
Biologics	J1440	Fecal microbiota, live - jslm, 1 ml	1 mL	7/1/2023	Rebyota™	fecal microbiota, live - jslm suspension, for rectal use	Indicated for the prevention of recurrence of Clostridioides difficile infection (CDI) in individuals 18 years of age and older, following antibiotic treatment for recurrent CDI. Limitation of Use: Rebyota is not indicated for treatment of CDI.	150	18 years	N/A	N/A	Y	Y		6/22/2023
Biologics	J1442	Injection, filgrastim (G-CSF), excludes biosimilars, 1 microgram	1 mcg	1/1/2016	Neupogen®	filgrastim injection, for subcutaneous or intravenous use	Indicated to: • Decrease the incidence of infection, as manifested by febrile neutropenia, in patients with nonmyeloid malignancies receiving myelosuppressive anti-cancer drugs associated with a significant incidence of severe neutropenia with fever. • Reduce the time to neutrophil recovery and the duration of fever, following induction or consolidation chemotherapy treatment of patients with acute myeloid leukemia (AML). Indicated for the replacement of iron to maintain hemoglobin in adult patients with hemodialysis-dependent chronic kidney disease (HDD-CKD). Limitations of Use: • Triferic is not intended for use in patients receiving peritoneal dialysis.	59,520	N/A	N/A	N/A	Y	Y		6/6/2019
Drugs	J1443	Injection, ferric pyrophosphate citrate solution (Triferic), 0.1 mg of iron	0.1 mg of iron	10/1/2021	Triferic®	ferric pyrophosphate citrate solution, for hemodialysis use, and powder for solution	Indicated for the replacement of iron to maintain hemoglobin in adult patients with hemodialysis-dependent chronic kidney disease (HDD-CKD). Limitations of Use: • Triferic is not intended for use in patients receiving peritoneal dialysis.	38,080	18 years	N/A	N/A	Y	Y		9/29/2021
Drugs	J1444	Injection, ferric pyrophosphate citrate powder, 0.1 mg of iron (This code would be used with the "JE" modifier, when administered via dialysate.)	0.1 mg	7/1/2019	Triferic®	ferric pyrophosphate citrate powder packet for hemodialysis use	Indicated for the replacement of iron to maintain hemoglobin in adult patients with hemodialysis-dependent chronic kidney disease (HDD-CKD). Limitations of Use: • Triferic is not intended for use in patients receiving peritoneal dialysis.	38,080	18 years	N/A	N/A	Y	Y		7/26/2019
Biologics	J1447	Injection, tbo-filgrastim, 1 microgram	1 mcg	1/1/2016	Granix®	tbo-filgrastim injection, for subcutaneous use	Indicated in adult and pediatric patients 1 month and older for reduction in the duration of severe neutropenia in patients with non-myeloid malignancies receiving myelosuppressive anti-cancer drugs associated with a clinically significant incidence of febrile neutropenia.	10,920	1 month	N/A	N/A	Y	Y		5/20/2019
Drugs	J1448	Injection, trilaciclib, 1mg	1 mg	10/1/2021	Cosela*	trilaciclib for injection, for intravenous use	Indicated to decrease the incidence of chemotherapy-induced myelosuppression in adult patients when administered prior to a platinum/etoposide-containing regimen or topotecan-containing regimen for extensive-stage small cell lung cancer.	9,000	18 years	N/A	N/A	Y	Y	12/2023: NC Suggested Max Monthly updated from 1,200 units to 9,000 units effective 5/1/2023 at DHB request.	12/1/2023
Biologics	J1449	Injection, eflapegrastim-xnst, 0.1 mg	0.1 mg	4/1/2023	Rolvedon™	eflapegrastim-xnst injection, for subcutaneous use	Indicated to decrease the incidence of infection, as manifested by febrile neutropenia, in adult patients with nonmyeloid malignancies receiving myelosuppressive anti-cancer drugs associated with clinically significant incidence of febrile neutropenia. Limitations of Use: Rolvedon is not indicated for the mobilization of peripheral blood progenitor cells for hematopoietic stem cell transplantation.	396	18 years	N/A	N/A	Y	Y		3/16/2023
Drugs	J1453	Injection, fosaprepitant, 1 mg	1 mg	1/1/2009	Emend®	fosaprepitant for injection, for intravenous use	Indicated in adults and pediatric patients 6 months of age and older, in combination with other antiemetic agents, for the prevention of: • acute and delayed nausea and vomiting associated with initial and repeat courses of highly emetogenic cancer chemotherapy (HEC) including high-dose cisplatin. • delayed nausea and vomiting associated with initial and repeat courses of moderately emetogenic cancer chemotherapy (MEC). Limitations of Use: Emend has not been studied for treatment of established nausea and vomiting.	750	6 months	N/A	N/A	Y	Y	9/2023: NC Suggested Max Monthly Units updated from 600 units to 750 units effective 1/1/2023 at DHB request	9/28/2023
Drugs	J1454	Injection, fosnetupitant 235 mg and palonosetron 0.25 mg	235.25 mg (1 vial)	1/1/2019	Akynzeo®	fosnetupitant and palonosetron for injection, for intravenous use	Indicated in combination with dexamethasone in adults for the prevention of acute and delayed nausea and vomiting associated with initial and repeat courses of highly emetogenic cancer chemotherapy. Limitations of Use: Akynzeo for injection has not been studied for the prevention of nausea and vomiting associated with anthracycline plus cyclophosphamide chemotherapy.	5	18 years	N/A	N/A	Y	Y	9/1/2023: NC Suggested Max Monthly Units updated to align with NCTracks, which has been set to 5 units/month since 1/1/2019.	9/13/2023
Drugs	J1455	Injection, foscarnet sodium, per 1,000 mg	1,000 mg	1/1/2000	Foscavir®	foscarnet sodium injection	Indicated for the treatment of: • CMV retinitis in patients with acquired immunodeficiency syndrome (AIDS). Combination therapy with	996	18 years	N/A	N/A	Y	Y		6/4/2019

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

- Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.
- 11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacture participating in the Medicaid Drug Rebate Program (MDRP).
- The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.
- The HCPCS Code effective date represents the date the HCPCS code was established
- Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacture as they are not classified as covered outpatient drugs.
- Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/nccli-medicaid/medicaid-nccli-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Drugs	J1456	Injection, fosaprepitant (teva), not therapeutically equivalent to J1453, 1 mg	1 mg	1/1/2023	N/A	fosaprepitant for injection, for intravenous use (Teva)	Indicated in adults, in combination with other antiemetic agents, for the prevention of: • acute and delayed nausea and vomiting associated with initial and repeat courses of highly emetogenic cancer chemotherapy (HEC) including high-dose cisplatin. • delayed nausea and vomiting associated with initial and repeat courses of moderately emetogenic cancer chemotherapy (MEC). Limitations of Use: Fosaprepitant for Injection has not been studied for treatment of established nausea and vomiting.	750	18 years	N/A	N/A	Y	Y	9/2023: NC Suggested Max Monthly Units updated from 600 units to 750 units effective 1/1/2023 at DHB request	9/28/2023
Biologicals	J1458	Injection, galsulfase, 1 mg	1 mg	1/1/2007	Naglazyme®	galsulfase injection for intravenous use	Indicated for patients with Mucopolysaccharidosis VI (MPS VI; Maroteaux-Lamy syndrome). Naglazyme has been shown to improve walking and stair-climbing capacity.	700	N/A	N/A	N/A	Y	Y		7/2/2018
Immune Globulins	J1459	Injection, immune globulin (Privigen), intravenous, non-lyophilized (e.g., liquid), 500	500 mg	1/1/2009	Privigen®	immune globulin intravenous (human), 10% liquid	Indicated for the treatment of: • Primary humoral immunodeficiency (PI) • Chronic immune thrombocytopenic purpura (ITP) in patients age 15 years and older	840	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	Indication specific age restrictions: • Primary Humoral	7/3/2018
Immune Globulins	J1460	Injection, gamma globulin, intramuscular, 1 cc	1 cc	1/1/2000	GamaSTAN® S/D, GamaSTAN®	immune globulin (human), solution for intramuscular injection, less than 10 cc	Indicated: • For prophylaxis following exposure to hepatitis A. • To prevent or modify measles in a susceptible person exposed fewer than 6 days previously.	10	18 years	N/A	N/A	Y	Y		10/25/2018
Immune Globulins	J1554	Injection, immune globulin (asceniv), 500 mg	500 mg	4/1/2021	Asceniv™	immune globulin intravenous, human – sira 10% liquid	Indicated for the treatment of primary humoral immunodeficiency (PI) in adults and adolescents (12 to 17 years of age).	460	12 years	N/A	N/A	Y	Y		3/25/2021
Immune Globulins	J1555	Injection, immune globulin (Cuvitru), 100 mg	100 mg	1/1/2018	Cuvitru	immune globulin subcutaneous (human), 20% solution	Indicated as replacement therapy for primary humoral immunodeficiency (PI) in adult and pediatric patients two years of age and older.	14,880	2 years	N/A	N/A	Y	Y		9/12/2018
Immune Globulins	J1556	Injection, immune globulin (Bivigam), 500 mg	500 mg	1/1/2014	Bivigam®	immune globulin intravenous (human), 10% liquid	Indicated for the treatment of adults and pediatric patients 2 years of age and older with primary humoral immunodeficiency (PI).	480	2 years	N/A	N/A	Y	Y		2/16/2024
Immune Globulins	J1557	Injection, immune globulin, (Gammaglex), intravenous, non-lyophilized, (e.g. liquid), 500 mg	500 mg	1/1/2012	Gammaglex®	immune globulin intravenous (human), 5% and 10% liquid, for intravenous use	Gammaglex 5%: Indicated for the treatment of: • Chronic immune thrombocytopenic purpura (ITP). • Primary humoral immunodeficiency (PI) in adults and pediatric patients 2 years of age and older. Gammaglex 10%: Indicated for the treatment of: • Primary humoral immunodeficiency (PI) in adults. • Chronic immune thrombocytopenic purpura (ITP) in adults.	560	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	Product specific age restrictions: Gammaglex 5%: 2 years of age and older Gammaglex 10%: 18 years of age and older	9/21/2018
Immune Globulins	J1558	Injection, immune globulin (xembify), 100 mg	100 mg	7/1/2020	Xembify®	immune globulin subcutaneous, human – kihw 20% solution	Indicated for treatment of Primary Humoral Immunodeficiency (PI) in patients 2 years of age and older.	14,880	2 years	N/A	N/A	Y	Y		6/17/2020
Immune Globulins	J1559	Injection, immune globulin (Hizentra), 100 mg	100 mg	1/1/2011	Hizentra®	immune globulin subcutaneous (human), 20% liquid	• Indicated as replacement therapy for primary immunodeficiency (PI) in adults and pediatric patients 2 years of age and older. This includes, but is not limited to, the humoral immune defect in congenital agammaglobulinemia, common variable immunodeficiency, X-linked agammaglobulinemia, Wiskott-Aldrich syndrome and severe combined immunodeficiencies. • Indicated as maintenance therapy for the treatment of adult patients with chronic inflammatory demyelinating polyneuropathy (CIDP) to prevent relapse of neuromuscular disability and impairment.	2,800	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	Indication specific age restrictions: • PI - 2 years of age and older • CDIP - 18 years of age and older	7/16/2018

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

•Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.

•11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacture participating in the Medicaid Drug Rebate Program (MDRP).

•The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.

•The HCPCS Code effective date represents the date the HCPCS code was established

•Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacture as they are not classified as covered outpatient drugs.

•Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/ncci-medicaid/medicaid-ncci-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Immune Globulins	J1560	Injection, gamma globulin, intramuscular, over 10 cc (always use for any amount injected over 10cc and place number of units)	10 cc	1/1/2000	GamaSTAN® S/D, GamaSTAN®	immune globulin (human), solution for intramuscular injection greater than 10 cc	Indicated: • For prophylaxis following exposure to hepatitis A. • To prevent or modify measles in a susceptible person exposed fewer than 6 days previously. • To modify varicella. • To modify rubella in exposed women who will not consider a therapeutic abortion. • Not indicated for routine prophylaxis or treatment of viral hepatitis type B, rubella, poliomyelitis, mumps or varicella.	17	18 years	N/A	N/A	Y	Y		9/21/2018
Immune Globulins	J1561	Injection, immune globulin, (Gamunex-C/Gammaked), non-lyophilized (e.g. liquid), 500 mg	500 mg	1/1/2013	Gammaked™, Gamunex®-C	immune globulin injection (human), 10% caprylate/chromatography purified	Gamunex-C is indicated for: • Primary Humoral Immunodeficiency (PI) in patients 2 years of age and older • Idiopathic Thrombocytopenic Purpura (ITP) in adults and children • Chronic Inflammatory Demyelinating Polyneuropathy (CIDP) in adults Gammaked is indicated for: • Primary Humoral Immunodeficiency (PI) in patients 2 years of age and older	840	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	Indication specific age restrictions: • Primary Humoral Immunodeficiency (PI): 2 years of age and older • Idiopathic Thrombocytopenic	9/12/2018
Immune Globulins	J1566	Injection, immune globulin, intravenous, lyophilized (e.g. powder), not otherwise specified, 500mg	500 mg	1/1/2006	Carimune NF®, Gammagard S/D	immune globulin intravenous (human), lyophilized, nanofiltered - Carimune NF immune globulin intravenous (human), solvent detergent treated - Gammagard S/D	Carimune NF: Indicated for the maintenance treatment of patients with primary immunodeficiencies (PID), e.g., common variable immunodeficiency, X-linked agammaglobulinemia, severe combined immunodeficiency. Gammagard S/D: Indicated for the treatment of Primary Immunodeficiency (PI) in adults and pediatric patients two years of age or older, prevention of bacterial infections in hypogammaglobulinemia and/or recurrent bacterial infections associated with B-cell Chronic Lymphocytic Leukemia (CLL), prevention and/or control of bleeding in adult Chronic Idiopathic Thrombocytopenic Purpura (ITP) patients and prevention of coronary artery aneurysms associated with Kawasaki syndrome in pediatric patients.	952	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	Indication specific age restrictions: • Carimune NF: - PID: None - ITP: None • Gammagard S/D: - PI: 2 years of age and older - Chronic ITP: 18 years of age and older - Kawasaki Disease: None - CLL: None	9/8/2021
Immune Globulins	J1568	Injection, immune globulin, (Octagam), intravenous, non-lyophilized (e.g. liquid), 500 mg	500 mg	1/1/2008	Octagam®	immune globulin intravenous (human) liquid solution for intravenous administration	Octagam 5%: Indicated for the treatment of primary humoral immunodeficiency. Octagam 10%: Indicated for the treatment of: • Chronic immune thrombocytopenic purpura (ITP) in adults. • Dermatomyositis (DM) in adults.	• Octagam 5%: 336 units • Octagam 10%: 1,120 units	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	Product specific age restrictions: • Octagam 5%: 6 years of age and older.	8/25/2021
Immune Globulins	J1569	Injection, immune globulin, (Gammagard liquid), non-lyophilized, (e.g. liquid), 500 mg	500 mg	1/1/2008	Gammagard Liquid	immune globulin infusion (human), 10% solution, for intravenous and subcutaneous administration	Indicated as a: - replacement therapy for primary humoral immunodeficiency (PI) in adult and pediatric patients two years of age or older - maintenance therapy to improve muscle strength and disability in adult patients with Multifocal Motor Neuropathy (MMN)	840	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	Indication specific age restrictions: • PI: 2 years and older • MMN and CIDP: 18 years and older	2/27/2024
Drugs	J1570	Injection, ganciclovir sodium, 500 mg	500 mg	1/1/2000	Cytovene®-IV	ganciclovir sodium for injection, for intravenous use	Indicated for: • Treatment of CMV retinitis in immunocompromised individuals, including patients with acquired immunodeficiency syndrome (AIDS). • Prevention of CMV disease in adult transplant recipients at risk for CMV disease.	104	18 years	N/A	N/A	Y	Y		12/19/2022
Immune Globulins	J1571	Injection, hepatitis B immune globulin (Hepagab B), intramuscular, 0.5 mL	0.5 mL	1/1/2008	Hepagab B®	hepatitis b immune globulin intramuscular (human)	Indicated for post exposure prophylaxis in the following settings: • Acute Exposure to Blood Containing HBsAg • Perinatal Exposure of Infants Born to HBsAg-positive Mothers • Sexual Exposure to HBsAg-positive Persons	34	N/A	N/A	N/A	Y	Y		9/12/2018
Immune Globulins	J1572	Injection, immune globulin, (Flebogamma/Flebogamma DIF), intravenous, non-lyophilized (e.g. liquid), 500	500 mg	1/1/2008	Flebogamma®	immune globulin intravenous (human) for intravenous administration, 10% liquid preparation	Indicated for the treatment of: • Primary (inherited) Immunodeficiency (PI). • Chronic Primary Immune Thrombocytopenia (ITP) in patients 2 years of age and older.	560	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	Indication specific age restrictions: • Primary (inherited) Immunodeficiency (PI): None	7/3/2018
Immune Globulins	J1573	Injection, hepatitis B immune globulin (Hepagab B), intravenous, 0.5 mL	0.5 mL	1/1/2008	HepaGam B®	hepatitis b immune globulin intravenous (human)	Indicated for the prevention of hepatitis B virus recurrence after liver transplantation in HBsAg-positive transplant patients (HepaGam B) – IV only.	1,290	N/A	N/A	N/A	Y	Y		7/3/2018
Drugs	J1574	Injection, ganciclovir sodium (exela), not therapeutically equivalent to J1570, 500 mg	500 mg	1/1/2023	Ganzylk-RTU	ganciclovir injection, for intravenous use (Exela)	Indicated for the: • Treatment of CMV retinitis in immunocompromised adult patients, including patients with acquired immunodeficiency syndrome (AIDS). • Prevention of CMV disease in adult transplant recipients at risk for CMV disease.	104	18 years	N/A	N/A	Y	Y		12/6/2022

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

- Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.
- 11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacturer participating in the Medicaid Drug Rebate Program (MDRP).
- The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.
- The HCPCS Code effective date represents the date the HCPCS code was established
- Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacture as they are not classified as covered outpatient drugs.
- Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/ncci-medicare/medicaid-ncci-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Immune Globulins	J1575	Injection, immune globulin,hyaluronidase, (Hyqvia), 100 mg immune globulin	100 mg	1/1/2016	HyQvia*	immune globulin infusion 10% (human) with recombinant human hyaluronidase solution for subcutaneous administration	Indicated for treatment of primary immunodeficiency (PI) in patients two years of age and older. Indicated for the treatment of maintenance therapy in adults with chronic inflammatory demyelinating polyneuropathy (CIDP).	1,300	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	Indication Specific Age Restrictions: PI: 2 years of age and older CIDP: 18 years of age and older	2/27/2024
Immune Globulins	J1576	Injection, immune globulin (panzyga), intravenous, non-lyophilized (e.g., liquid), 500 mg	500 mg	7/1/2023	Panzyga*	immune globulin intravenous, human - ifas	Indicated for the treatment of: • Primary humoral immunodeficiency (PI) in patients 2 years of age and older. • Chronic immune thrombocytopenia (ITP) in adults. • Chronic inflammatory demyelinating polyneuropathy (CIDP) in adults.	1,120	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	Indication specific age restrictions: • Primary humoral immunodeficiency (PI) - 2 years of age and older • Chronic immune thrombocytopenia (ITP) and chronic inflammatory demyelinating polyneuropathy	6/22/2023
Drugs	J1580	Injection, garamycin, gentamicin, up to 80 mg	up to 80 mg	1/1/2000	N/A	gentamicin sulfate injection, for intravenous infusion or intramuscular injection	• Indicated in the treatment of serious infections caused by susceptible strains of the following microorganisms: Pseudomonas aeruginosa, Proteus species (indole-positive and indole-negative), Escherichia coli, Klebsiella-Enterobacter-Serratia species, Citrobacter species, and Staphylococcus species (coagulase-positive and coagulase-negative). • Clinical studies have shown gentamicin to be effective in bacterial neonatal sepsis; bacterial septicemia; and serious bacterial infections of the central nervous system (meningitis), urinary tract, respiratory tract,	279	N/A	N/A	N/A	Y	Y		6/4/2019
Biologicals	J1602	Injection, golimumab, 1 mg, for intravenous use	1 mg	1/1/2014	Simponi Aria*	golimumab injection, for intravenous use	Indicated for treatment of adult patients with: • Moderately to severely active Rheumatoid Arthritis (RA) in combination with methotrexate. • Active Ankylosing Spondylitis (AS). Indicated for treatment in patients 2 years of age and older with: • Active Psoriatic Arthritis (PsA). • Active polyarticular Juvenile Idiopathic Arthritis (pJIA)	560	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	Indication specific age restrictions: Rheumatoid Arthritis and Ankylosing Spondylitis: 18 years of age and older Polyarticular Juvenile Idiopathic Arthritis and Psoriatic Arthritis: 2 years of age and older	10/21/2020
Drugs	J1610	Injection, glucagon hydrochloride, per 1 mg	1 mg	1/1/2000	GlucaGen*	glucagon for injection, for subcutaneous, intramuscular, or intravenous use	Indicated for: • Treatment of severe hypoglycemia. • Use as a diagnostic aid for use during radiologic examinations to temporarily inhibit movement of the gastrointestinal tract.	10	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	Indication specific age restrictions: • Treatment of severe hypoglycemia: None • Diagnostic aid: 18 years of age and old	10/26/2018
Drugs	J1611	Injection, glucagon hydrochloride (fresenius kabi), not therapeutically equivalent to J1610, per 1 mg	1 mg	1/1/2023	N/A	glucagon for injection, for subcutaneous, intramuscular or intravenous use (Fresenius Kabi)	Indicated: • for the treatment of severe hypoglycemia in pediatric and adult patients with diabetes • as a diagnostic aid for use during radiologic examinations to temporarily inhibit movement of the gastrointestinal tract in adult patients	10	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	Indication specific age restrictions: • Diagnostic aid during radiologic examinations to	12/12/2022
Drugs	J1626	Injection, granisetron hydrochloride, 100 mcg	100 mcg	1/1/2000	N/A	granisetron hydrochloride injection, for intravenous use	Indicated for: • Prevention of nausea and/or vomiting associated with initial and repeat courses of emetogenic cancer	294	Indication Specific Age Restrictions	N/A	N/A	Y	Y	Indication specific: • Chemotherapy Induced	6/4/2019
Drugs	J1627	Injection, granisetron, extended-release, 0.1 mg	0.1 mg	1/1/2018	Sustol*	granisetron extended-release injection, for subcutaneous use	Indicated in combination with other antiemetics in adults for the prevention of acute and delayed nausea and vomiting associated with initial and repeat courses of moderately emetogenic chemotherapy (MEC) or anthracycline and cyclophosphamide (AC) combination chemotherapy regimens	500	18 years	N/A	N/A	Y	Y		10/26/2018
Drugs	J1630	Injection, haloperidol, up to 5 mg	up to 5 mg	1/1/2000	Haldol*	haloperidol lactate injection	Indicated for use in the treatment of schizophrenia and for the control of tics and vocal utterances of Tourette's Disorder.	124	18 years	N/A	N/A	Y	Y		10/26/2018

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

•Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.

•11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacturer participating in the Medicaid Drug Rebate Program (MDRP).

•The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.

•The HCPCS Code effective date represents the date the HCPCS code was established

•Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacturer as they are not classified as covered outpatient drugs.

•Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/ncci-medicaid/medicaid-ncci-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Drugs	J1631	Injection, haloperidol decanoate, per 50 mg	per 50 mg	1/1/2000	Haldol® Decanoate	haloperidol decanoate injection, for intramuscular use	Indicated for the treatment of schizophrenic patients who require prolonged parenteral antipsychotic therapy.	18	18 years	N/A	N/A	Y	Y		6/4/2019
Drugs	J1640	Injection, hemin, 1 mg	1 mg	1/1/2006	Panhematin®	hemin for injection	Indicated for amelioration of recurrent attacks of acute intermittent porphyria temporally related to the menstrual cycle in susceptible women, after initial carbohydrate therapy is known or suspected to be inadequate. Limitations of Use: • Before administering Panhematin, consider an appropriate period of carbohydrate loading (i.e., 400 g glucose/day for 1 to 2 days). • Panhematin is not effective in repairing neuronal damage due to progression of porphyria attacks.	14,700	16 years	N/A	Females Only	Y	Y		11/30/2021
Drugs	J1642	Injection, heparin sodium (heparin lock flush), per 10 units	10 units	1/1/2000	Hep-Flush®, Hep-Lock®	heparin sodium injection (heparin lock flush)	Intended to maintain patency of an indwelling venipuncture device designed for intermittent injection or infusion therapy or blood sampling. Heparin lock flush solution may be used following initial placement of the device in the vein, after each injection of a medication or after withdrawal of blood for laboratory tests. Heparin lock flush solution is not to be used for anticoagulant therapy.	4,500	N/A	N/A	N/A	Y	Y		10/26/2018
Drugs	J1643	Injection, heparin sodium (pfizer), not therapeutically equivalent to J1644, per 1000 units	1,000 units	1/1/2023	N/A	heparin sodium injection, for intravenous or subcutaneous use (Pfizer)	Indicated for: • Prophylaxis and treatment of venous thrombosis and pulmonary embolism • Prophylaxis and treatment of the thromboembolic complications associated with atrial fibrillation • Prevention of acute and chronic consumption coagulopathies • Prevention of clotting in arterial and cardiac surgery • Prophylaxis and treatment of peripheral arterial embolism • Anticoagulant use in transfusion, extracorporeal circulation, and dialysis procedures	465	N/A	N/A	N/A	Y	Y		12/12/2022
Drugs	J1644	Injection, heparin sodium, per 1,000 units	per 1,000 units	1/1/2000	N/A	heparin sodium injection, for intravenous or subcutaneous use	Indicated for: • Prophylaxis and treatment of venous thrombosis and pulmonary embolism. • Prevention of postoperative deep venous thrombosis and pulmonary embolism in patients undergoing major abdominotherapeutic surgery or who, for other reasons, are at risk of developing thromboembolic disease. • Atrial fibrillation with embolization. • Treatment of acute and chronic consumptive coagulopathies (disseminated intravascular coagulation). • Prevention of clotting in arterial and cardiac surgery. • Prophylaxis and treatment of peripheral arterial embolism. • Use as an anticoagulant in blood transfusions, extracorporeal circulation, and dialysis procedures.	465	N/A	N/A	N/A	Y	Y		6/4/2019
Drugs	J1645	Injection, dalteparin sodium, per 2,500 IU	per 2,500 IU	1/1/2000	Fragmin®	dalteparin sodium injection, for subcutaneous use	Indicated for: • Prophylaxis of ischemic complications of unstable angina and non-Q-wave myocardial infarction. • Prophylaxis of deep vein thrombosis (DVT) in abdominal surgery, hip replacement surgery or medical patients with severely restricted mobility during acute illness. • Extended treatment of symptomatic venous thromboembolism (VTE) to reduce the recurrence in patients with cancer. In these patients, the Fragmin therapy begins with the initial VTE treatment and	372	1 month	N/A	N/A	Y	Y		6/4/2019
Drugs	J1650	Injection, enoxaparin sodium, 10 mg	10 mg	1/1/2000	Lovenox®	enoxaparin sodium injection, for subcutaneous and intravenous use	Indicated for: • Prophylaxis of deep vein thrombosis (DVT) in abdominal surgery, hip replacement surgery, knee replacement surgery, or medical patients with severely restricted mobility during acute illness. • Inpatient treatment of acute DVT with or without pulmonary embolism. • Outpatient treatment of acute DVT without pulmonary embolism. • Prophylaxis of ischemic complications of unstable angina and non-Q-wave myocardial infarction (MI). • Treatment of acute ST-segment elevation myocardial infarction (STEMI) managed medically or with subsequent percutaneous coronary intervention (PCI).	930	18 years	N/A	N/A	Y	Y		6/5/2019

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

•Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.

•11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacturer participating in the Medicaid Drug Rebate Program (MDRP).

•The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.

•The HCPCS Code effective date represents the date the HCPCS code was established

•Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacture as they are not classified as covered outpatient drugs.

•Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/nccli-medicaid/medicaid-nccli-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Drugs	J1652	Injection, fondaparinux sodium, 0.5 mg	0.5 mg	1/1/2003	Arixtra*	fondaparinux sodium injection solution for subcutaneous injection	Indicated for: • Prophylaxis of deep vein thrombosis (DVT) in patients undergoing hip fracture surgery (including extended prophylaxis), hip replacement surgery, knee replacement surgery, or abdominal surgery. • Treatment of DVT or acute pulmonary embolism (PE) when administered in conjunction with Coumadin.	520	18 years	N/A	N/A	Y	Y		10/10/2018
Drugs	J1720	Injection, hydrocortisone sodium succinate, up to 100 mg	up to 100 mg	1/1/2000	Solu-Cortef®	hydrocortisone sodium succinate for injection, for intravenous or intramuscular administration	When oral therapy is not feasible, and the strength, dosage form, and route of administration of the drug reasonably lend the preparation to the treatment of the condition, the intravenous or intramuscular use of Solu-Cortef is indicated as follows: • Allergic States: Control of severe or incapacitating allergic conditions intractable to adequate trials of conventional treatment in asthma, atopic dermatitis, contact dermatitis, drug hypersensitivity reactions, serum sickness, transfusion reactions.	155	N/A	N/A	N/A	Y	Y		6/28/2021
Drugs	J1729	Injection, hydroxyprogesterone caproate, Not Otherwise Specified, 10 mg	10 mg	1/1/2018	N/A	hydroxyprogesterone caproate injection	Indicated in non-pregnant women: • For the treatment of advanced adenocarcinoma of the uterine corpus (Stage III or IV) • In the management of amenorrhea (primary and secondary) and abnormal uterine bleeding due to hormonal imbalance in the absence of organic pathology, such as submucous fibroids or uterine cancer • As a test for endogenous estrogen production and for the production of secretory endometrium and indicated for use in adults for the management of moderate-to-severe pain, alone or in combination with non-NSAID analgesics.	3,100	N/A	N/A	Indicated only for non-pregnant women.	Y	Y		6/4/2019
Drugs	J1738	Injection, meloxicam, 1 mg	1 mg	10/1/2020	Anjeso™	meloxicam injection, for intravenous use	Limitation of Use: Indicated for the treatment of osteoporosis in postmenopausal women.	930	18 years	N/A	N/A	Y	Y		9/21/2020
Drugs	J1740	Injection, ibandronate sodium, 1 mg	1 mg	1/1/2007	Boniva*	ibandronate injection, for intravenous use	Limitations of Use: Optimal duration of use has not been determined. For patients at low-risk form fracture, consider drug discontinuation after 3 to 5 years of use.	3	40 years	N/A	Females Only	Y	Y		10/18/2018
Drugs	J1742	Injection, ibutilide fumarate, 1 mg	1 mg	1/1/2000	Corvert*	ibutilide fumarate injection, for intravenous infusion	Indicated for the rapid conversion of atrial fibrillation or atrial flutter of recent onset to sinus rhythm. Patients with atrial arrhythmias of longer duration are less likely to respond to ibutilide. The effectiveness	10	18 years	N/A	N/A	Y	Y		10/18/2018
Drugs	J1743	Injection, idursulfase, 1 mg	1 mg	1/1/2008	Elaforce*	idursulfase injection, for intravenous use	Indicated for patients with Hunter syndrome (Mucopolysaccharidosis II, MPS II). Elaforce has been shown to improve walking capacity in patients 5 years and older. In patients 16 months to 5 years of age, no data are available to demonstrate improvement in disease-related symptoms or long term clinical outcome; however, treatment with Elaforce has reduced spleen volume similarly to that of adults and children 5 years of age and older. The safety and efficacy of Elaforce have not been established in pediatric patients less than 16 months of age.	360	16 months	N/A	N/A	Y	Y		6/4/2019
Biologicals	J1744	Injection, icatibant, 1 mg	1 mg	1/1/2013	Firazyr*	icatibant injection, for subcutaneous use	Indicated for the treatment of acute attacks of hereditary angioedema (HAE).	2700	18 years	N/A	N/A	Y	Y		6/4/2019
Biologicals	J1745	Injection, infliximab, excludes biosimilar, 10 mg	10 mg	1/1/2017	Remicade®	infliximab lyophilized concentrate for injection, for intravenous use	Indicated for: • Crohn's Disease: reducing signs and symptoms and inducing and maintaining clinical remission in adult	140	6 years	N/A	N/A	Y	Y		6/6/2019
Biologicals	J1746	Injection, ibalizumab-uiyk, 10 mg	10 mg	1/1/2019	Trogarzo™	ibalizumab-uiyk injection, for intravenous use	Indicated for use in combination with other antiretroviral(s), for the treatment of human immunodeficiency virus type 1 (HIV-1) infection in heavily treatment-experienced adults with multidrug	360	18 years	N/A	N/A	Y	Y		2/16/2024
Drugs	J1750	Injection, iron dextran, 50 mg	50 mg	1/1/2009	INFeD*	iron dextran injection	Indicated for treatment of patients with documented iron deficiency in whom oral administration is unsatisfactory or impossible.	62	4 months	N/A	N/A	Y	Y		10/26/2018
Drugs	J1756	Injection, iron sucrose, 1 mg	1 mg	1/1/2003	Venofer*	iron sucrose injection for intravenous use	Indicated for the treatment of iron deficiency anemia in patients with chronic kidney disease (CKD).	2,000	2 years	N/A	N/A	Y	Y		7/29/2020
Biologicals	J1786	Injection, imiglucerase, 10 units	10 units	1/1/2011	Cerezyme®	imiglucerase for injection, for intravenous use	Indicated for long-term enzyme replacement therapy for pediatric and adult patients with a confirmed diagnosis of Type 1 Gaucher disease that results in one or more of the following conditions: • anemia	2,520	2 years	N/A	N/A	Y	Y		6/22/2023
Drugs	J1790	Injection, droperidol, up to 5 mg	up to 5 mg	1/1/2000	N/A	droperidol injection for intravenous or intramuscular use	Indicated to reduce nausea and vomiting associated with surgical and diagnostic procedures.	5	2 years	N/A	N/A	Y	Y		10/4/2018

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

•Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.

•11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacturer participating in the Medicaid Drug Rebate Program (MDRP).

•The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.

•The HCPCS Code effective date represents the date the HCPCS code was established

•Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacture as they are not classified as covered outpatient drugs.

•Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/ncci-medicaid/medicaid-ncci-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Drugs	J1800	Injection, propranolol HCl, up to 1 mg	up to 1 mg	1/1/2000	N/A	propranolol hydrochloride injection, solution	Indicated for supraventricular arrhythmias, ventricular tachycardias, tachyarrhythmias of digitalis intoxication and resistant tachyarrhythmias due to excessive catecholamine action during anesthesia.	N/A	18 years	N/A	N/A	Y	Y		8/29/2018
Biologicals	J1812	Insulin (fiasp), per 5 units	5 units	7/1/2023	Fiasp*	insulin aspart injection for subcutaneous or intravenous use	Indicated to improve glycemic control in adult and pediatric patients with diabetes mellitus.	N/A	2 years	N/A	N/A	Y	Y		6/19/2023
Biologicals	J1814	Insulin (lyumjev), per 5 units	5 units	7/1/2023	Lyumjev*	insulin lispro-aabc injection, for subcutaneous or intravenous use	Indicated to improve glycemic control in adult and pediatric patients with diabetes mellitus.	N/A	1 year	N/A	N/A	Y	Y		6/19/2023
Biologicals	J1815	Injection, insulin, per 5 units	5 units	1/1/2003	Various brand names	insulin, injectable suspension	Indicated to improve glycemic control in adults and pediatric patients with diabetes mellitus.	3,100	N/A	N/A	N/A	Y	Y		6/19/2023
Biologicals	J1823	Injection, inebilizumab-cdon, 1 mg	1 mg	1/1/2021	Uplizna™	inebilizumab-cdon injection, for intravenous use	Indicated for the treatment of neuromyelitis optica spectrum disorder (NMOSD) in adult patients who are anti-aquaporin-4 (AQP4) antibody positive.	600	18 years	N/A	N/A	Y	Y		12/28/2020
Biologicals	J1830	Injection, interferon beta-1B, 0.25 mg	0.25 mg	1/1/2000	Betaseron*, Extavia*	interferon beta-1b for injection, for subcutaneous use	Indicated for the treatment of relapsing forms of multiple sclerosis to reduce the frequency of clinical exacerbations. Patients with multiple sclerosis in whom efficacy has been demonstrated include patients who have experienced a first clinical episode and have MRI features consistent with multiple sclerosis.	16	18 years	N/A	N/A	Y	Y		6/4/2019
Drugs	J1833	Injection, isavuconazonium sulfate, 1 mg	1 mg	1/1/2016	Cresamba*	isavuconazonium sulfate for injection for intravenous administration	Indicated in adults and pediatric patients 1 year of age and older for the treatment of: • Invasive aspergillosis • Invasive mucormycosis	13,020	1 year	N/A	N/A	Y	Y		2/16/2024
Drugs	J1885	Injection, ketorolac tromethamine, per 15 mg	15 mg	1/1/2000	N/A	ketorolac tromethamine injection for intravenous or intramuscular use	Indicated for the short-term management (≤ 5 days) of moderately-severe acute pain requiring analgesia at the opioid level in adults, usually in a postoperative setting.	40	17 years	N/A	N/A	Y	Y		4/9/2019
Drugs	J1930	Injection, lanreotide, 1 mg	1 mg	1/1/2009	Somatuline® Depot	lanreotide injection, for subcutaneous use	Indicated for the long-term treatment of acromegalic patients who have had an inadequate response to or cannot be treated with surgery and/or radiotherapy. Indicated for the treatment of adult patients with unresectable, well- or moderately-differentiated, locally advanced or metastatic gastroenteropancreatic neuroendocrine tumors (GEP-NETs) to improve progression-free survival. Indicated for the treatment of adults with carcinoid syndrome; when used, it reduces the frequency of	240	18 years	N/A	N/A	Y	Y		10/26/2018
Biologicals	J1931	Injection, laronidase, 0.1 mg	0.1 mg	1/1/2005	Aldurazyme*	laronidase solution for intravenous infusion only	Indicated for patients with Hurler and Hurler-Scheie forms of Mucopolysaccharidosis I (MPS I) and for patients with the Scheie form who have moderate to severe symptoms. The risks and benefits of treating mildly affected patients with the Scheie form have not been established. Aldurazyme has been shown to improve pulmonary function and walking capacity. Aldurazyme has not been evaluated for effects on the central nervous system manifestations of the disorder.	4,060	6 months	N/A	N/A	Y	Y		4/10/2019
Drugs	J1932	Injection, lanreotide, (cipla), 1 mg	1 mg	10/1/2022	N/A	lanreotide injection, for subcutaneous use (Cipla)	Indicated for: • The long-term treatment of acromegalic patients who have had an inadequate response to or cannot be treated with surgery and/or radiotherapy. • The treatment of adult patients with unresectable, well- or moderately differentiated, locally advanced	240	18 years	N/A	N/A	Y	Y		9/15/2022
Drugs	J1940	Injection, furosemide, up to 20 mg	up to 20 mg	1/1/2000	Lasix*	furosemide injection	Indicated for the treatment of edema associated with congestive heart failure, cirrhosis of the liver, and renal disease, including the nephrotic syndrome. - Furosemide is particularly useful when an agent with greater diuretic potential is desired. Indicated as an adjunct in the treatment of pulmonary edema.	310	N/A	N/A	N/A	Y	Y		10/26/2018
Drugs	J1943	Injection, aripiprazole lauroxil, (aristada initio), 1 mg	1 mg	10/1/2019	Aristada Initio™	aripiprazole lauroxil extended release injectable suspension, for intramuscular use	Indicated for the initiation of Aristada when used for the treatment of schizophrenia in adults in combination with oral aripiprazole.	675	18 years	N/A	N/A	Y	Y	• Cervical Dystonia: Safety and effectiveness in pediatric patients have not been established.	9/27/2019
Drugs	J1944	Injection, aripiprazole lauroxil, (aristada), 1 mg	1 mg	10/1/2019	Aristada*	aripiprazole lauroxil extended release injectable suspension, for intramuscular use	Indicated for the treatment of schizophrenia.	1,064	18 years	65 years	N/A	Y	Y		9/27/2019

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

- Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.
- 11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacture participating in the Medicaid Drug Rebate Program (MDRP).
- The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.
- The HCPCS Code effective date represents the date the HCPCS code was established
- Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacture as they are not classified as covered outpatient drugs.
- Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/ncci-medicaid/medicaid-ncci-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Drugs	J1950	Injection, leuprolide acetate (for depot suspension), per 3.75 mg	per 3.75 mg	1/1/2000	Lupron Depot*, Lupron Depot-PED*	leuprolide acetate for depot suspension, for intramuscular use	Lupron Depot 3.75 mg and 11.25 mg are indicated for: • Endometriosis • Management of endometriosis, including pain relief and reduction of endometriotic lesions.	12	Product Specific Age Restrictions (see comments)	N/A	Lupron Depot: Females Only Lupron Depot-	Y	Y	Product specific age restrictions: Lupron Depot:	2/19/2024
Drugs	J1951	Injection, leuprolide acetate for depot suspension (fensolvi), 0.25 mg	0.25 mg	7/1/2021	Fensolvi*	leuprolide acetate for injectable suspension, for subcutaneous use	Indicated for the treatment of pediatric patients 2 years of age and older with central precocious puberty.	180	2 years	N/A	N/A	Y	Y		6/28/2021
Drugs	J1952	Leuprolide injectable, camcevi, 1 mg	1 mg	1/1/2022	Camcevi™	leuprolide injectable emulsion, for subcutaneous use	Indicated for the treatment of adult patients with advanced prostate cancer.	42	18 years	N/A	Males Only	Y	Y		5/16/2022
Drugs	J1953	Injection, levetiracetam, 10 mg	10 mg	1/1/2009	Keppra®	levetiracetam injection, for intravenous use	Indicated as an adjunctive therapy, as an alternative when oral administration is temporarily not feasible, for the treatment of: • Partial onset seizures in patients 1 month of age and older with epilepsy • Myoclonic seizures in patients 12 years of age and older with juvenile myoclonic epilepsy • Primary generalized tonic-clonic seizures in patients 6 years of age and older with idiopathic generalized	9,300	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	Indication specific age restrictions: • Partial Onset Seizures: 1 month of age and older • Myoclonic Seizures in	10/10/2018
Drugs	J1954	Injection, leuprolide acetate for depot suspension (lutrate), 7.5 mg	7.5 mg	1/1/2023	Lutate Depot	leuprolide acetate for depot suspension	Indicated for treatment of advanced prostate cancer.	3	18 years	N/A	Males Only	Y	Y		3/16/2023
Drugs	J1955	Injection, levocarnitine, per 1 g	1 g	1/1/2000	Carnitor®	levocarnitine injection for intravenous use	Indicated for: • the acute and chronic treatment of patients with an inborn error of metabolism which results in secondary carnitine deficiency.	1,302	N/A	N/A	N/A	Y	Y		4/10/2019
Drugs	J1956	Injection, levofloxacin, 250 mg	250 mg	1/1/2000	Levaquin®	levofloxacin injection for intravenous use	Indicated in adults (>=18 years of age) with infections caused by designated, susceptible bacteria: • Pneumonia: Nosocomial and Community Acquired • Skin and Skin Structure Infections: Complicated and Uncomplicated • Chronic bacterial prostatitis • Inhalational Anthrax, Post-Exposure • Plague • Urinary Tract Infections: Complicated and Uncomplicated • Acute Pyelonephritis • Acute Bacterial Exacerbation of Chronic Bronchitis • Acute Bacterial Sinusitis Usage: To reduce the development of drug-resistant bacteria and maintain the effectiveness of Levaquin and other antibacterial drugs, Levaquin should be used only to treat or prevent infections that are proven or strongly suspected to be caused by bacteria.	62	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	Indication specific: Inhalation Anthrax (Post-Exposure): 6 months and older. Plague: 6 months and older. All other indications: 18 years of age and older.	6/5/2019
Drugs	J1961	Injection, lenacapavir, 1 mg	1 mg	7/1/2023	Sunlenca®	lenacapavir injection, for subcutaneous use	Indicated for the treatment of HIV-1 infection in heavily treatment-experienced adults with multidrug resistant HIV-1 infection failing their current antiretroviral regimen due to resistance, intolerance, or safety considerations.	927	18 years	N/A	N/A	Y	Y		6/22/2023
Drugs	J1980	Injection, hyoscyamine sulfate, up to 0.25 mg	up to 0.25 mg	1/1/2000	Levsin®	hyoscyamine sulfate injection	• Is effective as adjunctive therapy in the treatment of peptic ulcer. • In acute episodes, Levsin injection can be used to control gastric secretion, visceral spasm and hypermotility in spastic colitis, spastic bladder, cystitis, pylorospasm, and associated abdominal cramps.	248	N/A	N/A	N/A	Y	Y		7/2/2018
Drugs	J2001	Injection, lidocaine HCL for intravenous infusion, 10 mg	10 mg	1/1/2004	N/A	lidocaine hydrochloride injection, solution	• Administered intravenously or intramuscularly, is specifically indicated in the acute management of ventricular arrhythmias such as those occurring in relation to acute myocardial infarction, or during cardiac manipulation, such as cardiac surgery. • Indicated for production of local or regional anesthesia by infiltration techniques such as percutaneous injection and intravenous regional anesthesia by peripheral nerve block techniques such as brachial plexus and intercostal and by central neural techniques such as lumbar and caudal epidural blocks, when the accepted procedures for these techniques as described in standard textbooks are observed.	35	N/A	N/A	N/A	Y	Y		10/31/2018

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

- Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.
- 11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacture participating in the Medicaid Drug Rebate Program (MDRP).
- The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.
- The HCPCS Code effective date represents the date the HCPCS code was established
- Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacture as they are not classified as covered outpatient drugs.
- Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/ncci-medicaid/medicaid-ncci-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Drugs	J2010	Injection, lincocin HCl, up to 300 mg	300 mg	1/1/2000	Lincocin*	lincocin hydrochloride injection, solution	Indicated for the treatment of serious infections due to susceptible strains of streptococci, pneumococci, and staphylococci. Its use should be reserved for penicillin-allergic patients or other patients for whom, in the judgment of the physician, a penicillin is inappropriate.	837	1 month	N/A	N/A	Y	Y		10/26/2018
Drugs	J2020	Injection, linezolid, 200 mg	200 mg	1/1/2002	Zyvox*	linezolid injection, solution	Indicated in adults and children for the treatment of the following infections caused by susceptible Gram-positive bacteria: nosocomial pneumonia; community-acquired pneumonia, complicated skin and skin structure infections, including diabetic foot infections, without concomitant osteomyelitis, uncomplicated skin and skin structure infections, vancomycin-resistant Enterococcus faecium infections.	168	N/A	N/A	N/A	Y	Y		10/26/2018
Drugs	J2021	Injection, linezolid (hospira), not therapeutically equivalent to J2020, 200 mg	200 mg	1/1/2023	N/A	linezolid injection, for intravenous use (Hospira)	Indicated in adults and children for the treatment of the following infections caused by susceptible Gram-positive bacteria: Nosocomial pneumonia; Community-acquired pneumonia; Complicated skin and skin structure infections, including diabetic foot infections, without concomitant osteomyelitis; Vancomycin-	168	N/A	N/A	N/A	Y	Y		12/12/2022
Drugs	J2060	Injection, lorazepam, 2 mg	2 mg	1/1/2000	Ativan*	lorazepam injection for intravenous or intramuscular use	Indicated: • In adult patients for preanesthetic medication, producing sedation (sleepiness or drowsiness), relief of anxiety and a decreased ability to recall events related to the day of surgery. • For treatment of status epilepticus.	124	18 years	N/A	N/A	Y	Y		4/10/2019
Drugs	J2150	Injection, mannitol, 25% in 50 mL	50 mL	1/1/2000	N/A	mannitol injection, for intravenous use	Indicated for the reduction of: • Intracranial pressure and treatment of cerebral edema • Elevated intraocular pressure	713	N/A	N/A	N/A	Y	Y		11/29/2021
Drugs	J2175	Injection, meperidine hydrochloride, per 100 mg	100 mg	1/1/2000	Demerol™	meperidine hydrochloride injection, for subcutaneous, intramuscular, and intravenous use	Indicated for preoperative medication, support of anesthesia, obstetrical analgesia, and for the management of pain severe enough to require an opioid analgesic and for which alternative treatments are inadequate. Limitations of Use:	124	N/A	N/A	N/A	Y	Y		10/26/2018
Drugs	J2186	Injection, meropenem and vaborbactam, 10mg/10mg (20mg)	1 vial	1/1/2019	Vabomere™	meropenem and vaborbactam for injection, for intravenous use	Indicated for the treatment of patients 18 years and older with complicated urinary tract infections (cUTI) including pyelonephritis caused by designated susceptible bacteria. To reduce the development of drug-resistant bacteria and maintain the effectiveness of Vabomere and other antibacterial drugs, Vabomere	8,400	18 years	N/A	N/A	Y	Y		10/26/2018
Drugs	J2210	Injection, methylergonovine maleate, up to 0.2 mg	up to 0.2 mg	1/1/2000	Methergine®	methylergonovine maleate injection	Indicated • Following delivery of the placenta, for routine management of uterine atony, hemorrhage, and subinvolution of the uterus. • For control of uterine hemorrhage in the second stage of labor following delivery of the anterior shoulder.	5	Women of childbearing age	Women of childbearing age	Females Only	Y	Y		10/31/2018
Drugs	J2249	Injection, remimazolam, 1 mg	1 mg	7/1/2023	Byfavo™	remimazolam for injection, for intravenous use	Indicated for the induction and maintenance of procedural sedation in adults undergoing procedures lasting 30 minutes or less.	200	18 years	N/A	N/A	Y	Y		6/22/2023
Drugs	J2250	Injection, midazolam hydrochloride, per 1 mg	1 mg	1/1/2000	N/A	midazolam hydrochloride injection for intravenous or	Indicated: • Intramuscularly or intravenously for preoperative sedation/anxiolysis/amnesia	25	N/A	N/A	N/A	Y	Y		10/31/2018

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

•Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.

•11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacturer participating in the Medicaid Drug Rebate Program (MDRP).

•The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.

•The HCPCS Code effective date represents the date the HCPCS code was established

•Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacturer as they are not classified as covered outpatient drugs.

•Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/ncci-medicaid/medicaid-ncci-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Drugs	J2251	Injection, midazolam hydrochloride (wg critical care), not therapeutically equivalent to J2250, per 1 mg	1 mg	1/1/2023	N/A	midazolam in sodium chloride injection for intravenous use (WG Critical Care)	Indicated for: • Continuous intravenous infusion for sedation of intubated and mechanically ventilated adult, pediatric, and neonatal patients as a component of anesthesia or during treatment in a critical care setting.	500	N/A	N/A	N/A	Y	Y		12/12/2022
Drugs	J2260	Injection, milirone lactate, per 5 mg	per 5 mg	1/1/2000	N/A	milirone lactate injection	Indicated for the short-term intravenous treatment of patients with acute decompensated heart failure.	64	18 years	N/A	N/A	Y	Y		6/6/2019
Drugs	J2270	Injection, morphine sulfate, up to 10 mg	up to 10 mg	1/1/2000	N/A	morphine sulfate injection, up to 10 mg	Indicated for the management of pain severe enough to require an opioid analgesic and for which alternative treatments are inadequate. [Limitations of Use: Because of the risks of addiction, abuse, and misuse with opioids, even at recommended doses, reserve Morphine Sulfate Injection, for use in patients for whom alternative treatment options (e.g., non-opioid analgesics or opioid combination products): • Have not been tolerated, or are not expected to be tolerated. • Have not provided adequate analgesia, or are not expected to provide adequate analgesia Prior: Indicated for: • the relief of severe acute and chronic pain • to relieve preoperative apprehension • to facilitate anesthesia induction • the treatment of dyspnea associated with acute left ventricular failure and pulmonary edema • analgesia during labor • anxiety • anesthesia • to control postoperative pain.	527	N/A	N/A	N/A	Y	Y		6/7/2019
Drugs	J2272	Injection, morphine sulfate (Fresenius Kabl), not therapeutically equivalent to J2270, up to 10 mg	10 mg	1/1/2023	N/A	morphine sulfate injection, for intravenous or intramuscular use, CII (Fresenius Kabl)	Indicated for the management of pain severe enough to require an opioid analgesic and for which alternative treatments are inadequate. Limitations of Use • Mitigo: for use in continuous microinfusion devices and indicated only for intrathecal or epidural infusion in the management of intractable chronic pain severe enough to require an opioid analgesic and for which alternative treatments are inadequate.	527	18 years	N/A	N/A	Y	Y		12/12/2022
Drugs	J2274	Injection, morphine sulfate, preservative-free for epidural or intrathecal use, 10 mg	10 mg	1/1/2015	Duramorph®, Infumorph®, Mitigo	morphine sulfate injection preservative-free	Indicated for the management of pain severe enough to require an opioid analgesic and for which alternative treatments are inadequate.	100	18 years	N/A	N/A	Y	Y		4/9/2022
Drugs	J2277	Injection, motixafortide, 0.25 mg	0.25 mg	4/1/2024	Aphexda™	motixafortide for injection, for subcutaneous use	Indicated in combination with filgrastim (G-CSF) to mobilize hematopoietic stem cells to the peripheral blood for collection and subsequent autologous transplantation in patients with multiple myeloma.	1,488	18 years	N/A	N/A	Y	Y		3/22/2024
Drugs	J2278	Injection, ziconotide, 1 microgram	1 mcg	1/1/2006	Prialt®	ziconotide solution, intrathecal infusion	Indicated for the management of severe chronic pain in patients for whom intrathecal therapy is warranted, and who are intolerant of or refractory to other treatment, such as systemic analgesics, adjunctive therapies, or intrathecal morphine.	620	18 years	N/A	N/A	Y	Y		9/21/2018
Drugs	J2300	Injection, nalbuphine hydrochloride, per 10 mg	10 mg	1/1/2000	N/A	nalbuphine hydrochloride injection, solution	Indicated for management of pain severe enough to require an opioid analgesic and for which alternative treatments are inadequate. Also can be used as a supplement to balanced anesthesia, for pre/post operative analgesia and obstetrical analgesia during labor and delivery.	248	18 years	N/A	N/A	Y	Y		10/26/2018

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

- Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.
- 11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacture participating in the Medicaid Drug Rebate Program (MDRP).
- The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.
- The HCPCS Code effective date represents the date the HCPCS code was established
- Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacture as they are not classified as covered outpatient drugs.
- Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/nccli-medicaid/medicaid-nccli-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Drugs	J2310	Injection, naloxone hydrochloride, per 1 mg	1 mg	1/1/2000	Narcan®	naloxone hydrochloride injection	Indicated for the complete or partial reversal of opioid depression, including respiratory depression, induced by natural and synthetic opioids including: propoxyphene, methadone, nalbuphine, butorphanol and pentazocine; It is also indicated for the diagnosis of suspected opioid tolerance or acute opioid overdose.	N/A	N/A	N/A	N/A	Y	Y		10/26/2018
Drugs	J2311	Injection, naloxone hydrochloride (zimhi), 1 mg	1 mg	1/1/2023	Zimhi™	naloxone hydrochloride injection for intramuscular or subcutaneous use	Indicated in adult and pediatric patients for the emergency treatment of known or suspected opioid overdose, as manifested by respiratory and/or central nervous system depression.	50	N/A	N/A	N/A	Y	Y		12/6/2022
Drugs	J2315	Injection, naltrexone, depot form, 1 mg	1 mg	1/1/2007	Vivitrol®	naltrexone for extended-release injectable suspension,	• Indicated for the treatment of alcohol dependence in patients who are able to abstain from alcohol in an outpatient setting prior to initiation of treatment with Vivitrol. Patients should not be actively drinking at	760	18 years	N/A	N/A	Y	Y	9/1/2023: Generic Name updated to align with	9/13/2023
Biologicals	J2323	Injection, natalizumab, 1 mg	1 mg	1/1/2008	Tysabri®	natalizumab injection, for intravenous use	Indicated for treatment of: Multiple Sclerosis (MS) • Tysabri is indicated as monotherapy for the treatment of patients with relapsing forms of multiple	600	18 years	N/A	N/A	Y	Y		10/26/2018
Drugs	J2326	Injection, nusinersen, 0.1 mg	0.1 mg	1/1/2018	Spinraza®	nusinersen injection, for intrathecal use	Indicated for the treatment of spinal muscular atrophy (SMA) in pediatric and adult patients.	360	N/A	N/A	N/A	Y	Y		5/6/2021
Biologicals	J2327	Injection, risankizumab-rzaa, intravenous, 1 mg	1 mg	1/1/2023	Skyrizi®	risankizumab-rzaa injection, for intravenous use	Indicated for the treatment of moderately to severely active Crohn's disease in adults.	1,200	18 years	N/A	N/A	Y	Y		12/6/2022
Biologicals	J2329	Injection, ublituximab-xiyy, 1mg	1 mg	7/1/2023	Briumvi™	ublituximab-xiyy injection, for intravenous use	Indicated for the treatment of relapsing forms of multiple sclerosis (MS), to include clinically isolated syndrome, relapsing-remitting disease, and active secondary progressive disease, in adults.	600	18 years	N/A	N/A	Y	Y		6/22/2023
Drugs	J2353	Injection, octreotide, depot form for intramuscular injection, 1 mg	1 mg	1/1/2004	Sandostatin® LAR Depot	octreotide acetate for injectable suspension	Indicated for treatment in patients who have responded to and tolerated sandostatin injection subcutaneous injection for: • Acromegaly	40	18 years	N/A	N/A	Y	Y		7/16/2018
Drugs	J2354	Injection, octreotide, non-depot form for subcutaneous	25 mcg	1/1/2004	Sandostatin®	octreotide acetate, injection	Indicated: • To reduce blood levels of growth hormone and IGF-I (somatomedin C) in acromegaly patients who have	1,860	18 years	N/A	N/A	Y	Y		7/16/2018
Drugs	J2358	Injection, olanzapine, long-acting, 1 mg	1 mg	1/1/2011	Zyprexa® Relprevv™	olanzapine pamoate for extended release injectable	Indicated for the treatment of schizophrenia.	900	18 years	N/A	N/A	Y	Y		9/21/2018
Drugs	J2359	Injection, olanzapine, 0.5 mg	0.5 mg	10/1/2023	Zyprexa®	olanzapine injection, powder, intramuscular	Indicated for the treatment of acute agitation associated with schizophrenia and bipolar I mania.	1,860	13 years	N/A	N/A	Y	Y		9/28/2023
Drugs	J2360	Injection, orphenadrine citrate, up to 60 mg	up to 60 mg	1/1/2000	Norflex®	orphenadrine citrate injection	Indicated as an adjunct to rest, physical therapy, and other measures for the relief of discomfort associated with acute painful musculoskeletal conditions.	20	18 years	N/A	N/A	Y	Y		7/16/2018
Drugs	J2371	Injection, phenylephrine hydrochloride, 20 micrograms	20 mcg	7/1/2023	Vaculep®	phenylephrine hydrochloride injection for intravenous use	Indicated for the treatment of clinically important hypotension resulting primarily from vasodilation in the setting of anesthesia.	31,000	18 years	N/A	N/A	Y	Y		6/22/2023
Drugs	J2372	Injection, phenylephrine hydrochloride (biorphen), 20 micrograms	20 mcg	7/1/2023	Biorphen®	phenylephrine hydrochloride injection, for intravenous use	Indicated for the treatment of clinically important hypotension resulting primarily from vasodilation in the setting of anesthesia.	31,000	18 years	N/A	N/A	Y	Y		6/22/2023
Drugs	J2401	Injection, chlorprocaine hydrochloride, per 1 mg	1 mg	1/1/2023	Nesacaine®, Nesacaine®-MPF	chlorprocaine HCl injection	Multidose vial with preservatives: Indicated for the production of local anesthesia by infiltration and peripheral nerve block. Single dose vial without preservatives and without EDTA: Indicated for the production of local anesthesia	1,000	N/A	N/A	N/A	Y	Y		12/6/2022

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

- Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.
- 11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacture participating in the Medicaid Drug Rebate Program (MDRP).
- The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.
- The HCPCS Code effective date represents the date the HCPCS code was established
- Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacture as they are not classified as covered outpatient drugs.
- Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/nccli-medicaid/medicaid-nccli-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Drugs	J2402	Injection, chloroprocaine hydrochloride (clorotekal), per 1 mg	1 mg	1/1/2023	Clorotekal*	chloroprocaine hydrochloride injection, for intrathecal use	Indicated for intrathecal injection in adults for the production of subarachnoid block (spinal anesthesia).	50	18 years	N/A	N/A	Y	Y		12/6/2022
Drugs	J2403	Chloroprocaine hcl ophthalmic, 3% gel, 1 mg	1 mg	4/1/2023	iheezo™	chloroprocaine hydrochloride ophthalmic gel 3%, for topical ophthalmic use	Chloroprocaine hydrochloride ophthalmic gel is indicated for ocular surface anesthesia.	4,000	18 years	N/A	N/A	Y	Y		12/1/2023
Drugs	J2405	Injection, ondansetron hydrochloride, per 1 mg	1 mg	1/1/2000	Zofran*	ondansetron hydrochloride injection, for intravenous or intramuscular use	Indicated for the prevention of: • Nausea and vomiting associated with initial and repeat courses of emetogenic cancer chemotherapy. • Postoperative nausea and/or vomiting.	720	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	Indication specific age restrictions: • Prevention of nausea and	9/27/2018
Drugs	J2406	Injection, oritavancin (kimyrsa), 10 mg	10 mg	10/1/2021	Kimyrsa™	oritavancin for injection, for intravenous use	Indicated for the treatment of adult patients with acute bacterial skin and skin structure infections (ABSSSI) caused by susceptible isolates of the following Gram-positive microorganisms: Staphylococcus aureus (including methicillin-susceptible and methicillin-resistant isolates), Streptococcus pyogenes, Streptococcus agalactiae, Streptococcus dysgalactiae, Streptococcus anginosus group (includes S. anginosus, S. intermedius, and S. constellatus), and Enterococcus faecalis (vancomycin-susceptible isolates only). To reduce the development of drug-resistant bacteria and maintain the effectiveness of Kimyrsa and other antibacterial drugs, Kimyrsa should be used only to treat or prevent infections that are proven or strongly suspected to be caused by bacteria.	120	18 years	N/A	N/A	Y	Y		9/29/2021
Drugs	J2407	Injection, oritavancin (orbactiv), 10 mg	10 mg	10/1/2021	Orbactiv*	oritavancin for injection, for intravenous use	Indicated for the treatment of adult patients with acute bacterial skin and skin structure infections caused or suspected to be caused by susceptible isolates of designated Gram-positive microorganisms.	120	18 years	N/A	N/A	Y	Y		9/29/2021
Drugs	J2425	Injection, palifermin, 50 micrograms	50 mcg	1/1/2006	Kepivance*	palifermin injection, for intravenous use	Indicated to decrease the incidence and duration of severe oral mucositis in patients with hematologic malignancies receiving myelotoxic therapy in the setting of autologous hematopoietic stem cell support. Kepivance is indicated as supportive care for preparative regimens predicted to result in ≥ WHO Grade 3 mucositis in the majority of patients. Limitations of Use: • The safety and efficacy of Kepivance have not been established in patients with non-hematologic malignancies. • Kepivance was not effective in decreasing the incidence of severe mucositis in patients with hematologic malignancies receiving myelotoxic therapy in the setting of allogeneic hematopoietic stem cell support. • Kepivance is not recommended for use with melphalan 200 mg/m ² as a conditioning regimen.	1,008	18 years	N/A	N/A	Y	Y		4/9/2019
Drugs	J2426	Injection, paliperidone palmitate extended release (invega sustenna), 1 mg	1 mg	1/1/2011	Invega Sustenna*	paliperidone palmitate extended-release injectable suspension, for intramuscular use	Indicated for: • Treatment of schizophrenia in adults. • Treatment of schizoaffective disorder in adults as monotherapy and as an adjunct to mood stabilizers or antidepressants. Invega Trinza: Indicated for the treatment of schizophrenia in patients after they have been adequately treated with Invega Sustenna* (1-month paliperidone palmitate extended-release injectable suspension) for at least four months.	624	18 years	N/A	N/A	Y	Y		6/22/2023
Drugs	J2427	Injection, paliperidone palmitate extended release (invega hafyera, or invega trinza), 1 mg	1 mg	7/1/2023	Invega Hafyera™, Invega Trinza*	paliperidone palmitate extended-release injectable suspension, for intramuscular use	Indicated for the treatment of schizophrenia in patients after they have been adequately treated with Invega Sustenna* (1-month paliperidone palmitate extended-release injectable suspension) for at least four months.	1,560	18 years	N/A	N/A	Y	Y		6/22/2023
Drugs	J2430	Injection, pamidronate disodium, per 30 mg	30 mg	1/1/2000	Aredia*	pamidronate disodium for injection for intravenous infusion	Indicated for: • Hypercalcemia of malignancy • Paget's disease • Osteolytic bone metastases of breast cancer and osteolytic lesions of multiple myeloma	6	18 years	N/A	N/A	Y	Y		9/21/2018
Drugs	J2440	Injection, papaverine HCl, up to 60 mg	up to 60 mg	1/1/2000	N/A – various generics	papaverine hydrochloride injection, solution	Indicated in various conditions accompanied by spasm of smooth muscle, such as vascular spasm associated with acute myocardial infarction (coronary occlusion), angina pectoris, peripheral and Indicated in adults for: • Moderately emetogenic cancer chemotherapy -- prevention of acute and delayed nausea and vomiting associated with initial and repeat courses. • Highly emetogenic cancer chemotherapy -- prevention of acute nausea and vomiting associated with initial and repeat courses.	80	18 years	N/A	N/A	Y	Y		7/16/2018
Drugs	J2469	Injection, palonosetron HCl, 25 mcg	25 mcg	1/1/2005	Aloxi*	palonosetron HCl injection for intravenous use	Indicated in adults for: • Moderately emetogenic cancer chemotherapy -- prevention of acute and delayed nausea and vomiting associated with initial and repeat courses. • Highly emetogenic cancer chemotherapy -- prevention of acute nausea and vomiting associated with initial and repeat courses.	50	1 month	N/A	N/A	Y	Y		7/16/2018

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

•Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.

•11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacturer participating in the Medicaid Drug Rebate Program (MDRP).

•The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.

•The HCPCS Code effective date represents the date the HCPCS code was established

•Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacturer as they are not classified as covered outpatient drugs.

•Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/ncci-medicare/medicaid-ncci-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Drugs	J2501	Injection, paricalcitol, 1 mcg	1 mcg	1/1/2003	Zemplar*	paricalcitol injection	Indicated for the prevention and treatment of secondary hyperparathyroidism associated with stage 5 chronic kidney disease (CKD).	420	18 years	N/A	N/A	Y	Y		7/16/2018
Drugs	J2502	Injection, pasireotide long acting, 1 mg	1 mg	1/1/2016	Signifor* LAR	pasireotide for injectable suspension, for intramuscular	Indicated for the treatment of: • Patients with acromegaly who have had an inadequate response to surgery and/or for whom surgery is indicated to: - To decrease the incidence of infection, as manifested by febrile neutropenia, in patients with nonmyeloid malignancies receiving myelosuppressive anticancer drugs associated with a clinically significant incidence of febrile neutropenia. - Increase survival in patients acutely exposed to myelosuppressive doses of radiation (Hematopoietic	120	18 years	N/A	N/A	Y	Y		7/26/2018
Biologicals	J2506	Injection, pegfilgrastim, excludes biosimilar, 0.5 mg	0.5 mg	1/1/2022	Neulasta*, Neulasta* Onpro*	pegfilgrastim injection, for subcutaneous use	Indicated for the treatment of chronic gout in adult patients refractory to conventional therapy.	36	N/A	N/A	N/A	Y	Y		12/14/2021
Biologicals	J2507	Injection, pegloticase, 1 mg	1 mg	1/1/2012	Krystexxa*	pegloticase injection, for intravenous infusion	Indicated for the treatment of adults with confirmed Fabry disease.	24	18 years	N/A	N/A	Y	Y		6/4/2019
Biologicals	J2508	Injection, pegunigalsidase alfa-ixwj, 1 mg	1 mg	1/1/2024	Elfabrio*	pegunigalsidase alfa-ixwj injection, for intravenous use	Indicated in the treatment of moderately severe infections in both adults and pediatric patients due to penicillin-G-susceptible microorganisms that are susceptible to the low and persistent serum levels common to this particular dosage form. Therapy should be guided by bacteriological studies (including susceptibility tests) and by clinical response. See package insert for list of infections and microorganisms.	420	18 years	N/A	N/A	Y	Y		12/22/2023
Drugs	J2510	Injection, penicillin G procaine, aqueous, up to 600,000 units	up to 600,000 units	1/1/2000	N/A	penicillin G procaine injectable suspension	Indicated for use as: • Sedatives • Hypnotics, for the short-term treatment of insomnia, since they appear to lose their effectiveness for sleep induction and sleep maintenance after 2 weeks • Preanesthetics • Anticonvulsant, in anesthetic doses, in the emergency control of certain acute convulsive episodes, e.g., those associated with status epilepticus, cholera, eclampsia, meningitis, tetanus, and toxic reactions to strychnine or local anesthetics	52	N/A	N/A	N/A	Y	Y		8/24/2018
Drugs	J2515	Injection, pentobarbital sodium, per 50 mg	50 mg	1/1/2000	Nembutal*	pentobarbital sodium injection, USP	Indicated in the therapy of severe infections caused by penicillin G-susceptible microorganisms when rapid and high penicillin levels are required. Therapy should be guided by bacteriological studies (including susceptibility tests) and by clinical response. See package insert for full list of microorganisms.	150	N/A	N/A	N/A	Y	Y		8/24/2018
Drugs	J2540	Injection, penicillin G potassium, up to 600,000 units	600,000 units	1/1/2000	Pfizerpen*	penicillin G potassium for injection	Indicated for treatment of: • Intra-abdominal infections • Skin and skin structure infections • Female pelvic infections • Community-acquired pneumonia • Nosocomial pneumonia • Usage To reduce the development of drug-resistant bacteria and maintain the effectiveness of Zosyn and other antibacterial drugs, Zosyn should be used only to treat or prevent infections that are proven or strongly suspected to be caused by bacteria.	1,240	N/A	N/A	N/A	Y	Y		8/24/2018
Drugs	J2543	Injection, piperacillin sodium/tazobactam sodium, 1 g/0.125 g (1.125 g)	1.125 g	1/1/2000	Zosyn*	piperacillin and tazobactam for injection, for intravenous use	Indicated for the prevention of Pneumocystis jiroveci pneumonia (PJP) in high-risk, HIV-infected patients defined by one or both of the following criteria: • a history of one or more episodes of PJP • a peripheral CD4+ (T4 helper/inducer) lymphocyte count less than or equal to 200/mm3	224	2 months	N/A	N/A	Y	Y		4/10/2019
Drugs	J2545	Pentamidine isethionate, inhalation solution, FDA-approved final product, non-compounded, administered through DME, unit dose form, per 300 mg	300 mg	1/1/2000	NebuPent*	pentamidine isethionate inhalant (DME) for oral inhalation only	Indicated for the prevention of Pneumocystis jiroveci pneumonia (PJP) in high-risk, HIV-infected patients defined by one or both of the following criteria: • a history of one or more episodes of PJP • a peripheral CD4+ (T4 helper/inducer) lymphocyte count less than or equal to 200/mm3	2	16 years	N/A	N/A	Y	Y		8/24/2018

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

•Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.

•11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacturer participating in the Medicaid Drug Rebate Program (MDRP).

•The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.

•The HCPCS Code effective date represents the date the HCPCS code was established

•Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacture as they are not classified as covered outpatient drugs.

•Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/nccli-medicaid/medicaid-nccli-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Drugs	J2547	Injection, peramivir, 1 mg	1 mg	1/1/2016	Rapivab®	peramivir injection, for intravenous use	Indicated for the treatment of acute uncomplicated influenza in patients 6 months and older who have been symptomatic for no more than two days. Limitations of Use: • Efficacy based on clinical trials in which the predominant influenza virus type was influenza A; a limited number of subjects infected with influenza B virus were enrolled. • Consider available information on influenza drug susceptibility patterns and treatment effects when deciding whether to use. • Efficacy could not be established in patients with serious influenza requiring hospitalization.	600	6 months	N/A	N/A	Y	Y		2/25/2021
Drugs	J2550	Injection, promethazine HCl, up to 50 mg	up to 50 mg	1/1/2000	Phenergan	promethazine hydrochloride injection	Indicated for the following conditions: • Amelioration of allergic reactions to blood or plasma. • In anaphylaxis as an adjunct to epinephrine and other standard measures after the acute symptoms have been controlled.	93	2 years	N/A	N/A	Y	Y		8/24/2018
Drugs	J2560	Injection, phenobarbital sodium, up to 120 mg	up to 120 mg	1/1/2000	N/A	phenobarbital sodium injection	Indicated for use as: • Sedative. Sedation is obtainable within an hour, and in adequate dosage, the duration of action is more than six hours. Included in the more common conditions in which the sedative action of this class of drugs is desired are anxiety-tension states, hyperthyroidism, essential hypertension, nausea and vomiting of functional origin, motion sickness, acute labyrinthitis, pylorospasm in infants, chorea and cardiac failure. Phenobarbital is also a useful adjunct in treatment of hemorrhage from the respiratory or gastrointestinal tract. Phenobarbital controls anxiety, decreases muscular activity and lessens nervous excitability in hyperthyroid patients. However, thyrotoxic individuals occasionally react poorly to barbiturates. • Hypnotic, for the short-term treatment of insomnia, since it appears to lose its effectiveness for sleep induction and sleep maintenance after 2 weeks. • Preanesthetic. • Long-term anticonvulsant, (phenobarbital, mephobarbital and metharbital) for the treatment of	N/A	N/A	N/A	N/A	Y	Y		8/29/2018
Drugs	J2562	Injection, plerixafor, 1 mg	1 mg	1/1/2010	Mozobil®	plerixafor injection, solution for subcutaneous use	Indicated in combination with granulocyte-colony stimulating factor (G-CSF) to mobilize hematopoietic stem cells (HSCs) to the peripheral blood for collection and subsequent autologous transplantation in patients with non-Hodgkin's lymphoma and multiple myeloma.	160	18 years	N/A	N/A	Y	Y		6/6/2019
Drugs	J2590	Injection, oxytocin, up to 10 units	up to 10 units	1/1/2000	Pitocin®	oxytocin injection, USP synthetic	Indicated for: • Antepartum - The initiation or improvement of uterine contractions, where there is desirable and considered suitable	12	N/A	N/A	Females Only	Y	Y		7/16/2018
Drugs	J2597	Injection, desmopressin acetate, per 1 mcg	1 mcg	1/1/2000	DDAVP®	desmopressin acetate injection	Indicated for patients with hemophilia A with factor VIII coagulant activity levels greater than 5%, patients with mild to moderate classic von Willebrand's disease (Type 1) with factor VIII levels greater than 5%, as an antidiuretic replacement therapy in the management of central (cranial) diabetes insipidus and for the management of the temporary polyuria and polydipsia following head trauma or surgery in the pituitary region. DDAVP is ineffective for the treatment of nephrogenic diabetes insipidus.	660	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	Indication age specific: Hemophilia A and von Willebrand's Disease: 3 months of age and older Diabetes Insipidus: 12 years of	7/2/2018
Drugs	J2675	Injection, progesterone, per 50 mg	per 50 mg	1/1/2003	N/A	progesterone injection, in sesame oil for intramuscular use only	Indicated in amenorrhea and abnormal uterine bleeding caused by hormonal imbalance in the absence of organic pathology, such as submucous fibroids or uterine cancer.	2	18 years	N/A	Females Only	Y	Y		6/6/2019
Drugs	J2679	Injection, fluphenazine hcl, 1.25 mg	1.25 mg	1/1/2024	N/A	fluphenazine hydrochloride injection, solution	Fluphenazine Hydrochloride Injection, USP is indicated in the management of manifestations of psychotic disorders. • Fluphenazine hydrochloride has not been shown effective in the management of behavioral complications in patients with mental retardation.	248	18 years	N/A	N/A	Y	Y		12/22/2023
Drugs	J2680	Injection, fluphenazine decanoate, up to 25 mg	up to 25 mg	1/1/2000	N/A	fluphenazine decanoate injection	intended for use in the management of patients requiring prolonged parenteral neuroleptic therapy (e.g. chronic schizophrenics). Fluphenazine decanoate has not been shown effective in the management of behavioral complications in patients with mental retardation.	8	12 years	N/A	N/A	Y	Y		6/4/2019
Drugs	J2690	Injection, procainamide HCl, up to 1 g	up to 1 g	1/1/2000	N/A	procainamide hydrochloride injection, solution	Indicated for the treatment of documented ventricular arrhythmias, such as sustained ventricular tachycardia, that, in the judgment of the physician, are life-threatening. Because of the proarrhythmic effects of procainamide, its use with lesser arrhythmias is generally not recommended. Treatment of	7	18 years	N/A	N/A	Y	Y		6/6/2019
Drugs	J2700	Injection, oxacillin sodium, up to 250 mg	up to 250 mg	1/1/2000	N/A, various generics	oxacillin sodium injection, powder, for solution for intramuscular or intravenous	Indicated for the treatment of infections caused by penicillinase-producing staphylococci which have demonstrated susceptibility to the drug. Cultures and susceptibility tests should be performed initially to determine the causative organism and their susceptibility to the drug.	744	N/A	N/A	N/A	Y	Y		9/21/2018
Drugs	J2710	Injection, neostigmine methylsulfate, up to 0.5 mg	up to 0.5 mg	1/1/2000	Bloxiverz®	neostigmine methylsulfate injection, for intravenous use	Indicated for the reversal of the effects of nondepolarizing neuromuscular blocking agents (NMBAs) after surgery.	50	N/A	N/A	N/A	Y	Y		4/10/2019

North Carolina Division of Health Benefits Physician Administered Drug Program Catalog

- Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.
- 11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacturer participating in the Medicaid Drug Rebate Program (MDRP).
- The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.
- The HCPCS Code effective date represents the date the HCPCS code was established
- Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacturer as they are not classified as covered outpatient drugs.
- Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/nccli-medicaid/medicaid-nccli-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Drugs	J2720	Injection, protamine sulfate, per 10 mg	10 mg	1/1/2000	N/A	protamine sulfate injection, solution for intravenous use	Indicated for the treatment of heparin overdose.	5	18 years	N/A	N/A	Y	Y		8/29/2018
Biologics	J2724	Injection, protein C concentrate, intravenous, human, 10 IU	10 IU	1/1/2008	Ceprotrin	protein c concentrate (human) lyophilized power for solution for injection	Indicated for pediatric and adult patients with severe congenital Protein C deficiency for the prevention and treatment of venous thrombosis and purpura fulminans.	105,840	N/A	N/A	N/A	Y	Y		6/4/2019
Drugs	J2730	Injection, pralidoxime chloride, up to 1 g	up to 1 g	1/1/2000	Protopam®	pralidoxime chloride for injection	Indicated as an antidote: • In the treatment of poisoning caused by those pesticides and chemicals of the organophosphate class which have anticholinesterase activity. • In the control of overdose by anticholinesterase drugs used in the treatment of myasthenia gravis.	20	N/A	N/A	N/A	Y	Y		8/24/2018
Drugs	J2760	Injection, phentolamine mesylate, up to 5 mg	up to 5 mg	1/1/2000	Regitine®	phentolamine mesylate injection, powder, lyophilized, for suspension	Indicated for: • The prevention or control of hypertensive episodes that may occur in a patient with pheochromocytoma as a result of stress or manipulation during preoperative preparation and surgical excision. • The prevention or treatment of dermal necrosis and sloughing following intravenous administration or	372	N/A	N/A	N/A	Y	Y		8/24/2018
Drugs	J2765	Injection, metoclopramide HCl, up to 10 mg	up to 10 mg	1/1/2000	N/A	metoclopramide hydrochloride injection	Indicated for: • The relief of symptoms associated with acute and recurrent diabetic gastric stasis • The prophylaxis of vomiting associated with emetogenic cancer chemotherapy • The prophylaxis of postoperative nausea and vomiting in those circumstances where nasogastric suction is undesirable • Facilitating small bowel intubation in adults and pediatric patients in whom the tube does not pass the	560	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	Indication specific: • Facilitating Small Bowel Intubation: 18 years of age and older • All other indications: None	6/6/2019
Biologics	J2777	Injection, faricimab-svoa, 0.1 mg	0.1 mg	10/1/2022	Vabysmo®	faricimab-svoa injection, for intravitreal use	Indicated for the treatment of patients with: • Neovascular (Wet) Age-Related Macular Degeneration (nAMD) • Diabetic Macular Edema (DME)	240	18 years	N/A	N/A	Y	Y		12/1/2023
Biologics	J2778	Injection, ranibizumab, 0.1 mg	0.1 mg	1/1/2008	Lucentis®	ranibizumab injection for intravitreal injection	Indicated for the treatment of patients with: • Neovascular (Wet) Age-Related Macular Degeneration (AMD)	20	18 years	N/A	N/A	Y	Y		10/31/2018
Biologics	J2779	Injection, ranibizumab, via intravitreal implant (susvimo), 0.1 mg	0.1 mg	1/1/2002	Susvimo™	ranibizumab injection for intravitreal use via ocular implant	Indicated for the treatment of patients with Neovascular (wet) Age-related Macular Degeneration (AMD) who have previously responded to at least two intravitreal injections of a VEGF inhibitor.	100	18 years	N/A	N/A	Y	Y		6/6/2022
Drugs	J2781	Injection, pegcetacoplan, intravitreal, 1 mg	1 mg	10/1/2023	Syfovre™	pegcetacoplan injection, for intravitreal use	Indicated for the treatment of geographic atrophy (GA) secondary to age-related macular degeneration (AMD).	60	18 years	N/A	N/A	Y	Y		9/28/2023
Drugs	J2782	Injection, avacincaptad pegol, 0.1 mg	0.1 mg	4/1/2024	Izervay™	avacincaptad pegol intravitreal solution	Indicated for the treatment of geographic atrophy (GA) secondary to age-related macular degeneration (AMD).	80	18 years	N/A	N/A	Y	Y		4/12/2024
Biologics	J2783	Injection, rasburicase, 0.5 mg	0.5 mg	1/1/2004	Elitek®	rasburicase for injection, for intravenous use	Indicated for the initial management of plasma uric acid levels in pediatric and adult patients with leukemia, lymphoma, and solid tumor malignancies who are receiving anti-cancer therapy expected to result in tumor lysis and subsequent elevation of plasma uric acid. Limitation of Use: Elitek is indicated for a single course of treatment.	280	N/A	N/A	N/A	Y	Y		6/4/2019
Drugs	J2785	Injection, regadenoson, 0.1 mg	0.1 mg	1/1/2009	Lexiscan®	regadenoson injection for intravenous use	Indicated for radionuclide myocardial perfusion imaging (MPI) in patients unable to undergo adequate exercise stress.	4	18 years	N/A	N/A	Y	Y		6/4/2021
Biologics	J2786	Injection, reslizumab, 1 mg	1 mg	1/1/2017	Cinqair®	reslizumab injection, for intravenous use	Indicated for add-on maintenance treatment of patients with severe asthma aged 18 years and older, and with an eosinophilic phenotype. Limitations of Use: Cinqair is not indicated for:	840	18 years	N/A	N/A	Y	Y		7/2/2018

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

•Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.

•11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacture participating in the Medicaid Drug Rebate Program (MDRP).

•The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.

•The HCPCS Code effective date represents the date the HCPCS code was established

•Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacture as they are not classified as covered outpatient drugs.

•Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/nccli-medicaid/medicaid-nccli-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Immune Globulins	J2788	Injection, Rho d immune globulin, human, minidose, 50 micrograms (250 IU)	50 mcg	1/1/2003	HyperRHO® S/D Mini Dose, MICRhoGAM®	rho(D) immune globulin (human), mini dose	HyperRHO S/D Mini Dose: recommended to prevent the isoimmunization of Rho(D) negative women at the time of spontaneous or induced abortion of up to 12 weeks' gestation provided the following criteria are met: 1. The mother must be Rho(D) negative and must not already be sensitized to the Rho(D) antigen. 2. The father is not known to be Rho(D) negative. 3. Gestation is not more than 12 weeks at termination. **See package insert for full usage criteria.** MICRhoGAM: For use in preventing Rh immunization. • Pregnancy and other obstetrical conditions in Rh-negative women unless the father or baby are conclusively Rh-negative, e.g. delivery of an Rh-positive baby irrespective of the ABO groups of the mother	1	N/A	N/A	HyperRHO: Females Only	Y	Y		7/3/2018
Immune Globulins	J2790	Injection, Rho d immune globulin, human, full dose, 300 micrograms (1500 IU)	300 mcg (1500 IU)	1/1/2003	HyperRho® S/D Full Dose, RhoGAM®	rho(d) immune globulin (human), full dose	Indicated for use in preventing Rh immunization: • In pregnancy and other obstetrical conditions (see full prescribing information). • In any Rh-negative person after incompatible transfusion of Rh-positive blood or blood products.	3	N/A	N/A	N/A	Y	Y		4/9/2022
Immune Globulins	J2791	Injection, Rho(D) immune globulin (human), (Rhophylac), intramuscular or intravenous, 100 IU	100 IU	1/1/2008	Rhophylac®	rho(d) immune globulin intravenous (human) 1500 IU (300 mcg) solution for intravenous (IV) or intramuscular (IM) injection	Indicated for: Suppression of Rhesus (Rh) Isoimmunization in: • Pregnancy and obstetrical conditions in non-sensitized, Rho (D)-negative women with an Rh-incompatible pregnancy, including: -Routine antepartum and postpartum Rh prophylaxis -Rh prophylaxis in obstetric complications or invasive procedures • Incompatible transfusions in Rho (D)-negative individuals transfused with blood components containing Rho (D)-positive red blood cells (RBCs).	350	N/A	N/A	N/A	Y	Y	12/2023: Age restrictions updated to align with other rho(D) immune globulin products effective 12/20/2023.	1/26/2024
Immune Globulins	J2792	Injection, rho D immune globulin, intravenous, human, solvent detergent, 100 IU	100 IU	1/1/2000	WinRho SDF®	rho(D) immune globulin intravenous (human) solution for intravenous or intramuscular injection	Indicated for: Immune Thrombocytopenic Purpura (ITP) Raising platelet counts in Rho(D) positive, non-splenectomized: • Children with chronic or acute ITP,	1,500	N/A	N/A	N/A	Y	Y		9/12/2018
Biologics	J2793	Injection, rilonacept, 1 mg	1 mg	1/1/2010	Arcalyst®	rilonacept injection for subcutaneous use	Indicated for: • the treatment of patients with Cryopyrin-Associated Periodic Syndromes (CAPS), including Familial Cold Autoinflammatory Syndrome (FACS) and Muckle-Wells Syndrome (MWS) in adults and children 12 years of age and older. • maintenance of remission of Deficiency of Interleukin-1 Receptor Antagonist (DIRA) in adults and pediatric patients weighing at least 10 kg.	1,600	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	Indication specific age restrictions: CAPS and RP: 12 years of age and older DIRA: N/A	4/26/2021
Drugs	J2794	Injection, risperidone (risperdal consta), 0.5 mg	0.5 mg	1/1/2005	Risperdal Consta®	risperidone long-acting injection	Indicated: • For the treatment of schizophrenia. • as monotherapy or as adjunctive therapy to lithium or valproate for the maintenance treatment of Bipolar I Disorder.	300	N/A	N/A	N/A	Y	Y		10/3/2019
Drugs	J2795	Injection, ropivacaine hydrochloride, 1 mg	1 mg	1/1/2001	Naropin®	ropivacaine HCl injection	Indicated for the production of local or regional anesthesia for surgery and for acute pain management. Surgical Anesthesia: epidural block for surgery including cesarean section; major nerve block; local infiltration. Acute pain management: epidural continuous infusion or intermittent bolus, eg, postoperative or labor; Indicated for the treatment of thrombocytopenia in: • Adult patients with immune thrombocytopenia (ITP) who have had an insufficient response to corticosteroids, immunoglobulins, or splenectomy. • Pediatric patients 1 year of age and older with ITP for at least 6 months who have had an insufficient response to corticosteroids, immunoglobulins, or splenectomy.	2,166	18 years	N/A	N/A	Y	Y		8/29/2018
Drugs	J2796	Injection, romiplostim, 10 micrograms	10 mcg	1/1/2010	Nplate®	romiplostim for injection, for subcutaneous use	Indicated for the treatment of thrombocytopenia in: • Adult patients with immune thrombocytopenia (ITP) who have had an insufficient response to corticosteroids, immunoglobulins, or splenectomy. • Pediatric patients 1 year of age and older with ITP for at least 6 months who have had an insufficient response to corticosteroids, immunoglobulins, or splenectomy.	700	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	Indication Specific Age Restrictions: ITP: 1 year of age and older HS-ARS: None	2/25/2021
Drugs	J2798	Injection, risperidone, (perseris), 0.5 mg	0.5 mg	10/1/2019	Perseris™	risperidone for extended-release injectable suspension, for subcutaneous use	Indicated for the treatment of schizophrenia in adults.	480	18 years	N/A	N/A	Y	Y		10/3/2019
Drugs	J2799	Injection, risperidone (uzedy), 1 mg	1 mg	1/1/2024	Uzedy™	risperidone extended-release injectable suspension, for subcutaneous use	Indicated for the treatment of schizophrenia in adults.	250	18 years	N/A	N/A	Y	Y		12/22/2023

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

•Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.

•11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacturer participating in the Medicaid Drug Rebate Program (MDRP).

•The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.

•The HCPCS Code effective date represents the date the HCPCS code was established

•Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacturer as they are not classified as covered outpatient drugs.

•Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/ncci-medicare/medicaid-ncci-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Drugs	J2800	Injection, methocarbamol, up to 10 mL	up to 10 mL	1/1/2000	Robaxin®	methocarbamol injection for intravenous or intramuscular use	Indicated as an adjunct to rest, physical therapy, and other measures for the relief of discomfort associated with acute, painful, musculoskeletal conditions; supportive therapy in tetanus.	54	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	Indication specific. Relief of discomfort associated with acute, painful, musculoskeletal conditions: 18 years of age and older. Tetanus: None	6/8/2019
Drugs	J2801	Injection, risperidone (rykindo), 0.5 mg	0.5 mg	4/1/2024	Rykindo®	risperidone for extended-release injectable suspension.	Risperidone for extended-release injectable suspension is indicated: • For the treatment of schizophrenia in adults.	300	18 years	N/A	N/A	Y	Y		4/12/2024
Biologicals	J2820	Injection, sargramostim (GM-CSF), 50 mcg	50 mcg	1/1/2000	Leukine®	sargramostim injection, for subcutaneous or intravenous use	Indicated: • To shorten time to neutrophil recovery and to reduce the incidence of severe and life-threatening infections and infections resulting in death following induction chemotherapy in adult patients 55 years and older with acute myeloid leukemia (AML). • For the mobilization of hematopoietic progenitor cells into peripheral blood for collection by	620	Indication Specific Age Restrictions (see comments)	Indication Specific (see comments)	N/A	Y	Y	Indication specific age restrictions: • To shorten time to neutrophil recovery and to reduce the incidence of severe	8/29/2018
Biologicals	J2840	Injection, sebelipase alfa, 1 mg	1 mg	1/1/2017	Kanuma®	sebelipase alfa injection, for intravenous use	Indicated for the treatment of patients with a diagnosis of Lysosomal Acid Lipase (LAL) deficiency.	1,260	1 month	N/A	N/A	Y	Y		12/16/2021
Biologicals	J2860	Injection, siltuximab, 10 mg	10 mg	1/1/2016	Sylvant®	siltuximab for injection, for intravenous use	Indicated for treatment of patients with multicentric Castleman's disease (MCD) who are human immunodeficiency virus (HIV) negative and human herpesvirus-8 (HHV-8) negative. Limitations of Use: Sylvant was not studied in patients with MCD who are HIV positive or HHV-8 positive because Sylvant did not bind to virally produced IL-6 in a non-clinical study.	400	18 years	N/A	N/A	Y	Y		6/7/2019
Drugs	J2916	Injection, sodium ferric gluconate complex in sucrose injection, 12.5 mg	12.5 mg	1/1/2003	Ferrlecit®	sodium ferric gluconate complex in sucrose injection, for intravenous (IV) use	Indicated for the treatment of iron deficiency anemia in patients 6 years of age and older with chronic kidney disease receiving hemodialysis who are receiving supplemental epoetin therapy.	80	6 years	N/A	N/A	Y	Y		9/21/2018
Drugs	J2919	Injection, methylprednisolone sodium succinate, 5 mg	5 mg	4/1/2024	Solu-Medrol®	methylprednisolone sodium succinate for injection	When oral therapy is not feasible, and the strength, dosage form, and route of administration of the drug reasonably lend the preparation to the treatment of the condition, the intravenous or intramuscular use of Solu-Medrol is indicated as follows: • Allergic states: Control of severe or incapacitating allergic conditions intractable to adequate trials of conventional treatment in asthma, atopic dermatitis, contact dermatitis, drug hypersensitivity reactions, serum sickness, transfusion reactions. • Dermatologic diseases: Bullous dermatitis herpetiformis, exfoliative erythroderma, mycosis fungoides, pemphigus, severe erythema multiforme (Stevens-Johnson syndrome). • Endocrine disorders: Primary or secondary adrenocortical insufficiency (hydrocortisone or cortisone is	4,500	N/A	N/A	N/A	Y	Y		3/22/2024

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

- Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.
- 11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacture participating in the Medicaid Drug Rebate Program (MDRP).
- The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.
- The HCPCS Code effective date represents the date the HCPCS code was established
- Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacture as they are not classified as covered outpatient drugs.
- Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/nccli-medicaid/medicaid-nccli-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Biologics	J2993	Injection, reteplase, 18.1 mg	18.1 mg	1/1/2002	Retavase®	reteplase for injection, for intravenous use	Indicated for treatment of acute ST-elevation myocardial infarction (STEMI) to reduce the risk of death and heart failure. Limitation of Use: The risk of stroke may outweigh the benefit produced by thrombolytic therapy in	2	18 years	N/A	N/A	Y	Y		10/31/2018
Biologics	J2997	Injection, alteplase recombinant, 1 mg	1 mg	1/1/2001	Activase®, Cathflo® Activase®	alteplase for injection, for intravenous use	Cathflo Activase: Indicated for the restoration of function to central venous access devices as assessed by the ability to withdraw blood. Activase: Indicated for the treatment of: • Acute Ischemic Stroke (AIS) • Acute Myocardial Infarction (AMI) to reduce mortality and incidence of heart failure. Limitation of use in AMI: The risk of stroke may be greater than the benefit in patients at low risk of death from cardiac causes. • Acute Massive Pulmonary Embolism (PE) for lysis.	3,100	18 years	N/A	N/A	Y	Y	1/2024: Category corrected from Drugs to Biologics.	1/26/2024
Biologics	J2998	Injection, plasminogen, human-tvmh, 1 mg	1 mg	1/1/2002	Ryplazim®	plasminogen, human-tvmh lyophilized powder for reconstitution, for intravenous use	Indicated for the treatment of patients with plasminogen deficiency type 1 (hypoplasminogenemia).	15,411.2	11 months	N/A	N/A	Y	Y		6/6/2022
Drugs	J3000	Injection, streptomycin, up to 1 gram	up to 1 g	1/1/2000	N/A	streptomycin for injection for intramuscular use	Indicated for the treatment of individuals with moderate to severe infections caused by susceptible strains of microorganisms in the specific conditions of Mycobacterium tuberculosis and Non-tuberculosis infections: Mycobacterium tuberculosis, and other sensitive non tuberculosis pathogens including Pasteurella pestis (plague); Francisella tularensis (tularemia); Brucella; Calymmatobacterium granulomatis (donovanosis, granuloma inguinale); H. ducreyi (chancroid); H. influenzae (in respiratory, endocardial, and meningial infections, concomitantly with another antibacterial agent); K. pneumoniae pneumonia (concomitantly with another antibacterial agent); E. coli, Proteus, A. aerogenes, K. pneumoniae, and	62	N/A	N/A	N/A	Y	Y		6/7/2019
Drugs	J3010	Injection, fentanyl citrate, 0.1 mg	0.1 mg	1/1/2000	N/A	fentanyl citrate injection, for intravenous or intramuscular use	Indicated for: • analgesic action of short duration during the anesthetic periods, premedication, induction and maintenance, and in the immediate postoperative period (recovery room) as the need arises. • use as an opioid analgesic supplement in general or regional anesthesia. • administration with a neuroleptic as an anesthetic premedication, for the induction of anesthesia and as an adjunct in the maintenance of general and regional anesthesia. • use as an anesthetic agent with oxygen in selected high risk patients, such as those undergoing open heart surgery or certain complicated neurological or orthopedic procedures.	210	2 years	N/A	N/A	Y	Y		6/4/2019
Drugs	J3030	Injection, sumatriptan, succinate, 6 mg	6 mg	1/1/2000	Imitrex®	sumatriptan succinate injection, for subcutaneous use	Indicated for: • Acute treatment of migraine with or without aura in adults • Acute treatment of cluster headache in adults	8	18 years	N/A	N/A	Y	Y		9/21/2018
Biologics	J3055	Injection, talquetamab-tgvs, 0.25 mg	0.25 mg	4/1/2024	Talvey™	talquetamab-tgvs injection, for subcutaneous use	Indicated for the treatment of adult patients with relapsed or refractory multiple myeloma who have received at least four prior lines of therapy, including a proteasome inhibitor, an immunomodulatory agent and an anti-CD38 monoclonal antibody.	1,808	18 years	N/A	N/A	Y	Y		4/12/2024
Biologics	J3060	Injection, taliglucerase alfa, 10 units	10 units	1/1/2014	Elelyso®	taliglucerase alfa for injection, for intravenous use	Indicated for the treatment of patients with a confirmed diagnosis of Type 1 Gaucher disease.	2,520	4 years	N/A	N/A	Y	Y		6/4/2019
Drugs	J3090	Injection, tedizolid phosphate, 1 mg	1 mg	1/1/2016	Sivextro®	tedizolid phosphate for injection, for intravenous use	Indicated in adults and pediatric patients 12 years of age and older for the treatment of acute bacterial skin and skin structure infections (ABSSSI) caused by designated susceptible bacteria. Indicated for the treatment of the following infections in adult patients caused by designated susceptible bacteria: • Complicated skin and skin structure infections (cSSSI) • Hospital-acquired and ventilator-associated bacterial pneumonia (HABP/VABP) caused by susceptible isolates of Staphylococcus aureus. Vibativ should be reserved for use when alternative treatments are not suitable.	1,200	12 years	N/A	N/A	Y	Y		7/28/2020
Drugs	J3095	Injection, telavancin, 10 mg	10 mg	1/1/2011	Vibativ®	telavancin for injection, for intravenous use	Indicated for the prevention and reversal of bronchospasm in patients 12 years of age and older with asthma and reversible bronchospasm associated with bronchitis and emphysema.	3,150	18 years	N/A	N/A	Y	Y		6/8/2019
Drugs	J3105	Injection, terbutaline sulfate, up to 1 mg	up to 1 mg	1/1/2000	N/A	terbutaline sulfate injection, solution	Indicated for the prevention and reversal of bronchospasm in patients 12 years of age and older with asthma and reversible bronchospasm associated with bronchitis and emphysema.	45	12 years	N/A	N/A	Y	Y		9/12/2018
Biologics	J3111	Injection, romosozumab-aqqg, 1 mg	1 mg	10/1/2019	Evenity™	romosozumab-aqqg injection, for subcutaneous use	Indicated for the treatment of osteoporosis in postmenopausal women at high risk for fracture, defined as a history of osteoporotic fracture, or multiple risk factors for fracture; or patients who have failed or are intolerant to other available osteoporosis therapy. Limitations of Use: Limit duration of use to 12 monthly doses. If osteoporosis therapy remains warranted, continued therapy with an anti-resorptive agent should be considered	420	Not for use in premenopausal women.	N/A	Females Only	Y	Y		10/3/2019

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

- Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.
- 11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacturer participating in the Medicaid Drug Rebate Program (MDRP).
- The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.
- The HCPCS Code effective date represents the date the HCPCS code was established
- Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacturer as they are not classified as covered outpatient drugs.
- Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/ncci-medicaid/medicaid-nci-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Drugs	J3121	Injection, testosterone enanthate, 1 mg	1 mg	1/1/2015	N/A	testosterone enanthate injection, solution	Indicated for replacement therapy in conditions associated with a deficiency or absence of endogenous testosterone including primary hypogonadism (congenital or acquired), hypogonadotropic hypogonadism (congenital or acquired), and delayed puberty. Testosterone Enanthate injection may be used secondarily in women with advancing inoperable metastatic (skeletal) mammary cancer who are 1 – 5 years postmenopausal.	1,200	N/A	N/A	N/A	Y	Y		9/12/2018
Drugs	J3145	Injection, testosterone undecanoate, 1mg	1 mg	1/1/2015	Aveed*	testosterone undecanoate injection for intramuscular use	Indicated for testosterone replacement therapy in adult males for conditions associated with a deficiency or absence of endogenous testosterone: primary hypogonadism (congenital or acquired) or hypogonadotropic hypogonadism (congenital or acquired). Limitations of Use: • Safety and efficacy of Aveed in men with "age-related hypogonadism" have not been established. • Safety and efficacy of Aveed in males less than 18 years old have not been established.	1,500	18 years	N/A	Males Only	Y	Y		9/21/2018
Drugs	J3230	Injection, chlorpromazine HCl, up to 50 mg	50 mg	1/1/2000	N/A	chlorpromazine hydrochloride injection	Indicated for the treatment of schizophrenia; to control nausea and vomiting; for relief of restlessness and apprehension before surgery; for acute intermittent porphyria; as an adjunct in the treatment of tetanus; to control the manifestations of the manic type of manic-depressive illness; for relief of intractable	248	6 months	N/A	N/A	Y	Y		9/27/2018
Biologicals	J3240	Injection, thyrotropin alfa, 0.9 mg, provided in 1.1 mg vial	0.9 mg	1/1/2003	Thyrogen®	thyrotropin alfa for injection, for intramuscular use	Indicated for: • Diagnostic: Use as an adjunctive diagnostic tool for serum thyroglobulin (Tg) testing with or without radioiodine imaging in the follow-up of patients with well-differentiated thyroid cancer who have previously undergone thyroidectomy. • Ablation: Use as an adjunctive treatment for radioiodine ablation of thyroid tissue remnants in patients who have undergone a near-total or total thyroidectomy for well-differentiated thyroid cancer and who do not have evidence of distant metastatic thyroid cancer.	2	18 years	N/A	N/A	Y	Y		6/19/2023
Biologicals	J3241	Injection, teprotumumab-trbw, 10 mg	10 mg	10/1/2020	Tepezza®	teprotumumab-trbw for injection, for intravenous use	Indicated for the treatment of Thyroid Eye Disease regardless of Thyroid Eye Disease activity or duration.	600	18 years	N/A	N/A	Y	Y		5/25/2023
Drugs	J3243	Injection, tigecycline, 1 mg	1 mg	1/1/2007	Tygacil®	tigecycline for injection, for intravenous use	Indicated in patients 18 years of age and older for: • Complicated skin and skin structure infections • Complicated intra-abdominal infections Indicated in patients 18 years of age and older for: • Complicated skin and skin structure infections • Complicated intra-abdominal infections • Community-acquired bacterial pneumonia Limitations of Use: Tigecycline for injection is not indicated for treatment of diabetic foot infection or hospital-acquired pneumonia, including ventilator-associated pneumonia. To reduce the development of drug-resistant bacteria and maintain the effectiveness of tigecycline for injection and other antibacterial drugs, Tigecycline for injection should be used only to treat infections that are proven or strongly suspected to be caused by bacteria.	1,450	18 years	N/A	N/A	Y	Y		9/21/2018
Drugs	J3244	Injection, tigecycline (accord) not therapeutically equivalent to j3243, 1 mg	1 mg	1/1/2023	N/A	tigecycline for injection, for intravenous use (Accord)	Indicated in patients 18 years of age and older for: • Complicated skin and skin structure infections • Complicated intra-abdominal infections • Community-acquired bacterial pneumonia Limitations of Use: Tigecycline for injection is not indicated for treatment of diabetic foot infection or hospital-acquired pneumonia, including ventilator-associated pneumonia. To reduce the development of drug-resistant bacteria and maintain the effectiveness of tigecycline for injection and other antibacterial drugs, Tigecycline for injection should be used only to treat infections that are proven or strongly suspected to be caused by bacteria.	1,450	18 years	N/A	N/A	Y	Y		12/12/2022
Drugs	J3250	Injection, trimethobenzamide HCl, up to 200 mg	up to 200 mg	1/1/2000	Tigan®	trimethobenzamide hydrochloride	Indicated for the treatment of postoperative nausea and vomiting and for nausea associated with gastroenteritis.	124	18 years	N/A	N/A	Y	Y		9/12/2018
Drugs	J3260	Injection, tobramycin sulfate, up to 80 mg	up to 80 mg	1/1/2000	N/A	tobramycin sulfate injection	Indicated for the treatment of serious bacterial infections caused by susceptible strains of the designated microorganisms in the diseases listed below: • Septicemia in the neonate, child, and adult caused by P. aeruginosa, E. coli, and Klebsiella sp	558	N/A	N/A	N/A	Y	Y		9/12/2018

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

- Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.
- 11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacturer participating in the Medicaid Drug Rebate Program (MDRP).
- The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.
- The HCPCS Code effective date represents the date the HCPCS code was established
- Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacture as they are not classified as covered outpatient drugs.
- Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/nc-ci-medicaid/medicaid-nc-ci-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Biologics	J3262	Injection, tocilizumab, 1 mg	1 mg	1/1/2011	Actemra®	tocilizumab injection, for intravenous use	Indicated for the treatment of: • Adult patients with moderately to severely active rheumatoid arthritis (RA) who have had an inadequate response to one or more Disease-Modifying Anti-Rheumatic Drugs (DMARDs).	3,200	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	Indication specific age restrictions: • 2 years of age and older:	3/17/2022
Drugs	J3285	Injection, treprostinil, 1 mg	1 mg	1/1/2006	Remodulin®	treprostinil injection, for subcutaneous or intravenous use	Indicated for treatment of pulmonary arterial hypertension (PAH) (WHO Group 1) to diminish symptoms associated with exercise and to reduce the rate of clinical deterioration in patients requiring transition from epoprostenol.	1,813	17 years	N/A	N/A	Y	Y		5/14/2019
Drugs	J3299	Injection, triamcinolone acetonide (xipere), 1 mg	1 mg	1/1/2000	Xipere™	triamcinolone acetonide injectable suspension, for suprachoroidal use	Indicated for the treatment of macular edema associated with uveitis.	80	18 years	N/A	N/A	Y	Y		6/6/2022
Drugs	J3300	Injection, triamcinolone acetonide, preservative free, 1 mg	1 mg	1/1/2009	Triesence®	triamcinolone acetonide injectable suspension	Indicated for: • Treatment of the following ophthalmic diseases: sympathetic ophthalmia, temporal arteritis, uveitis, and ocular inflammatory conditions unresponsive to topical corticosteroids. • Visualization during vitrectomy	8	N/A	N/A	N/A	Y	Y		6/7/2019
Drugs	J3301	Injection, triamcinolone acetonide, Not Otherwise Specified, per 10 mg	10 mg	1/1/2000	Kenalog-10®, Kenalog-40®	triamcinolone acetonide injectable suspension, for intra-articular or intralesional use only	Kenalog-40 Indicated for intramuscular use as follows: • Allergic states: Control of severe or incapacitating allergic conditions intractable to adequate trials of conventional treatment in asthma, atopic dermatitis, contact dermatitis, drug hypersensitivity reactions, perennial or seasonal allergic rhinitis, serum sickness, transfusion reactions. • Dermatologic diseases: Bullous dermatitis herpetiformis, exfoliative erythroderma, mycosis fungoides,	150	N/A	N/A	N/A	Y	Y		9/12/2018
Drugs	J3304	Injection, triamcinolone acetonide, preservative-free, extended-release, microsphere formulation, 1 mg	1 mg	1/1/2019	Zilretta™	triamcinolone acetonide extended-release injectable suspension, for intra-articular use	Indicated as an intra-articular injection for the management of osteoarthritis pain of the knee. Limitation of Use: Zilretta is not intended for repeat administration.	64	18 years	N/A	N/A	Y	Y		9/12/2018
Drugs	J3315	Injection, triptorelin pamoate, 3.75 mg	3.75 mg	1/1/2003	Trelstar®	triptorelin pamoate for injectable suspension	Indicated for the palliative treatment of advanced prostate cancer.	6	18 years	N/A	Males Only	Y	Y		2/19/2024
Drugs	J3316	Injection, triptorelin, extended-release, 3.75 mg	3.75 mg	1/1/2019	Triptodur™	triptorelin for extended-release injectable suspension, for intramuscular use	Indicated for the treatment of pediatric patients 2 years and older with central precocious puberty.	6	2 years	N/A	N/A	Y	Y		2/19/2024

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

•Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.

•11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacture participating in the Medicaid Drug Rebate Program (MDRP).

•The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.

•The HCPCS Code effective date represents the date the HCPCS code was established

•Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacture as they are not classified as covered outpatient drugs.

•Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/ncci-medicaid/medicaid-ncci-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Biologics	J3357	Ustekinumab, for subcutaneous injection, 1 mg	1 mg	1/1/2017	Stelara® for subcutaneous use	ustekinumab injection, for subcutaneous use	Indicated for the treatment of: Adult patients with: • Moderate to severe plaque psoriasis (Ps) who are candidates for phototherapy or systemic therapy • Active psoriatic arthritis (PsA) • Moderately to severely active Crohn's disease (CD) • Moderately to severely active ulcerative colitis Pediatric patients 6 to 17 years of age with: • Moderate to severe plaque psoriasis, who are candidates for phototherapy or systemic therapy • Active psoriatic arthritis (PsA)	180	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	Indication specific age restrictions. • 6 years of age and older: plaque psoriasis (Ps), psoriatic arthritis (PsA) • 18 years of age and older: Crohn's disease (CD), ulcerative colitis	8/16/2022
Biologics	J3358	Ustekinumab, for intravenous injection, 1 mg	1 mg	1/1/2018	Stelara® for intravenous use	ustekinumab injection, for intravenous use	Indicated for the treatment of adult patients with: • Moderately to severely active Crohn's disease (CD) • Moderately to severely active ulcerative colitis	520	18 years	N/A	N/A	Y	Y		12/3/2019
Drugs	J3360	Injection, diazepam, up to 5 mg	up to 5 mg	1/1/2000	N/A	diazepam injection	Indicated: • For the management of anxiety disorders or for the short-term relief of the symptoms of anxiety. Anxiety or tension associated with the stress of everyday life usually does not require treatment with an anxiolytic. • In acute alcohol withdrawal, diazepam may be useful in the symptomatic relief of acute agitation, tremor, impending or acute delirium tremens and hallucinosis. • As an adjunct prior to endoscopic procedures if apprehension, anxiety or acute stress reactions are present, and to diminish the patient's recall of the procedures. • As a useful adjunct for the relief of skeletal muscle spasm due to reflex spasm to local pathology (such as inflammation of the muscles or joints, or secondary to trauma); spasticity caused by upper motor neuron disorders (such as cerebral palsy and paraplegia); athetosis; stiff-man syndrome; and tetanus.	250	31 days	N/A	N/A	Y	Y		10/10/2018
Drugs	J3370	Injection, vancomycin HCl, 500 mg	500 mg	1/1/2000	N/A	vancomycin hydrochloride for injection, USP for intravenous use	Indicated for the treatment of serious or severe infections caused by susceptible strains of methicillin-resistant (β-lactam-resistant) staphylococci. It is indicated for penicillin-allergic patients, for patients who cannot receive or who have failed to respond to other drugs, including the penicillins or cephalosporins, and for infections caused by vancomycin-susceptible organisms that are resistant to other antimicrobial drugs. Vancomycin hydrochloride for injection is indicated for initial therapy when methicillin-resistant staphylococci are suspected, but after susceptibility data are available, therapy should be adjusted accordingly. To reduce the development of drug-resistant bacteria and maintain the effectiveness of vancomycin hydrochloride for injection USP and other antibacterial drugs, vancomycin hydrochloride for injection should be used only to treat or prevent infections that are proven or strongly suspected to be caused by susceptible bacteria. When culture and susceptibility information are available, they should be considered in selecting or modifying antibacterial therapy. In the absence of such data, local epidemiology and susceptibility patterns may contribute to the empiric selection of therapy. See package insert for list of infections.	124	N/A	N/A	N/A	Y	Y		6/8/2019
Drugs	J3371	Injection, vancomycin hcl (mylan), not therapeutically equivalent to J3370, 500 mg	500 mg	1/1/2023	N/A	vancomycin hydrochloride for injection, for intravenous use (Mylan)	Indicated in adult and pediatric patients (neonates and older) for the treatment of: • Septicemia • Infective Endocarditis • Skin and Skin Structure Infections • Bone Infections • Lower Respiratory Tract Infections	124	N/A	N/A	N/A	Y	Y		12/6/2022
Drugs	J3372	Injection, vancomycin hcl (xellia), not therapeutically equivalent to J3370, 500 mg	500 mg	1/1/2023	N/A	vancomycin injection, for intravenous use (Xellia)	Indicated in adult and pediatric patients less than 18 years of age as follows: • Vancomycin Injection administered intravenously is indicated for the treatment of: • Septicemia	124	N/A	N/A	N/A	Y	Y		12/6/2022

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

•Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.

•11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacturer participating in the Medicaid Drug Rebate Program (MDRP).

•The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.

•The HCPCS Code effective date represents the date the HCPCS code was established

•Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacturer as they are not classified as covered outpatient drugs.

•Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/ncci-medicare/medicaid-ncci-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Biologicals	J3380	Injection, vedolizumab, intravenous, 1 mg	1 mg	1/1/2016	Entyvio®	vedolizumab for injection, for intravenous use	Indicated in adults for the treatment of: • moderately to severely active ulcerative colitis (UC). • moderately to severely active Crohn's disease (CD).	600	18 years	N/A	N/A	Y	Y	4/2024: Subcutaneous formulation removed from coverage effective 3/31/2024 due to HCPCS code description change effective 4/1/2024.	3/22/2024
Biologicals	J3385	Injection, velaglucerase alfa, 100 units	100 units	1/1/2011	VPRIV®	velaglucerase alfa for injection, for intravenous use	Indicated for long-term enzyme replacement therapy (ERT) for patients with type 1 Gaucher disease.	252	4 years	N/A	N/A	Y	Y		6/8/2019
Drugs	J3396	Injection, verteporfin, 0.1 mg	0.1 mg	1/1/2005	Visudyne®	verteporfin for injection, for intravenous use	Indicated for the treatment of patients with predominantly classic subfoveal choroidal neovascularization due to age-related macular degeneration, pathologic myopia or presumed ocular histoplasmosis.	150	18 years	N/A	N/A	Y	Y		9/12/2018
Biologicals	J3397	Injection, vestronidase alfa-vjkb, 1 mg	1 mg	1/1/2019	Mepsevii™	vestronidase alfa-vjkb injection, for intravenous use	Indicated in pediatric and adult patients for the treatment of Mucopolysaccharidosis VII (MPS VII, Sly syndrome). Limitations of Use: The effect of Mepsevii on the central nervous system manifestations of MPS VII has not been determined.	1,680	N/A	N/A	N/A	Y	Y		8/5/2021
Biologicals	J3398	Injection, voretigene neparovvec-rzyl, 1 billion vector genomes	1 billion vector genomes (vg)	1/1/2019	Luxturna™	voretigene neparovvec-rzyl intraocular suspension for subretinal injection	Indicated for the treatment of patients with confirmed biallelic RPE65 mutation-associated retinal dystrophy. Patients must have viable retinal cells as determined by the treating physician(s).	300	1 year	N/A	N/A	Y	Y		9/17/2021
Biologicals	J3401	Beremagene geperpavec-svdt for topical administration, containing nominal 5 x 10 ⁹ pfu/ml vector genomes, per 0.1 ml	0.1 mL	1/1/2024	Vyjuvek™	beremagene geperpavec-svdt biological suspension mixed with excipient gel for topical application	Indicated for the treatment of wounds in patients 6 months of age and older with dystrophic epidermolysis bullosa with mutation(s) in the collagen type VII alpha 1 chain (COL7A1) gene.	125	6 months	N/A	N/A	Y	Y		12/22/2023
Drugs	J3410	Injection, hydroxyzine HCl, up to 25 mg	up to 25 mg	1/1/2000	Vistaril®	hydroxyzine hydrochloride injection for intramuscular use	• The total management of anxiety, tension, and psychomotor agitation in conditions of emotional stress requires in most instances a combined approach of psychotherapy and chemotherapy. Hydroxyzine has been found to be particularly useful for this latter phase of therapy in its ability to render the disturbed patient more amenable to psychotherapy in long term treatment of the psychoneurotic and psychotic, although it should not be used as the sole treatment of psychosis or of clearly demonstrated cases of depression. • Also useful in alleviating the manifestations of anxiety and tension as in the preparation for dental procedures and in acute emotional problems. It has also been recommended for the management of anxiety associated with organic disturbances and as adjunctive therapy in alcoholism and allergic conditions with strong emotional overlay, such as in asthma, chronic urticaria, and pruritus. • Hydroxyzine hydrochloride intramuscular solution is useful in treating the following types of patients	240	N/A	N/A	N/A	Y	Y		10/26/2018
Drugs	J3420	Injection, vitamin B-12 cyanocobalamin, up to 1,000 mcg	up to 1,000 mcg	1/1/2000	N/A	cyanocobalamin injection, USP (vitamin B-12)	Indicated for vitamin B12 deficiencies due to malabsorption which may be associated with the following conditions: • Addisonian (pernicious) anemia • Gastrointestinal pathology, dysfunction, or surgery, including gluten enteropathy or sprue, small bowel Indicated in the following coagulation disorders which are due to faulty formation of factors II, VII, IX and X when caused by vitamin K deficiency or interference with vitamin K activity: • anticoagulant-induced prothrombin deficiency caused by coumarin or indanedione derivatives; • prophylaxis and therapy of hemorrhagic disease of the newborn; • hypoprothrombinemia due to antibacterial therapy; • hypoprothrombinemia secondary to factors limiting absorption or synthesis of vitamin K, e.g., obstructive jaundice, biliary fistula, sprue, ulcerative colitis, celiac disease, intestinal resection, cystic fibrosis of the pancreas, and regional enteritis; • other drug-induced hypoprothrombinemia where it is definitely shown that the result is due to	10	N/A	N/A	N/A	Y	Y		9/27/2018
Drugs	J3430	Injection, phytanadione (vitamin K) per 1 mg	1 mg	1/1/2000	Mephyton®	phytanadione injectable emulsion, USP	Indicated in the following coagulation disorders which are due to faulty formation of factors II, VII, IX and X when caused by vitamin K deficiency or interference with vitamin K activity: • anticoagulant-induced prothrombin deficiency caused by coumarin or indanedione derivatives; • prophylaxis and therapy of hemorrhagic disease of the newborn; • hypoprothrombinemia due to antibacterial therapy; • hypoprothrombinemia secondary to factors limiting absorption or synthesis of vitamin K, e.g., obstructive jaundice, biliary fistula, sprue, ulcerative colitis, celiac disease, intestinal resection, cystic fibrosis of the pancreas, and regional enteritis; • other drug-induced hypoprothrombinemia where it is definitely shown that the result is due to	50	N/A	N/A	N/A	Y	Y		6/5/2019

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

•Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.

•11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacturer participating in the Medicaid Drug Rebate Program (MDRP).

•The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.

•The HCPCS Code effective date represents the date the HCPCS code was established

•Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacture as they are not classified as covered outpatient drugs.

•Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/nccli-medicaid/medicaid-nccli-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Biologics	J3470	Injection, hyaluronidase, up to 150 units	up to 150 units	1/1/2000	Amphadase®	hyaluronidase injection	Indicated as an adjuvant: • In subcutaneous fluid administration for achieving hydration. • To increase absorption and dispersion of other injected drugs. • In subcutaneous urography for improving resorption of radiopaque agents.	93	N/A	N/A	N/A	Y	Y		6/19/2023
Biologics	J3473	Injection, hyaluronidase, recombinant, 1 USP unit	1 USP unit	1/1/2007	Hylenex® Recombinant	hyaluronidase human injection, for infiltration use.	Indicated as an: • Adjuvant to increase the dispersion and absorption of other injected drugs.	2,250	N/A	N/A	N/A	Y	Y		6/19/2023
Drugs	J3475	Injection, magnesium sulfate, per 500 mg	500 mg	1/1/2000	N/A	magnesium sulfate injection	Indicated for replacement therapy in magnesium deficiency, especially in acute hypomagnesemia accompanied by signs of tetany similar to those observed in hypocalcemia. In such cases, the serum magnesium level is usually below the lower limit of normal (1.5 to 2.5 mEq/L) and the serum calcium level	560	N/A	N/A	N/A	Y	Y		6/5/2019
Drugs	J3480	Injection, potassium chloride, per 2 mEq	2 mEq	1/1/2000	N/A	potassium chloride injection	Indicated for the treatment or prevention of hypokalemia when oral treatment is not feasible.	1,240	N/A	N/A	N/A	Y	Y		8/24/2018
Drugs	J3486	Injection, ziprasidone mesylate, 10 mg	10 mg	1/1/2004	Geodon®	ziprasidone mesylate for injection, for intramuscular use	Indicated for the acute treatment of agitation in schizophrenic patients.	124	18 years	N/A	N/A	Y	Y		3/17/2022
Drugs	J3489	Injection, zoledronic acid, 1 mg	1 mg	1/1/2014	Reclast®; Zometa®	zoledronic acid injection, for intravenous use	Reclast is indicated for: • Treatment and prevention of postmenopausal osteoporosis • Treatment to increase bone mass in men with osteoporosis • Treatment and prevention of glucocorticoid-induced osteoporosis • Treatment of Paget's disease of bone in men and women Limitations of Use: Optimal duration of use has not been determined. For patients at low-risk for fracture, consider drug discontinuation after 3 to 5 years of use. Zometa is indicated for the treatment of: • Hypercalcemia of malignancy. • Patients with multiple myeloma and patients with documented bone metastases from solid tumors, in conjunction with standard antineoplastic therapy. Prostate cancer should have progressed after treatment with at least one hormonal therapy. Limitations of Use: The safety and efficacy of Zometa has not been established for use in hyperparathyroidism or non-tumor-related hypercalcemia.	20	18 years	N/A	N/A	Y	Y		9/21/2018
Drugs	J3490	Unclassified drugs	1 mg	1/1/2000	Aponvie™	aprepitant injectable emulsion, for intravenous use	Indicated for the prevention of postoperative nausea and vomiting (PONV) in adults. Limitations of Use: Aponvie has not been studied for treatment of established nausea and vomiting.	160	18 years	N/A	N/A	Y	Y		3/16/2023
Drugs	J3490	Unclassified drugs	1 mg	1/1/2000	Baxdela™	delafloxacin for injection, for intravenous use	Indicated in adults for the treatment of acute bacterial skin and skin structure infections (ABSSSI) caused by susceptible isolates of the following:	8,400	18 years	N/A	N/A	Y	Y		12/3/2019
Drugs	J3490	Unclassified drugs	1 mg	1/1/2000	Bludigo™	indigotindisulfonate sodium injection, for intravenous use	Indicated for use as a visualization aid in the cystoscopic assessment of the integrity of the ureters in adults following urological and gynecological open, robotic, or endoscopic surgical procedures.	40	18 years	N/A	N/A	Y	Y		10/20/2022
Drugs	J3490	Unclassified drugs	1 mg	1/1/2000	Bridion®	sugammadex injection, for intravenous use	Indicated for the reversal of neuromuscular blockade induced by rocuronium bromide and vecuronium bromide in adults undergoing surgery.	12,500	18 years	N/A	N/A	Y	Y		11/14/2019
Drugs	J3490	Unclassified drugs	1 mg	1/1/2000	Cleviprex®	clevitidine injectable emulsion, for intravenous use	Indicated for the reduction of blood pressure when oral therapy is not feasible or not desirable.	1,500	18 years	N/A	N/A	Y	Y		10/4/2018
Drugs	J3490	Unclassified drugs	1 mL	1/1/2000	Defitelio®	defibrotide sodium injection, for intravenous use	Indicated for the treatment of adult and pediatric patients with hepatic veno-occlusive disease (VOD), also known as sinusoidal obstruction syndrome (SOS), with renal or pulmonary dysfunction following hematopoietic stem-cell transplantation (HSCT).	1,395	18 years	N/A	N/A	Y	Y		6/10/2019
Drugs	J3490	Unclassified drugs	1 mg	1/1/2000	Depacon®	valproate sodium, for intravenous injection	Indicated as an intravenous alternative in patients in whom oral administration of valproate products is temporarily not feasible in the following conditions: • Monotherapy and adjunctive therapy of complex partial seizures and simple and complex absence seizures; adjunctive therapy in patients with multiple seizure types that include absence seizures.	119,000	2 years	N/A	N/A	Y	Y		5/30/2019

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

- Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.
- 11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacture participating in the Medicaid Drug Rebate Program (MDRP).
- The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.
- The HCPCS Code effective date represents the date the HCPCS code was established
- Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacture as they are not classified as covered outpatient drugs.
- Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/ncci-medicare/medicaid-ncci-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Drugs	J3490	Unclassified drugs	1 mg lidocaine USP base	1/1/2000	Lidocaine (various topical formulations)	lidocaine (various topical formulations)	Indicated for production of anesthesia of accessible mucous membranes of the oropharynx. It is also useful as an anesthetic lubricant for intubation and for the temporary relief of pain associated with minor burns, including sunburn, abrasions of the skin, and insect bites.	31,000	N/A	N/A	N/A	Y	Y		10/26/2018
Drugs	J3490	Unclassified drugs	1 mg	1/1/2000	N/A	nalmefene hydrochloride injection	Indicated: - for the complete or partial reversal of opioid drug effects, including respiratory depression, induced by either natural or synthetic opioids - in the management of known or suspected opioid overdose	20	18 years	N/A	N/A	Y	Y	12/2023: Due to NDA product Revex no longer being marketed, recommended dosing updated to align with ANDA product Prescribing Information and brand name Revex updated to N/A effective 6/22/2022.	1/26/2024
Drugs	J3490	Unclassified drugs	50 mL	1/1/2000	N/A	sodium bicarbonate injection, solution	Indicated in: • The treatment of metabolic acidosis which may occur in severe renal disease, uncontrolled diabetes, circulatory insufficiency due to shock or severe dehydration, extracorporeal circulation of blood, cardiac. Indicated for the prophylaxis of invasive Aspergillus and Candida infections in patients who are at high risk of developing these infections due to being severely immunocompromised, such as HSCT recipients with GVHD or those with hematologic malignancies with prolonged neutropenia from chemotherapy. Indicated for the treatment of invasive aspergillosis in adults and pediatric patients 13 years of age and older.	403	N/A	N/A	N/A	Y	Y		10/31/2018
Drugs	J3490	Unclassified drugs	1 mg	1/1/2000	Noxafi®	posaconazole injection, for intravenous use	Indicated for the prophylaxis of invasive Aspergillus and Candida infections in patients who are at high risk of developing these infections due to being severely immunocompromised, such as HSCT recipients with GVHD or those with hematologic malignancies with prolonged neutropenia from chemotherapy. Indicated for the treatment of invasive aspergillosis in adults and pediatric patients 13 years of age and older.	9,600	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	Indication specific age restrictions: Prophylaxis of invasive Aspergillus and Candida infections: 2 years of age and	7/27/2021
Drugs	J3490	Unclassified drugs	1 mg	1/1/2000	Opvee®	nalmefene nasal spray	Indicated for the emergency treatment of known or suspected overdose induced by natural or synthetic opioids in adults and pediatric patients aged 12 years and older, as manifested by respiratory and/or central nervous system depression.	27	12 years	N/A	N/A	Y	Y		10/26/2023
Drugs	J3490	Unclassified drugs	1 vial	1/1/2000	Prevymis®	letermovir injection, for intravenous use	Indicated for: - prophylaxis of cytomegalovirus (CMV) infection and disease in adult CMV-seropositive recipients [R+] of an allogeneic hematopoietic stem cell transplant (HSCT). - prophylaxis of CMV disease in adult kidney transplant recipients at high risk (Donor CMV seropositive/Recipient CMV seronegative [D+/R-]).	31	18 years	N/A	N/A	Y	Y		7/26/2023
Drugs	J3490	Unclassified drugs	1 mL	1/1/2000	Proyavblue®	methylene blue injection, for intravenous use	Indicated for the treatment of pediatric and adult patients with acquired methemoglobinemia. This indication is approved under accelerated approval. Continued approval for this indication may be contingent upon verification of clinical benefit in subsequent trials.	60	N/A	N/A	N/A	Y	Y		3/17/2022
Drugs	J3490	Unclassified drugs	10 mg	1/1/2000	Revatio®	sildenafil injection, for intravenous use	Indicated for the treatment of pulmonary arterial hypertension (PAH) (WHO Group I) in adults to improve exercise ability and delay clinical worsening. Studies establishing effectiveness were short-term (12 to 16 weeks), and included predominately patients with NYHA Functional Class II-III symptoms. Etiologies were idiopathic (71%) or associated with connective tissue disease (25%).	93	3 years	N/A	N/A	Y	Y		3/17/2022
Drugs	J3490	Unclassified drugs	1 mg	1/1/2000	Rezipres®	ephedrine hydrochloride injection, for intravenous use	Indicated for the treatment of clinically important hypotension occurring in the setting of anesthesia.	1,457	18 years	N/A	N/A	Y	Y		4/17/2022
Drugs	J3490	Unclassified drugs	1 mcg	1/1/2000	Upravi®	selexipag for injection, for intravenous use	Indicated for the treatment of pulmonary arterial hypertension (PAH, WHO Group I) to delay disease progression and reduce the risk of hospitalization for PAH.	111,600	18 years	N/A	N/A	Y	Y		9/28/2021
Drugs	J3490	Unclassified drugs	10 mg	1/1/2000	Vimpat®	lacosamide injection, for intravenous use	Vimpat is indicated for: • Treatment of partial-onset seizures in patients 1 month of age and older. • Adjunctive therapy in the treatment of primary generalized tonic-clonic seizures in patients 4 years of age and older.	1,240	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	Indication specific age restrictions: Partial-onset seizures: 1 month of age and older Primary generalized tonic-clonic seizures: 4 years of age and older	11/17/2021
Drugs	J3490	Unclassified drugs	0.6 mg	1/1/2000	Zegalogue®	dasiglucagon injection, for subcutaneous use	Indicated for the treatment of severe hypoglycemia in pediatric and adult patients with diabetes aged 6 years and above.	10	6 years	N/A	N/A	Y	Y		7/27/2021
Drugs	J3490	Unclassified drugs	1 mcg	1/1/2000	iDose® TR	travoprost intracameral implant, for intracameral administration	Travoprost intracameral implant is indicated for the reduction of intraocular pressure (IOP) in patients with open-angle glaucoma (OAG) or ocular hypertension (OHT).	150	18 years	N/A	N/A	Y	Y		2/27/2024

North Carolina Division of Health Benefits Physician Administered Drug Program Catalog

- Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.
- 11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacturer participating in the Medicaid Drug Rebate Program (MDRP).
- The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.
- The HCPCS Code effective date represents the date the HCPCS code was established
- Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacture as they are not classified as covered outpatient drugs.
- Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/nccli-medicaid/medicaid-nccli-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Drugs	J3490	Unclassified drugs	1 mg	1/1/2000	Wainua™	epilontersen injection, for subcutaneous use	Epilontersen injection is indicated for the treatment of the polyneuropathy of hereditary transthyretin-mediated amyloidosis in adults.	45	18 years	N/A	N/A	Y	Y		3/25/2024
Biologics	J3590	Unclassified biologics	1 IU	1/1/2002	Adzyna	ADAMTS13, recombinant-krhn lyophilized powder for injection, for intravenous use	ADAMTS13, recombinant-krhn lyophilized powder for injection is indicated for prophylactic or on demand enzyme replacement therapy (ERT) in adult and pediatric patients with congenital thrombotic thrombocytopenic purpura (cTTP).	30,000	2 years	N/A	N/A	Y	Y		12/21/2023
Biologics	J3590	Unclassified biologics	11 mg (1 kit)	1/1/2002	Cabivi®	caplacizumab-yhdp for injection, for intravenous or subcutaneous use	Indicated for the treatment of adult patients with acquired thrombotic thrombocytopenic purpura (aTTP), in combination with plasma exchange and immunosuppressive therapy.	32	18 years	N/A	N/A	Y	Y		3/26/2019
Biologics	J3590	Unclassified biologics	1 mg	1/1/2002	Cosentyx®	secukinumab injection, for intravenous use	Secukinumab intravenous injection is indicated for the treatment of: • Adults with active psoriatic arthritis (PsA) • Adults with active ankylosing spondylitis (AS). • Adults with active non-radiographic axial spondyloarthritis (nr-axSpA) with objective signs of inflammation.	1,125	18 years	N/A	N/A	Y	Y	3/2024: Removal of subcutaneous formulations from PADP effective 3/31/2024 per DHB request 3/20/2024.	4/12/2024
Biologics	J3590	Unclassified biologics	1 mg	1/1/2002	Orvoh™	mirikizumab-mrzk injection, for intravenous or subcutaneous use	Mirikizumab-mrzk injection is indicated for the treatment of moderately to severely active ulcerative colitis in adults.	600	18 years	N/A	N/A	Y	Y		12/1/2023
Biologics	J3590	Unclassified biologics	per daily dose	1/1/2002	Palforzia™	peanut (Arachis hypogaea) allergen powder-drfp powder for oral administration	Indicated for the mitigation of allergic reactions, including anaphylaxis, that may occur with accidental exposure to peanut. Limitation of Use: Not indicated for the emergency treatment of allergic reactions, including anaphylaxis.	31	4 years	N/A	N/A	Y	Y	Initial Dose Escalation may be administered to patients aged 4 through 17 years. Up-Dosing and Maintenance may be	4/29/2020
Biologics	J3590	Unclassified biologics	0.5 mL	1/1/2002	Plegridy™	peginterferon beta-1a injection, for subcutaneous or intramuscular use	Indicated for the treatment of patients with relapsing forms of multiple sclerosis.	3	18 years	N/A	N/A	Y	Y		2/25/2021
Biologics	J3590	Unclassified biologics	50 mL	1/1/2002	Praxbind®	idarucizumab injection, for intravenous use	Indicated in patients treated with Pradaxa when reversal of the anticoagulant effects of dabigatran is needed: • For emergency surgery/urgent procedures • In life-threatening or uncontrolled bleeding	4	18 years	N/A	N/A	Y	Y		7/16/2018
Biologics	J3590	Unclassified biologics	1 IU	1/1/2002	Recothrom®	thrombin topical (recombinant) lyophilized powder for solution - for elapagademase-ivlr injection, for intramuscular use	Indicated to aid hemostasis whenever oozing blood and minor bleeding from capillaries and small venules is accessible and control of bleeding by standard surgical techniques is ineffective or impractical in adults and pediatric populations greater than or equal to one month of age.	80,000	1 month	N/A	N/A	Y	Y		4/10/2019
Biologics	J3590	Unclassified biologics	1 mg	1/1/2002	Revcomi™	elapagademase-ivlr injection, for intramuscular use	Indicated for the treatment of adenosine deaminase severe combined immune deficiency (ADA-SCID) in pediatric and adult patients.	288	N/A	N/A	N/A	Y	Y		12/28/2018
Biologics	J3590	Unclassified biologics	1 mg	1/1/2002	Strensiq®	asfotase alfa injection, for subcutaneous use	Treatment of patients with perinatal/infantile-onset and juvenile-onset hypophosphatasia (HPP).	5,460	N/A	N/A	N/A	Y	Y		4/10/2019
Biologics	J3590	Unclassified biologics	1 mcg	1/1/2002	Sylatron™	peginterferon alfa-2b for injection, for subcutaneous use	Indicated for the adjuvant treatment of melanoma with microscopic or gross nodal involvement within 84 days of definitive surgical resection including complete lymphadenectomy.	4,500	18 years	N/A	N/A	Y	Y		6/7/2019
Biologics	J3590	Unclassified biologics	1 mg	1/1/2002	Tyenne®	tocilizumab-aazg injection, for intravenous use	Tocilizumab-aazg injection is indicated for treatment of: - Rheumatoid Arthritis (RA) • Adult patients with moderately to severely active rheumatoid arthritis who have had an inadequate response to one or more Disease-Modifying Anti-Rheumatic Drugs (DMARDs). - Giant Cell Arteritis (GCA) • Adults with moderate to severe active disease. Indicated as a source of water and electrolytes. Also indicated for use as a priming solution in hemodialysis procedures.	1,600	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	Indication specific: RA, GCA: 18 years of age and older PIJA, SJIA: 2 years of age and older	5/3/2024
Drugs	J7030	Infusion, normal saline solution, 1,000 cc	1,000 cc	1/1/2000	N/A	normal saline solution 1,000 cc (sodium chloride injection)	Indicated as a source of water and electrolytes. Also indicated for use as a priming solution in hemodialysis procedures.	N/A	N/A	N/A	N/A	Y	Y		10/26/2018
Drugs	J7040	Infusion, normal saline solution, sterile	500 mL	1/1/2000	N/A	normal saline solution 500 cc (sodium chloride injection)	Indicated as a source of water and electrolytes. Also indicated for use as a priming solution in hemodialysis procedures.	186	N/A	N/A	N/A	Y	Y		6/7/2019
Drugs	J7042	5% Dextrose/normal saline (500 mL = 1 unit)	500 mL	1/1/2000	N/A	dextrose 5% / normal saline	Indicated for use in adults and pediatric patients as sources of calories and water for hydration.	200	N/A	N/A	N/A	Y	Y		10/10/2018
Drugs	J7050	Infusion, normal saline solution, 250 cc	250 cc	1/1/2000	N/A	normal saline solution 250 cc (sodium chloride injection)	Indicated as a source of water and electrolytes. Also indicated for use as a priming solution in hemodialysis procedures.	186	N/A	N/A	N/A	Y	Y		6/7/2019
Drugs	J7060	5% Dextrose/water (500 mL = 1 unit)	500 mL	1/1/2000	N/A	dextrose 5% / water	Indicated for use in adults and pediatric patients as sources of calories and water for hydration.	200	N/A	N/A	N/A	Y	Y		10/10/2018
Drugs	J7070	Infusion, D5W, 1,000 cc	1,000 cc	1/1/2000	N/A	D5W (dextrose injection)	Indicated for parenteral replenishment of fluid and minimal carbohydrate calories as required by clinical condition of the patient.	124	N/A	N/A	N/A	Y	Y		10/4/2018
Drugs	J7120	Ringer's lactate infusion, up to 1,000 cc	up to 1,000 cc	1/1/2000	N/A	lactated ringer's infusion	Indicated as a source of water and electrolytes or as an alkalinizing agent.	124	N/A	N/A	N/A	Y	Y		8/29/2018
Drugs	J7121	5% dextrose in lactated ringers infusion, up to 1,000 cc	up to 1,000 cc	1/1/2016	N/A	D5LR (5% dextrose in lactated ringer's injection)	Indicated for parenteral replacement of extracellular losses of fluid and electrolytes, with or without minimal carbohydrate calories, as required by the clinical condition of the patient.	124	N/A	N/A	N/A	Y	Y		10/4/2018

North Carolina Division of Health Benefits Physician Administered Drug Program Catalog

- Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.
- 11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacturer participating in the Medicaid Drug Rebate Program (MDRP).
- The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.
- The HCPCS Code effective date represents the date the HCPCS code was established
- Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacturer as they are not classified as covered outpatient drugs.
- Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/ncci-medicaid/medicaid-ncci-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Biologicals	J7165	Injection, prothrombin complex concentrate, human-lans, per i.u. of factor ix activity	1 IU	4/1/2024	Balfaxar*	prothrombin complex concentrate, human-lans lyophilized powder for solution, for intravenous use	Prothrombin complex concentrate, human-lans is a blood coagulation factor replacement product indicated for the urgent reversal of acquired coagulation factor deficiency induced by Vitamin K antagonist (VKA, e.g., warfarin) therapy in adult patients with need for an urgent surgery/invasive procedure.	5,000	18 years	N/A	N/A	Y	Y		3/22/2024
Biologicals	J7168	Prothrombin complex concentrate (human), kcentra, per i.u. of factor ix activity	1 IU	7/1/2021	Kcentra*	prothrombin complex concentrate (human) for intravenous use, lyophilized	Indicated for the urgent reversal of acquired coagulation factor deficiency induced by Vitamin K antagonist (VKA, e.g., warfarin) therapy in adult patients with acute major bleeding or need for an urgent surgery/invasive procedure.	5,000	18 years	N/A	N/A	Y	Y		6/28/2021
Biologicals	J7169	Injection, coagulation factor xa (recombinant), inactivated-zhzo (andexxa), 10 mg	10 mg	7/1/2020	Andexxa*	coagulation factor Xa (recombinant), inactivated-zhzo lyophilized powder for solution for intravenous injection	Indicated for patients treated with rivaroxaban and apixaban, when reversal of anticoagulation is needed due to life-threatening or uncontrolled bleeding.	180	18 years	N/A	N/A	Y	Y		6/17/2020
Biologicals	J7170	Injection, emicizumab-kxwh, 0.5 mg	0.5 mg	1/1/2019	Hemlibra*	emicizumab-kxwh injection, for subcutaneous use	Indicated for routine prophylaxis to prevent or reduce the frequency of bleeding episodes in adult and pediatric patients ages newborn and older with hemophilia A (congenital factor VIII deficiency) with or without factor VIII inhibitors.	5,040	N/A	N/A	N/A	Y	Y		7/2/2018
Biologicals	J7175	Injection, factor X, (human), 1 IU	1 IU	1/1/2017	Coagadex*	coagulation factor X (human) lyophilized powder for solution for intravenous injection	Indicated in adults and children with hereditary Factor X deficiency for: • On-demand treatment and control of bleeding episodes • Perioperative management of bleeding in patients with mild, moderate and severe hereditary Factor X deficiency	84,000	N/A	N/A	N/A	Y	Y		5/25/2023
Biologicals	J7177	Injection, human fibrinogen concentrate (fibruga), 1 mg	1 mg	1/1/2019	Fibryga*	fibrinogen (human) lyophilized powder for reconstitution, for intravenous use	Indicated for the treatment of acute bleeding episodes in adults and children with congenital fibrinogen deficiency, including afibrinogenemia and hypofibrinogenemia. Fibryga is not indicated for dysfibrinogenemia.	9,800	N/A	N/A	N/A	Y	Y		11/29/2021
Biologicals	J7178	Injection, human fibrinogen concentrate, not otherwise specified, 1 mg	1 mg	1/1/2013	RiaSTAP*	fibrinogen concentrate (human) for intravenous use, lyophilized powder for	Indicated for the treatment of acute bleeding episodes in patients with congenital fibrinogen deficiency, including afibrinogenemia and hypofibrinogenemia.	9,800	N/A	N/A	N/A	Y	Y		6/8/2019
Biologicals	J7179	Injection, Von Willebrand factor (recombinant), (Vonvendil), 1IU VWF:RCO	1 IU	1/1/2017	Vonvendil*	von Willebrand factor (recombinant) lyophilized powder for solution, for intravenous injection	Indicated for use in adults (age 18 and older) diagnosed with von Willebrand disease (VWD) for: • On-demand treatment and control of bleeding episodes. • Perioperative management of bleeding. • Routine prophylaxis to reduce the frequency of bleeding episodes in patients with severe Type 3 von	254,800	18 years	N/A	N/A	Y	Y		2/11/2022
Biologicals	J7180	Injection, factor XIII (antihemophilic factor, human), 1 IU	1 IU	1/1/2012	Corifact	factor XIII concentrate (human) injection for intravenous use	Indicated for adult and pediatric patients with congenital Factor XIII deficiency for: • Routine prophylactic treatment • Peri-operative management of surgical bleeding.	10,000	N/A	N/A	N/A	Y	Y		10/10/2018
Biologicals	J7181	Injection, factor XIII A-subunit, (recombinant), per IU	per IU	1/1/2015	Tretten*	coagulation factor XIII a-subunit (recombinant)	Indicated for routine prophylaxis of bleeding in patients with congenital factor XIII A-subunit deficiency. Not for use in patients with congenital factor XIII B-subunit deficiency.	9,800	N/A	N/A	N/A	Y	Y		6/8/2019
Biologicals	J7182	Injection, factor VIII, (antihemophilic factor, recombinant), (Novoeight), per IU	1 IU	1/1/2015	Novoeight*	antihemophilic factor (recombinant) for intravenous injection lyophilized powder for solution	Adults and children with hemophilia A for: Control and prevention of bleeding; Perioperative management; Routine prophylaxis to prevent or reduce the frequency of bleeding episodes.	168,000	N/A	N/A	N/A	Y	Y		6/6/2019
Biologicals	J7183	Injection, Von Willebrand factor complex (human), Wilate, 1 IU VWF:RCO	1 IU VWF:RCO	1/1/2012	Wilate*	von willebrand factor/coagulation factor VIII complex (human) lyophilized powder for solution for intravenous injection	Von Willebrand disease: Indicated in children and adults with von Willebrand disease for: • On-demand treatment and control of bleeding episodes. • Perioperative management of bleeding. • Routine prophylaxis to reduce the frequency of bleeding episodes. - Wilate is indicated for routine prophylaxis in children 6 years of age and older and adults with von Willebrand disease. Hemophilia A: Indicated in adolescents and adults with hemophilia A for:	90,000	N/A	N/A	N/A	Y	Y		2/16/2024

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

- Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.
- 11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacturer participating in the Medicaid Drug Rebate Program (MDRP).
- The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.
- The HCPCS Code effective date represents the date the HCPCS code was established
- Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacture as they are not classified as covered outpatient drugs.
- Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/ncci-medicaid/medicaid-ncci-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Biologics	J7185	Injection, factor VIII (antihemophilic factor, recombinant) (Xyntha), per IU	1 IU	1/1/2010	Xyntha*	factor VIII (antihemophilic factor, recombinant) for intravenous injection	<ul style="list-style-type: none"> • Indicated in adults and children with hemophilia A for control and prevention of bleeding episodes and for perioperative management. • Indicated in adults and children with hemophilia A for routine prophylaxis to reduce the frequency of bleeding episodes. • Xyntha is not indicated in patients with von Willebrand's disease. 	58,800	N/A	N/A	N/A	Y	Y		9/21/2020
Biologics	J7186	Injection, antihemophilic factor VIII/Von Willebrand factor complex (human), per factor VIII IU	1 IU	1/1/2009	Alphanate*	antihemophilic factor/von Willebrand factor complex (human) lyophilized powder for solution for intravenous injection	<p>Indicated for:</p> <ul style="list-style-type: none"> • Control and prevention of bleeding in adult and pediatric patients with hemophilia A. • Surgical and/or invasive procedures in adult and pediatric patients with von Willebrand Disease in whom desmopressin (DDAVP) is either ineffective or contraindicated. It is not indicated for patients with severe VWD [Type 3] undergoing major surgery. 	133,250	N/A	N/A	N/A	Y	Y	Max Units: Although the monthly dose can exceed this amount, use of higher doses administered by a provider must be supported with	9/21/2018
Biologics	J7187	Injection, Von Willebrand factor complex (Humate-P), per IU, VWF:RCO	1 IU	1/1/2007	Humate-P*	antihemophilic factor/von Willebrand factor complex (human), lyophilized powder for reconstitution for intravenous use only	<p>Indicated for:</p> <ul style="list-style-type: none"> • Hemophilia A – Treatment and prevention of bleeding in adults. • Von Willebrand disease (VWD) – in adults and pediatric patients in the (1) Treatment of spontaneous and trauma-induced bleeding episodes, and (2) Prevention of excessive bleeding during and after surgery. This applies to patients with severe VWD as well as patients with mild to moderate VWD where the use of desmopressin is known or suspected to be inadequate. Humate-P is not indicated for the prophylaxis of 	136,250	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	Indication specific age restrictions: • Hemophilia A: 18 years of age and older • Von Willebrand disease (VWD): None	9/21/2018
Biologics	J7188	Injection, factor VIII (antihemophilic factor, recombinant), (Obizur), per IU	1 IU	1/1/2016	Obizur*	antihemophilic factor (recombinant), porcine sequence lyophilized powder for solution for intravenous injection	Treatment of bleeding episodes in adults with acquired hemophilia A.	630,000	18 years	N/A	N/A	Y	Y		4/10/2019
Biologics	J7189	Factor viia (antihemophilic factor, recombinant), (novoseven rt), 1 microgram	1 mcg	1/1/2006	NovoSeven*, NovoSeven* RT	coagulation factor VIIa (recombinant) for intravenous use	<p>Indicated for:</p> <ul style="list-style-type: none"> • Treatment of bleeding episodes and peri-operative management in adults and children with hemophilia A or B with inhibitors, congenital Factor VII (FVII) deficiency, and Glanzmann's thrombasthenia with refractoriness to platelet transfusions, with or without antibodies to platelets. • Treatment of bleeding episodes and peri-operative management in adults with acquired hemophilia. 	96,000	N/A	N/A	N/A	Y	Y		12/28/2020
Biologics	J7190	Factor VIII (antihemophilic factor [human]) per IU	1 IU	1/1/2000	Hemofil® M, Koate®-DVI, Monoclate-P®	factor VIII (antihemophilic factor, human) for intravenous injection	<p>Koate: Indicated for the control and prevention of bleeding episodes or in order to perform emergency and elective surgery in patients with hemophilia A (hereditary Factor VIII deficiency).</p> <p>Limitation of Use: Koate is not indicated for the treatment of von Willebrand disease.</p> <p>Kogenate: indicated for:</p>	24,000	N/A	N/A	N/A	Y	Y		10/10/2018
Biologics	J7192	Factor VIII (antihemophilic factor, recombinant) per IU, not otherwise specified	1 IU	1/1/2000	Advate®, Bioclata®, Helixate® FS, Kogenate® FS, Recombinate™, ReFacto®	factor VIII (antihemophilic factor, recombinant) for intravenous use	<p>Advate: Indicated for use in children and adults with hemophilia A for:</p> <ul style="list-style-type: none"> • On-demand treatment and control of bleeding episodes in adults and children with hemophilia A. • Perioperative management of bleeding in adults and children with hemophilia A. • Routine prophylaxis to reduce the frequency of bleeding episodes in children with hemophilia A and to reduce the risk of joint damage in children without pre-existing joint damage. • Routine prophylaxis to reduce the frequency of bleeding episodes in adults with hemophilia A. <p>Kogenate is not indicated for the treatment of von Willebrand disease.</p>	54,000	N/A	N/A	N/A	Y	Y		10/10/2018

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

•Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.

•11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacturer participating in the Medicaid Drug Rebate Program (MDRP).

•The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.

•The HCPCS Code effective date represents the date the HCPCS code was established

•Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacture as they are not classified as covered outpatient drugs.

•Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/nccli-medicare/medicaid-nccli-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Biologicals	J7193	Factor IX (antihemophilic factor, purified, non-recombinant) per IU	1 IU	1/1/2002	AlphaNine® SD, Mononine®	coagulation factor IX (human)	Indicated for the prevention and control of bleeding episodes in patients with Factor IX deficiency (hemophilia B, Christmas disease).	42,000	N/A	N/A	N/A	Y	Y		10/10/2018
Biologicals	J7194	Factor IX, complex, per IU	per IU	1/1/2000	Bebulin® VH, Profilnine® SD, Profilnine®	factor IX complex for intravenous administration	Bebulin: Indicated for the prevention and control of bleeding episodes in adult patients with hemophilia B (congenital Factor IX deficiency or Christmas disease). Bebulin is not indicated for use in the treatment of Factor VII deficiency. No clinical studies have been conducted to show benefit from this product for treating deficiencies other than Factor IX deficiency. Indicated for: • Control and prevention of bleeding episodes in adult and pediatric patients with hemophilia B. • Peri-operative management in adult and pediatric patients with hemophilia B.	59,500	18 years	N/A	N/A	Y	Y		10/26/2018
Biologicals	J7195	Injection factor IX (antihemophilic factor, recombinant) per IU, not otherwise specified	1 IU	1/1/2002	BeneFIX®	coagulation factor IX (recombinant) for intravenous use	Indicated for: • Control and prevention of bleeding episodes in adult and pediatric patients with hemophilia B. • Peri-operative management in adult and pediatric patients with hemophilia B.	42,000	N/A	N/A	N/A	Y	Y		10/10/2018
Biologicals	J7196	Injection, antithrombin recombinant, 50 IU	50 IU	1/1/2011	ATryn®	antithrombin (recombinant) lyophilized powder for reconstitution	Indicated for the prevention of peri-operative and peri-partum thromboembolic events in hereditary antithrombin deficient patients.	1,100	18 years	N/A	N/A	Y	Y		9/25/2018
Biologicals	J7197	Antithrombin III (human), per IU	1 IU	1/1/2000	Thrombate III®	antithrombin III (human) lyophilized powder for solution for intravenous injection	Indicated in patients with hereditary antithrombin deficiency for: • Treatment and prevention of thromboembolism • Prevention of peri-operative and peri-partum thromboembolism	40,000	18 years	N/A	N/A	Y	Y		9/25/2018
Biologicals	J7198	Anti-inhibitor, per IU	per IU	1/1/2000	Feiba	anti-inhibitor coagulant complex, for intravenous use, lyophilized powder for solution	Indicated for use in hemophilia A and B patients with inhibitors for: • Control and prevention of bleeding episodes • Perioperative management • Routine prophylaxis to prevent or reduce the frequency of bleeding episodes. Feiba is not indicated for the treatment of bleeding episodes resulting from coagulation factor deficiencies in the absence of inhibitors to factor VIII or factor IX.	560,000	N/A	N/A	N/A	Y	Y		9/21/2018
Biologicals	J7200	Injection, factor IX, (antihemophilic factor, recombinant), Rixubis, per IU	1 IU	1/1/2015	Rixubis®	coagulation factor IX (recombinant) for intravenous injection	Indicated in adults and children with hemophilia B for control and prevention of bleeding episodes, perioperative management, and routine prophylaxis. Rixubis is not indicated for induction of immune tolerance in patients with Hemophilia B.	60,300	N/A	N/A	N/A	Y	Y		10/10/2018
Biologicals	J7201	Injection, factor IX, Fc fusion protein, (recombinant), Alprolix, 1 IU	1 IU	1/1/2017	Alprolix®	coagulation factor IX (recombinant), Fc fusion protein, lyophilized powder for solution for intravenous injection	Indicated for adults and children with hemophilia B for: • On-demand treatment and control of bleeding episodes. • Perioperative management of bleeding. • Routine prophylaxis to reduce the frequency of bleeding episodes. Limitations of Use: Alprolix is not indicated for induction of immune tolerance in patients with hemophilia	72,000	N/A	N/A	N/A	Y	Y		4/10/2019
Biologicals	J7202	Injection, factor IX, albumin fusion protein, (recombinant), Idelvion, 1 IU	1 IU	1/1/2017	Idelvion®	coagulation factor IX (recombinant), albumin fusion protein lyophilized powder for solution for intravenous use	Indicated in children and adults with hemophilia B (congenital Factor IX deficiency) for: • On-demand treatment and control and prevention of bleeding episodes • Perioperative management of bleeding • Routine prophylaxis to reduce the frequency of bleeding episodes Limitations of Use: Idelvion is not indicated for immune tolerance induction in patients with Hemophilia B.	96,921	N/A	N/A	N/A	Y	Y		6/6/2019
Biologicals	J7203	Injection factor ix, (antihemophilic factor, recombinant), glycopegylated, (rebiny), 1 iu	1 IU	1/1/2019	Rebiny®	coagulation factor IX (recombinant), glycoPEGylated, lyophilized powder for solution for intravenous injection	Indicated for use in adults and children with hemophilia B for: • On-demand treatment and control of bleeding episodes • Perioperative management of bleeding Limitations of Use: Rebiny is not indicated for routine prophylaxis in the treatment of patients with hemophilia B or for immune tolerance induction in patients with hemophilia B.	67,200	N/A	N/A	N/A	Y	Y		7/2/2018
Biologicals	J7204	Injection, factor viii, antihemophilic factor (recombinant), (esperoct), glycopegylated-exei, per iu	1 IU	7/1/2020	Esperoct®	antihemophilic factor (recombinant), glycopegylated-exei lyophilized powder for solution, for intravenous use	Indicated for use in adults and children with hemophilia A for: • On-demand treatment and control of bleeding episodes • Perioperative management of bleeding • Routine prophylaxis to reduce the frequency of bleeding episodes Limitation of Use: Esperoct is not indicated for the treatment of von Willebrand disease.	133,000	N/A	N/A	N/A	Y	Y		6/17/2020
Biologicals	J7205	Injection, factor VIII Fc fusion protein (recombinant), per IU	1 IU	1/1/2016	Eloctate®	antihemophilic factor (recombinant) Fc fusion protein lyophilized powder for solution for intravenous injection	Indicated in adults and children with Hemophilia A (congenital Factor VIII deficiency) for: • On-demand treatment and control of bleeding episodes. • Perioperative management of bleeding. • Routine prophylaxis to reduce the frequency of bleeding episodes. Limitation of Use: Eloctate is not indicated for the treatment of von Willebrand disease.	140,000	N/A	N/A	N/A	Y	Y		7/2/2018

North Carolina Division of Health Benefits Physician Administered Drug Program Catalog

•Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.

•11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacture participating in the Medicaid Drug Rebate Program (MDRP).

•The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.

•The HCPCS Code effective date represents the date the HCPCS code was established

•Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacture as they are not classified as covered outpatient drugs.

•Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/nccli-medicaid/medicaid-nccli-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Biologicals	J7207	Injection, factor VIII, (antihemophilic factor, recombinant), pegylated, 1 IU	1 IU	1/1/2017	Adynovate®	antihemophilic factor (recombinant), PEGylated lyophilized powder for solution for intravenous injection	Indicated in children and adult patients with hemophilia A (congenital factor VIII deficiency) for: • On-demand treatment and control of bleeding episodes • Perioperative management • Routine prophylaxis to reduce the frequency of bleeding episodes Adynovate is not indicated for the treatment of von Willebrand disease.	210,000	N/A	N/A	N/A	Y	Y		9/25/2018
Biologicals	J7208	Injection, factor VIII, (antihemophilic factor, recombinant), pegylated-aucI, (jivi), 1 i.u.	1 IU	7/1/2019	Jivi®	antihemophilic factor (recombinant) PEGylated-aucI, for intravenous use	Indicated for use in previously treated adults and adolescents (12 years of age and older) with hemophilia A (congenital Factor VIII deficiency) for: • On-demand treatment and control of bleeding episodes • Perioperative management of bleeding • Routine prophylaxis to reduce the frequency of bleeding episodes Limitations of use: Indicated in adults and children with Hemophilia A for: • On-demand treatment and control of bleeding episodes • Perioperative management of bleeding • Routine prophylaxis to reduce the frequency of bleeding episodes	180,000	12 years	N/A	N/A	Y	Y		9/25/2018
Biologicals	J7209	Injection, factor VIII, (antihemophilic factor, recombinant), (Nuwiq), 1 IU	1 IU	1/1/2017	Nuwiq®	antihemophilic factor (recombinant), lyophilized powder for solution for intravenous injection	Indicated in adults and children with Hemophilia A for: • On-demand treatment and control of bleeding episodes • Perioperative management of bleeding • Routine prophylaxis to reduce the frequency of bleeding episodes Nuwiq is not indicated for the treatment of von Willebrand Disease.	210,000	N/A	N/A	N/A	Y	Y		4/10/2019
Biologicals	J7210	Injection, factor VIII, (antihemophilic factor, recombinant), (Afstyla), 1 IU	1 IU	1/1/2018	Afstyla®	antihemophilic factor (recombinant), single chain for intravenous injection, lyophilized powder for solution	Indicated in adults and children with hemophilia A (congenital Factor VIII deficiency) for: • On-demand treatment and control of bleeding episodes. • Routine prophylaxis to reduce the frequency of bleeding episodes. • Perioperative management of bleeding.	210,000	N/A	N/A	N/A	Y	Y		4/10/2019
Biologicals	J7211	Injection, factor VIII, (antihemophilic factor, recombinant), (Kovaltry), 1 IU	1 IU	1/1/2018	Kovaltry®	factor VIII (antihemophilic factor, recombinant) for intravenous injection	Indicated for use in adults and children with hemophilia A (congenital Factor VIII deficiency) for: • On-demand treatment and control of bleeding episodes • Perioperative management of bleeding • Routine prophylaxis to reduce the frequency of bleeding episodes Kovaltry is not indicated for the treatment of von Willebrand disease.	210,000	N/A	N/A	N/A	Y	Y		10/10/2018
Biologicals	J7212	Factor viia (antihemophilic factor, recombinant)-jncw (sevenfact), 1 microgram	1 mcg	1/1/2021	Sevenfact®	[coagulation factor VIIa (recombinant)-jncw] lyophilized powder for solution, for intravenous use	Indicated for the treatment and control of bleeding episodes occurring in adults and adolescents (12 years of age and older) with hemophilia A or B with inhibitors. Limitation of Use: Sevenfact is not indicated for treatment of congenital factor VII deficiency.	1,260,000	12 years	N/A	N/A	Y	Y		12/28/2020
Biologicals	J7213	Injection, coagulation factor ix (recombinant), ixinity, 1 i.u.	1 IU	7/1/2023	Ixinity®	coagulation factor IX (recombinant) lyophilized powder for solution for intravenous injection	Indicated in adults and children (< 12 years of age) with hemophilia B for: • On-demand treatment and control of bleeding episodes • Perioperative management	322,000	N/A	N/A	N/A	Y	Y		5/3/2024
Biologicals	J7214	Injection, factor VIII/von Willebrand factor complex, recombinant (altuviiio), per factor VIII i.u.	1 IU	10/1/2023	Altuviiio™	antihemophilic factor (recombinant), Fc-VWF-XTEN fusion protein-eHI, lyophilized powder for solution	Indicated for use in adults and children with hemophilia A (congenital factor VIII deficiency) for: • Routine prophylaxis to reduce the frequency of bleeding episodes • On-demand treatment & control of bleeding episodes • Perioperative management of bleeding	112,000	N/A	N/A	N/A	Y	Y		9/28/2023
Drugs	J7296	Levonorgestrel-releasing intrauterine contraceptive system, (Kyleena), 19.5 mg	19.5 mg	1/1/2018	Kyleena®	levonorgestrel-releasing intrauterine system	Indicated for prevention of pregnancy for up to 5 years.	1	After menarche	N/A	Females Only	Y	Y		10/26/2018
Drugs	J7297	Levonorgestrel-releasing intrauterine contraceptive	52 mg	1/1/2017	Liletta®	levonorgestrel-releasing intrauterine system	Indicated for the prevention of pregnancy for up to 8 years.	1	After menarche	N/A	Females Only	Y	Y		7/26/2023
Drugs	J7298	Levonorgestrel-releasing intrauterine contraceptive system (Mirena), 52 mg	52 mg	1/1/2017	Mirena®	levonorgestrel-releasing intrauterine system	Indicated for: • Pregnancy prevention for up to 8 years. • Treatment of heavy menstrual bleeding in women who choose to use intrauterine contraception as their	1	After menarche	N/A	Females Only	Y	Y		9/15/2022
Miscellaneous	J7300	Intrauterine copper contraceptive	1 intrauterine device	1/1/2000	Paragard®	intrauterine copper contraceptive	Indicated for intrauterine contraception for up to 10 years.	1	16 years	N/A	Females Only	Y	Y		7/16/2018
Drugs	J7301	Levonorgestrel-releasing intrauterine contraceptive system (Skyla), 13.5 mg	13.5 mg	1/1/2017	Skyla®	levonorgestrel-releasing intrauterine system	Indicated for the prevention of pregnancy for up to 3 years.	1	After menarche	N/A	Females Only	Y	Y		10/26/2018
Drugs	J7307	Etonogestrel (contraceptive) implant system, including implant and supplies	1 implant	1/1/2008	Nexplanon®	etonogestrel implant for subdermal use	Indicated for use by women to prevent pregnancy.	1	After menarche	N/A	Females Only	Y	Y		10/10/2018
Drugs	J7308	Aminolevulinic acid HCl for topical administration, 20%, single unit dosage form (354)	354 mg	1/1/2004	Levulan® Kerastick®	aminolevulinic acid HCl for topical solution, 20%	Indicated for photodynamic therapy (treatment) of minimally to moderately thick actinic keratoses of the face or scalp, or actinic keratoses of the upper extremities. FDA approval of upper extremity treatment approved 3/6/2018.	1	18 years	N/A	N/A	Y	Y		9/25/2018
Drugs	J7311	Injection, fluocinolone acetonide, intravitreal implant (retisert), 0.01 mg	0.01 mg	1/1/2007	Retisert®	fluocinolone acetonide intravitreal implant	Indicated for the treatment of chronic noninfectious uveitis affecting the posterior segment of the eye.	118	12 years	N/A	N/A	Y	Y		10/10/2018
Drugs	J7312	Injection, dexamethasone, intravitreal implant, 0.1 mg	0.1 mg	1/1/2011	Ozurdex®	dexamethasone intravitreal implant	Indicated for the treatment of macular edema following branch retinal vein occlusion (BRVO) or central retinal vein occlusion (CRVO), non-infectious uveitis affecting the posterior segment of the eye and diabetic macular edema.	14	18 years	N/A	N/A	Y	Y		6/6/2019
Drugs	J7313	Injection, fluocinolone acetonide, intravitreal implant (Iluvien), 0.01 mg	0.01 mg	1/1/2016	Iluvien®	fluocinolone acetonide intravitreal implant	Indicated for the treatment of diabetic macular edema in patients who have been previously treated with a course of corticosteroids and did not have a clinically significant rise in intraocular pressure.	38	18 years	N/A	N/A	Y	Y		10/16/2019

North Carolina Division of Health Benefits Physician Administered Drug Program Catalog

- Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.
- 11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacturer participating in the Medicaid Drug Rebate Program (MDRP).
- The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.
- The HCPCS Code effective date represents the date the HCPCS code was established
- Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacture as they are not classified as covered outpatient drugs.
- Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/ncci-medicare/medicaid-ncci-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Drugs	J7314	Injection, fluocinolone acetonide, intravitreal implant (Yutiq), 0.01 mg	0.01 mg	10/1/2019	Yutiq™	fluocinolone acetonide intravitreal implant 0.18 mg, for intravitreal injection	Indicated for the treatment of non-infectious uveitis affecting the posterior segment of the eye.	36	18 years	N/A	N/A	Y	Y		9/27/2019
Drugs	J7336	Capsaicin 8% patch, per square centimeter	per square centimeter	1/1/2015	Qutenza®	capsaicin 8% patch	<ul style="list-style-type: none"> • Indicated for the management of neuropathic pain associated with postherpetic neuralgia (PHN). • Indicated for the treatment of neuropathic pain associated with diabetic peripheral neuropathy (DPN) of the feet. 	1,120	18 years	N/A	N/A	Y	Y		8/25/2020
Drugs	J7342	Installation, ciprofloxacin otic suspension, 6 mg	6 mg	1/1/2017	Otiprio®	ciprofloxacin otic suspension, for intratympanic or otic use	• Indicated for the treatment of pediatric patients (age 6 months and older) with bilateral otitis media with effusion undergoing tympanostomy tube placement.	10	6 months	N/A	N/A	Y	Y		9/27/2018
Drugs	J7351	Injection, bimatoprost, intracameral implant, 1 microgram	1 mcg	10/1/2020	Durysta™	bimatoprost implant, for intracameral administration	Indicated for the reduction of intraocular pressure (IOP) in patients with open angle glaucoma (OAG) or ocular hypertension (OHT).	20	18 years	N/A	N/A	Y	Y		9/21/2020
Drugs	J7352	Afamelanotide implant, 1 mg	1 mg	1/1/2021	Scenesse®	afamelanotide implant, for subcutaneous use	Indicated to increase pain free light exposure in adult patients with a history of phototoxic reactions from erythropoietic protoporphyria (EPP).	16	18 years	N/A	N/A	Y	Y		11/17/2021
Drugs	J7354	Cantharidin for topical administration, 0.7%, single unit dose applicator (3.2 mg)	3.2 mg (1 ampule)	4/1/2024	Ycanth™	cantharidin topical solution	Indicated for the topical treatment of molluscum contagiosum in adult and pediatric patients 2 years of age and older.	4	2 years	N/A	N/A	Y	Y		3/22/2024
Drugs	J7402	Mometasone furoate sinus implant, (Sinuva), 10 micrograms	10 mcg	4/1/2021	Sinuva™	mometasone furoate sinus implant	Indicated for the treatment of chronic rhinosinusitis with nasal polyps in patients ≥ 18 years of age who have had ethmoid sinus surgery.	270	18 years	N/A	N/A	Y	Y		2/23/2023
Immune Globulins	J7504	Lymphocyte immune globulin, anti-thymocyte globulin, equine, parenteral, 250 mg	250 mg	1/1/2000	Atgam®	lymphocyte immune globulin, anti-thymocyte globulin (equine), sterile solution for intravenous use only	Indicated for: <ul style="list-style-type: none"> •Renal transplant rejection. •Aplastic anemia (moderate to severe) in patients unsuitable for bone marrow transplantation. Limitations of Use: The usefulness of Atgam has not been demonstrated in patients with aplastic anemia who are suitable candidates for bone marrow transplantation or in patients with aplastic anemia	235.2	N/A	N/A	N/A	Y	Y		9/12/2018
Drugs	J7613	Albuterol, inhalation solution, FDA-approved final product, non-compounded, administered through DME, unit dose, 1 mg	1 mg	4/1/2008	N/A	albuterol sulfate inhalation solution (0.021%, 0.042% and 0.083%)	0.63 mg/3 mL solution (0.021%) and 1.25 mg/3 mL solution (0.042%) formulations: Indicated for the relief of bronchospasm in patients 2 to 12 years of age with asthma (reversible obstructive airway disease). 2.5 mg/3 mL solution (0.083%) formulation: Indicated for the relief of bronchospasm in patients 2 years of age and older with reversible obstructive airway disease and acute attacks of bronchospasm.	310	2 years	Formulation Specific Age Restrictions (see comments)	N/A	Y	Y	Formulation Specific: 0.63 mg/3 mL solution (0.021%) and 1.25 mg/3 mL solution (0.042%) formulations: 2 to 12 years of age 2.5 mg/3 mL solution (0.083%) formulation: 2 years of age	9/21/2022
Drugs	J7614	Levalbuterol, inhalation solution, FDA-approved final product, non-compounded, administered through DME, unit dose, 0.5 mg	0.5 mg	4/1/2008	Xopenex®	levalbuterol hydrochloride inhalation solution	Indicated for the treatment or prevention of bronchospasm in adults, adolescents, and children 6 years of age and older with reversible obstructive airway disease.	310	6 years	N/A	N/A	Y	Y		9/23/2022
Drugs	J7620	Albuterol, up to 2.5 mg and ipratropium bromide, up to 0.5 mg, FDA-approved final product, non-compounded, administered through DME	2.5 mg/0.5 mg	1/1/2006	N/A	ipratropium bromide/albuterol sulfate inhalation solution	FDA Approved Indication: Indicated for the treatment of bronchospasm associated with COPD in patients requiring more than one bronchodilator. Recommended Uses from the National Heart, Lung, and Blood Institute: Asthma exacerbations for children through 12 years of age and adults.	186	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	Indication Specific Age Restrictions: Treatment of bronchospasm associated with COPD: 18 years of age and older Asthma exacerbations: N/A	9/21/2022

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

•Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.

•11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacture participating in the Medicaid Drug Rebate Program (MDRP).

•The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.

•The HCPCS Code effective date represents the date the HCPCS code was established

•Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacture as they are not classified as covered outpatient drugs.

•Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/ncci-medicaid/medicaid-ncci-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Drugs	J7644	Ipratropium bromide, inhalation solution, FDA-approved final product, non-compounded, administered through DME, unit dose form, per milligram	1 mg	1/1/2000	N/A	ipratropium bromide inhalation solution, 0.02%	FDA Approved Indication: Indicated as a bronchodilator for maintenance treatment of bronchospasm associated with chronic obstructive pulmonary disease, including chronic bronchitis and emphysema. Recommended Uses from the National Heart, Lung, and Blood Institute: Asthma exacerbations for children through 12 years of age and adults.	93	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	Indication Specific Age Restrictions: Maintenance treatment of bronchospasm associated with chronic obstructive pulmonary disease: 18 years of age and older Asthma exacerbations: N/A	9/23/2022
Drugs	J8499	Prescription drug, oral, non-chemotherapeutic, Not	2 grams	1/1/2000	Flagyl®, Likmez™	metronidazole, oral	Approved indications for use in the PADP: • Symptomatic Trichomoniasis: Metronidazole is indicated for the treatment of <i>T. vaginalis</i> infection in Indicated: • As a component of multiagent adjuvant chemotherapy for treatment of women with axillary lymph node involvement following resection of primary breast cancer. • For the treatment of: acute lymphoblastic leukemia, acute myeloblastic leukemia, Hodgkin lymphoma, Non-Hodgkin lymphoma, metastatic breast cancer, metastatic Wilms' tumor, metastatic neuroblastoma, metastatic soft tissue sarcoma, metastatic bone sarcomas, metastatic ovarian carcinoma, metastatic transitional cell bladder carcinoma, metastatic thyroid carcinoma, metastatic gastric carcinoma,	2	N/A	N/A	N/A	Y	Y		12/1/2023
Drugs	J9000	Injection, doxorubicin hydrochloride, 10 mg	10 mg	1/1/2000	Adriamycin®	doxorubicin hydrochloride for injection, for intravenous use	Indicated for the treatment of adults with metastatic renal cell carcinoma and metastatic melanoma.	38	N/A	N/A	N/A	Y	Y		4/10/2019
Drugs	J9015	Injection, aldesleukin, per single-use via	per single use vial	1/1/2000	Proleukin®	aldesleukin for injection, for intravenous infusion	Indicated for induction of remission and consolidation in patients with acute promyelocytic leukemia (APL) who are refractory to, or have relapsed from, retinoid and anthracycline chemotherapy, and whose APL is characterized by the presence of the t(15;17) translocation or PML/RAR-alpha gene expression. • Indicated in combination with tretinoin for treatment of adults with newly-diagnosed low-risk acute promyelocytic leukemia (APL) whose APL is characterized by the presence of the t(15;17) translocation or	112	18 years	N/A	N/A	Y	Y		6/6/2019
Drugs	J9017	Injection, arsenic trioxide, 1 mg	1 mg	1/1/2000	Trisenox®	arsenic trioxide injection, for intravenous use	Indicated as a component of a multi-agent chemotherapeutic regimen for the treatment of patients with acute lymphoblastic leukemia (ALL) who have developed hypersensitivity to E. coli-derived asparaginase.	651	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	Indication specific age restrictions: • In combination with tretinoin: 18 years of age and older	9/25/2018
Drugs	J9019	Injection, asparaginase (Erwinaze), 1,000 IU	1,000 units	1/1/2013	Erwinaze®	asparaginase erwinia chrysanthemi for injection, for intramuscular (IM) or intravenous (IV) use	Indicated as a component of a multi-agent chemotherapeutic regimen for the treatment of acute lymphoblastic leukemia (ALL) and lymphoblastic lymphoma (LBL) in adult and pediatric patients 1 month or older who have developed hypersensitivity to E. coli-derived asparaginase.	420	1 year	N/A	N/A	Y	Y		6/4/2019
Biologicals	J9021	Injection, asparaginase, recombinant, (rylaze), 0.1 mg	0.1 mg	1/1/2022	Rylaze™	asparaginase erwinia chrysanthemi (recombinant)-rywn injection, for intramuscular use	Indicated for the treatment of patients with: • Non-Small Cell Lung Cancer (NSCLC) o Metastatic non-small cell lung cancer who have disease progression during or following platinum-containing chemotherapy. Patients with EGFR or ALK genomic tumor aberrations should have disease progression on FDA approved therapy for these aberrations prior to receiving Tecentriq. o in combination with bevacizumab, paclitaxel, and carboplatin, for the first-line treatment of patients with metastatic non-squamous NSCLC with no EGFR or ALK genomic tumor aberrations. o in combination with paclitaxel protein-bound and carboplatin for the first-line treatment of adult	12,200	1 month	N/A	N/A	Y	Y		12/20/2022
Biologicals	J9022	Injection, atezolizumab, 10 mg	10 mg	1/1/2018	Tecentriq®	atezolizumab injection, for intravenous use	Indicated for the treatment of patients with: • Non-Small Cell Lung Cancer (NSCLC) o Metastatic non-small cell lung cancer who have disease progression during or following platinum-containing chemotherapy. Patients with EGFR or ALK genomic tumor aberrations should have disease progression on FDA approved therapy for these aberrations prior to receiving Tecentriq. o in combination with bevacizumab, paclitaxel, and carboplatin, for the first-line treatment of patients with metastatic non-squamous NSCLC with no EGFR or ALK genomic tumor aberrations. o in combination with paclitaxel protein-bound and carboplatin for the first-line treatment of adult	336	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	NSCLC, SCLC, HCC, melanoma: 18 years of age and older ASPS: 2 years of age and older	1/23/2023

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

•Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.

•11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacture participating in the Medicaid Drug Rebate Program (MDRP).

•The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.

•The HCPCS Code effective date represents the date the HCPCS code was established

•Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacture as they are not classified as covered outpatient drugs.

•Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/nci-medicaid/medicaid-nci-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Biologics	J9023	Injection, avelumab, 10 mg	10 mg	1/1/2018	Bavencio®	avelumab injection, for intravenous use	Indicated for: • Adults and pediatric patients 12 years and older with metastatic Merkel cell carcinoma (MCC). • Patients with locally advanced or metastatic urothelial carcinoma (UC) who have disease progression during or following platinum-containing chemotherapy or have disease progression within 12 months of neoadjuvant or adjuvant treatment with platinum-containing chemotherapy. • Maintenance treatment of patients with locally advanced or metastatic UC that has not progressed with first-line platinum-containing chemotherapy.	240	12 years	N/A	N/A	Y	Y		7/28/2020
Drugs	J9025	Injection, azacitidine, 1 mg	1 mg	1/1/2006	Vidaza®	azacitidine for injection, for subcutaneous or intravenous use	Indicated for the treatment of: - Adult patients with the following FAB myelodysplastic syndrome (MDS) subtypes: refractory anemia (RA) or refractory anemia with ringed sideroblasts (RARS) (if accompanied by neutropenia or thrombocytopenia or requiring transfusions), refractory anemia with excess blasts (RAEB), refractory anemia with excess blasts in transformation (RAEB-T) and chronic myelomonocytic leukemia (CMML).	3,000	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	Indication specific age restrictions: • Adult patients with FAB myelodysplastic syndrome (MDS) subtypes - 18 years of	6/9/2022
Biologics	J9029	Intravesical instillation, nadofaragene firadenovec-	1 therapeutic dose	7/1/2023	Adstiladrin®	nadofaragene firadenovec-vncg suspension, for	Indicated for the treatment of adult patients with high-risk Bacillus Calmette-Guérin (BCG)-unresponsive non-muscle invasive bladder cancer (NMIBC) with carcinoma in situ (CIS) with or without papillary tumors.	1	18 years	N/A	N/A	Y	Y		3/22/2024
Biologics	J9030	Bcg live intravesical instillation, 1 mg	per installation	1/1/2000	Tice BCG®	BCG Live (intravesical)	Indicated for the treatment and prophylaxis of carcinoma in situ (CIS) of the urinary bladder, and for the prophylaxis of primary or recurrent stage Ta and/or T1 papillary tumors following transurethral resection (TUR). Tice BCG is not recommended for stage TaG1 papillary tumors, unless they are judged to be at high risk of tumor recurrence. Tice BCG is not indicated for papillary tumors of stages higher than T1.	5	18 years	N/A	N/A	Y	Y		6/8/2019
Drugs	J9032	Injection, belinostat, 10 mg	10 mg	1/1/2016	Beleodaq®	belinostat for injection, for intravenous use	Indicated for the treatment of patients with relapsed or refractory peripheral T-cell lymphoma (PTCL).	2,500	18 years	N/A	N/A	Y	Y		4/10/2019
Drugs	J9033	Injection, bendamustine HCl (Treanda), 1 mg	1 mg	1/1/2017	Treanda®	bendamustine hydrochloride injection, for intravenous use	Indicated for treatment of patients with: • Chronic lymphocytic leukemia (CLL). Efficacy relative to first line therapies other than chlorambucil has not been established. • Indolent B-cell non-Hodgkin lymphoma (NHL) that has progressed during or within six months of treatment with rituximab or a rituximab-containing regimen.	1,200	18 years	N/A	N/A	Y	Y		9/25/2018
Drugs	J9034	Injection, bendamustine HCl (Bendeka), 1 mg	1 mg	1/1/2017	Bendeka®	bendamustine hydrochloride injection, for intravenous use	Indicated for treatment of patients with: • Chronic lymphocytic leukemia (CLL). Efficacy relative to first line therapies other than chlorambucil has not been established. • Indolent B-cell non-Hodgkin lymphoma (NHL) that has progressed during or within six months of treatment with rituximab or a rituximab-containing regimen.	1,200	18 years	N/A	N/A	Y	Y		9/25/2018
Biologics	J9035	Injection, bevacizumab, 10 mg	10 mg	1/1/2005	Avastin®	bevacizumab injection, for intravenous use	Indicated for the treatment of: • Metastatic colorectal cancer, in combination with intravenous 5-fluorouracil-based chemotherapy for first- or second-line treatment. • Metastatic colorectal cancer, in combination with fluoropyrimidine-irinotecan- or fluoropyrimidine-	420	18 years	N/A	N/A	Y	Y		10/20/2022
Drugs	J9036	Injection, bendamustine hydrochloride, (Belrapzo/bendamustine), 1 mg	1 mg	7/1/2019	Belrapzo™	bendamustine hydrochloride injection for intravenous use	Indicated for treatment of patients with: • Chronic lymphocytic leukemia (CLL). Efficacy relative to first line therapies other than chlorambucil has not been established. • Indolent B-cell non-Hodgkin lymphoma (NHL) that has progressed during or within six months of treatment with rituximab or a rituximab-containing regimen.	1,200	18 years	N/A	N/A	Y	Y		8/26/2019
Biologics	J9039	Injection, blinatumomab, 1 mcg	1 mcg	1/1/2016	Blinicyto®	blinatumomab for injection, for intravenous use	Treatment of adults and children with: • Relapsed or refractory CD19-positive B-cell precursor acute lymphoblastic leukemia (ALL). • CD19-positive B-cell precursor acute lymphoblastic leukemia (ALL) in first or second complete remission	784	N/A	N/A	N/A	Y	Y		4/26/2021

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

•Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.

•11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacture participating in the Medicaid Drug Rebate Program (MDRP).

•The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.

•The HCPCS Code effective date represents the date the HCPCS code was established

•Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacture as they are not classified as covered outpatient drugs.

•Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/nccli-medicaid/medicaid-nccli-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Drugs	J9040	Injection, bleomycin sulfate, 15 units	15 units	1/1/2000	N/A	bleomycin for injection	Considered a palliative treatment shown to be useful in the management of: <ul style="list-style-type: none"> • Squamous Cell Carcinoma: Head and neck (including mouth, tongue, tonsil, nasopharynx, oropharynx, sinus, palate, lip, buccal mucosa, gingivae, epiglottis, skin, larynx), penis, cervix, and vulva. The response to bleomycin is poorer in patients with previously irradiated head and neck cancer. • Lymphomas: Hodgkin's disease, non-Hodgkin's disease • Testicular Carcinoma: Embryonal cell, choriocarcinoma, and teratocarcinoma • Malignant Pleural Effusion: Bleomycin is effective as a sclerosing agent for the treatment of malignant pleural effusion and prevention of recurrent pleural effusions. 	27	N/A	N/A	N/A	Y	Y		4/10/2019
Drugs	J9041	Injection, bortezomib, 0.1 mg	0.1 mg	1/1/2005	Velcade*	bortezomib for injection, for subcutaneous or intravenous use	Indicated for treatment of patients with: <ul style="list-style-type: none"> • Multiple myeloma • Mantle cell lymphoma 	245	18 years	N/A	N/A	Y	Y		12/12/2022
Biologicals	J9042	Injection, brentuximab vedotin, 1 mg	1 mg	1/1/2013	Adcetris*	brentuximab vedotin for injection, for intravenous use	Indicated for: <ul style="list-style-type: none"> • Previously untreated Stage III or IV classical Hodgkin lymphoma (cHL), in combination with doxorubicin, vinblastine, and dacarbazine. • Classical Hodgkin Lymphoma (cHL) at high risk of relapse or progression as post-autologous 	360	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	Indication specific age restrictions: <ul style="list-style-type: none"> • Previously untreated high risk classical Hodgkin 	12/20/2022
Drugs	J9043	Injection, cabazitaxel, 1 mg	1 mg	1/1/2012	Jevtana*	cabazitaxel injection, for intravenous use	Indicated in combination with prednisone for treatment of patients with hormone-refractory metastatic prostate cancer previously treated with a docetaxel-containing treatment regimen.	240	18 years	N/A	Males Only	Y	Y		9/27/2018
Drugs	J9045	Injection, carboplatin, 50 mg	50 mg	1/1/2000	N/A	carboplatin injection for intravenous use	Indicated for the initial treatment of advanced ovarian carcinoma in established combination with other approved chemotherapeutic agents and for the palliative treatment of patients with ovarian carcinoma recurrent after prior chemotherapy, including patients who have previously been treated with cisplatin.	36	18 years	N/A	N/A	Y	Y		4/10/2019
Drugs	J9046	Injection, bortezomib (dr. reddy's), not therapeutically equivalent to J9041, 0.1 mg	0.1 mg	1/1/2023	N/A	bortezomib for injection, for intravenous use (Dr. Reddy's)	Indicated for: <ul style="list-style-type: none"> • treatment of adult patients with multiple myeloma • treatment of adult patients with mantle cell lymphoma who have received at least 1 prior therapy 	245	18 years	N/A	N/A	Y	Y		12/12/2022
Drugs	J9047	Injection, carfilzomib, 1 mg	1 mg	1/1/2014	Kyprolis*	carfilzomib for injection, for intravenous use	Indicated for the treatment of adult patients with relapsed or refractory multiple myeloma who have received one to three lines of therapy in combination with: <ul style="list-style-type: none"> o Lenalidomide and dexamethasone; or o Dexamethasone; or 	1060	18 years	N/A	N/A	Y	Y		7/20/2022
Drugs	J9048	Injection, bortezomib (fresenius kabi), not therapeutically equivalent to J9041, 0.1 mg	0.1 mg	1/1/2023	N/A	bortezomib for injection, for intravenous use (Fresenius Kabi)	Indicated for: <ul style="list-style-type: none"> • treatment of adult patients with multiple myeloma • treatment of adult patients with mantle cell lymphoma who have received at least 1 prior therapy 	245	18 years	N/A	N/A	Y	Y		12/12/2022
Drugs	J9049	Injection, bortezomib (hospira), not therapeutically equivalent to J9041, 0.1 mg	0.1 mg	1/1/2023	N/A	bortezomib for injection, for subcutaneous or intravenous use (Hospira)	Indicated for: <ul style="list-style-type: none"> • treatment of adult patients with multiple myeloma • treatment of adult patients with mantle cell lymphoma 	245	18 years	N/A	N/A	Y	Y		12/19/2022
Drugs	J9050	Injection, carmustine, 100 mg	100 mg	1/1/2000	BiCNU*	carmustine for injection	Indicated as palliative therapy as a single agent or in established combination therapy with other approved chemotherapeutic agents in the following: <ul style="list-style-type: none"> • Brain tumors - glioblastoma, brainstem glioma, medulloblastoma, astrocytoma, ependymoma, and metastatic brain tumors. • Multiple myeloma - in combination with prednisone. • Hodgkin's disease - as secondary therapy in combination with other approved drugs in patients who relapse while being treated with primary therapy, or who fail to respond to primary therapy. • Non-Hodgkin's lymphomas - as secondary therapy in combination with other approved drugs for 	5	18 years	N/A	N/A	Y	Y		5/20/2019

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

•Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.

•11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacture participating in the Medicaid Drug Rebate Program (MDRP).

•The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.

•The HCPCS Code effective date represents the date the HCPCS code was established

•Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacture as they are not classified as covered outpatient drugs.

•Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/ncci-medicaid/medicaid-ncci-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Drugs	J9051	Injection, bortezomib (maia), not therapeutically equivalent to J9041, 0.1 mg	0.1 mg	10/1/2023	N/A	bortezomib injection, for intravenous use (Maia)	Indicated for: • treatment of adult patients with multiple myeloma • treatment of adult patients with mantle cell lymphoma	245	18 years	N/A	N/A	Y	Y		9/28/2023
Drugs	J9052	Injection, carmustine (accord), not therapeutically equivalent to J9050, 100 mg	100 mg	1/1/2024	N/A	carmustine for injection, for intravenous use (Accord)	Carmustine for injection is indicated as palliative therapy as a single agent or in established combination therapy with other approved chemotherapeutic agents in the following: • Brain tumors glioblastoma, brainstem glioma, medulloblastoma, astrocytoma, ependymoma, and metastatic brain tumors • Multiple myeloma-in combination with prednisone	5	18 years	N/A	N/A	Y	Y		12/22/2023
Biologicals	J9055	Injection, cetuximab, 10 mg	10 mg	1/1/2005	Erbix*	cetuximab injection, for intravenous use	Indicated for: • Squamous Cell Carcinoma of the Head and Neck (SCCHN): - Locally or regionally advanced squamous cell carcinoma of the head and neck in combination with radiation therapy	390	18 years	N/A	N/A	Y	Y		10/26/2021
Drugs	J9056	Injection, bendamustine hydrochloride (vivimusta), 1 mg	1 mg	7/1/2023	Vivimusta	bendamustine hydrochloride injection, for intravenous use	Indicated for treatment of patients with: • Chronic lymphocytic leukemia (CLL). Efficacy relative to first line therapies other than chlorambucil has not been established. • Indolent B-cell non-Hodgkin lymphoma (NHL) that has progressed during or within six months of treatment with rituximab or a rituximab-containing regimen.	1,200	18 years	N/A	N/A	Y	Y		6/22/2023
Drugs	J9057	Injection, copanlisib, 1 mg	1 mg	1/1/2019	Alliqpa™	copanlisib injection, for intravenous use	Indicated for the treatment of adult patients with relapsed follicular lymphoma (FL) who have received at least two prior systemic therapies. Accelerated approval was granted for this indication based on overall response rate. Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial.	240	18 years	N/A	N/A	Y	Y		8/5/2021
Drugs	J9058	Injection, bendamustine hydrochloride (apotex), 1 mg	1 mg	7/1/2023	N/A	bendamustine hydrochloride injection, for intravenous use (Apotex)	Indicated for treatment of adult patients with: • Chronic lymphocytic leukemia (CLL). Efficacy relative to first line therapies other than chlorambucil has not been established. • Indolent B-cell non-Hodgkin lymphoma (NHL) that has progressed during or within six months of treatment with rituximab or a rituximab-containing regimen.	1,200	18 years	N/A	N/A	Y	Y		6/22/2023
Drugs	J9059	Injection, bendamustine hydrochloride (baxter), 1 mg	1 mg	7/1/2023	N/A	bendamustine hydrochloride injection, for intravenous use (Baxter)	Indicated for treatment of adult patients with: • Chronic lymphocytic leukemia (CLL). Efficacy relative to first line therapies other than chlorambucil has not been established. • Indolent B-cell non-Hodgkin lymphoma (NHL) that has progressed during or within six months of	1,200	18 years	N/A	N/A	Y	Y		6/22/2023
Drugs	J9060	Injection, cisplatin, powder or solution, per 10 mg	10 mg	1/1/2000	N/A	cisplatin injection	Indicated as therapy for: • Metastatic Testicular Tumors: In established combination therapy with other approved chemotherapeutic agents in patients with metastatic testicular tumors who have already received appropriate surgical and/or radiotherapeutic procedures.	50	18 years	N/A	N/A	Y	Y		9/27/2018
Biologicals	J9061	Injection, amivantamab-vmjw, 2 mg	2 mg	1/1/2022	Rybreant™	amivantamab-vmjw injection, for intravenous use	Indicated: - as a single agent for the treatment of adult patients with locally advanced or metastatic non-small cell lung cancer (NSCLC) with epidermal growth factor receptor (EGFR) exon 20 insertion mutations, as detected by an FDA-approved test, whose disease has progressed on or after platinum-based chemotherapy. - in combination with carboplatin and pemetrexed for the first-line treatment of adult patients with locally advanced or metastatic non-small cell lung cancer (NSCLC) with epidermal growth factor receptor (EGFR) exon 20 insertion mutations, as detected by an FDA-approved test.	2,800	18 years	N/A	N/A	Y	Y		5/3/2024

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

- Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.
- 11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacture participating in the Medicaid Drug Rebate Program (MDRP).
- The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.
- The HCPCS Code effective date represents the date the HCPCS code was established
- Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacture as they are not classified as covered outpatient drugs.
- Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/ncci-medicare/medicaid-ncci-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Biologicals	J9063	Injection, mirvetuximab soravtansine-gynx, 1 mg	1 mg	7/1/2023	Elahere™	mirvetuximab soravtansine-gynx injection, for intravenous use	Indicated for the treatment of adult patients with FRα positive, platinum-resistant epithelial ovarian, fallopian tube, or primary peritoneal cancer, who have received one to three prior systemic treatment regimens. Select patients for therapy based on an FDA-approved test.	1,800	18 years	N/A	N/A	Y	Y		6/22/2023
Drugs	J9065	Injection, cladribine, per 1 mg	1 mg	1/1/2000	N/A	cladribine injection	Indicated for the treatment of active Hairy Cell Leukemia as defined by clinically significant anemia, neutropenia, thrombocytopenia, or disease-related symptoms.	91	18 years	N/A	N/A	Y	Y		6/4/2019
Drugs	J9071	Injection, cyclophosphamide (auromedics), 5 mg	5 mg	4/1/2022	N/A	cyclophosphamide for injection, for intravenous use (AuroMedics)	Indicated for the treatment of: Malignant Diseases: malignant lymphomas, Hodgkin's disease, lymphocytic lymphoma, mixed-cell type lymphoma, histiocytic lymphoma, Burkitt's lymphoma; multiple myeloma, leukemias, mycosis fungoides, neuroblastoma, adenocarcinoma of ovary, retinoblastoma, breast carcinoma.	2,500	N/A	N/A	N/A	Y	Y		3/17/2022
Drugs	J9072	Injection, cyclophosphamide (dr. reddy's), 5 mg	5 mg	1/1/2024	N/A	cyclophosphamide injection, for intravenous use (Dr. Reddy's)	Cyclophosphamide injection is indicated for treatment of adult and pediatric patients with: • Malignant Diseases: - malignant lymphomas: Hodgkin's disease, lymphocytic lymphoma, mixed-cell type lymphoma, histiocytic lymphoma, Burkitt's lymphoma; - multiple myeloma, leukemias, mycosis fungoides, neuroblastoma, adenocarcinoma of ovary.	2,500	N/A	N/A	N/A	Y	Y		12/22/2023
Drugs	J9073	Injection, cyclophosphamide (ingenus), 5 mg	5 mg	4/1/2024	N/A	cyclophosphamide injection, for intravenous use (Ingenus)	Cyclophosphamide is indicated for treatment of: • Malignant Diseases: malignant lymphomas: Hodgkin's disease, lymphocytic lymphoma, mixed-cell type lymphoma, histiocytic lymphoma, Burkitt's lymphoma; multiple myeloma, leukemias, mycosis fungoides, neuroblastoma, adenocarcinoma of ovary, retinoblastoma, breast carcinoma.	2,100	N/A	N/A	N/A	Y	Y		3/27/2024
Drugs	J9074	Injection, cyclophosphamide (sandoz), 5 mg	5 mg	4/1/2024	N/A	cyclophosphamide injection, for intravenous use (Sandoz)	Cyclophosphamide injection is an alkylating drug indicated for treatment of adult patients with: Malignant Diseases: malignant lymphomas: Hodgkin's lymphoma, lymphocytic lymphoma, mixed-cell type lymphoma, histiocytic lymphoma, Burkitt's lymphoma; multiple myeloma, leukemias, mycosis fungoides.	2,100	18 years	N/A	N/A	Y	Y		5/3/2024
Drugs	J9075	Injection, cyclophosphamide, not otherwise specified, 5 mg	5 mg	4/1/2024	N/A	cyclophosphamide for injection, for intravenous use	Indicated for the treatment of: Malignant Diseases: malignant lymphomas: Hodgkin's disease, lymphocytic lymphoma, mixed-cell type lymphoma, histiocytic lymphoma, Burkitt's lymphoma; multiple myeloma, leukemias, mycosis fungoides, neuroblastoma, adenocarcinoma of ovary, retinoblastoma, breast carcinoma.	2,500	N/A	N/A	N/A	Y	Y		3/22/2024
Drugs	J9100	Injection, cytarabine, 100 mg	100 mg	1/1/2000	N/A	cytarabine injection	In combination with other approved anticancer drugs, is indicated for remission induction in acute non-lymphocytic leukemia of adults and pediatric patients. It has also been found useful in the treatment of acute lymphocytic leukemia and the blast phase of chronic myelocytic leukemia. Intrathecal administration of cytarabine injection (preservative-free preparations only) is indicated in the prophylaxis and treatment of meningeal leukemia.	35	N/A	N/A	N/A	Y	Y		7/2/2018
Biologicals	J9118	Injection, calaspargase pegol-mknl, 10 units	10 units	10/1/2019	Asparlas™	calaspargase pegol-mknl injection, for intravenous use	Indicated for the treatment of acute lymphoblastic leukemia in pediatric and young adult patients age 1 month to 21 years.	1,500	1 month	21 years	N/A	Y	Y		12/3/2019

North Carolina Division of Health Benefits Physician Administered Drug Program Catalog

- Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.
- 11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacturer participating in the Medicaid Drug Rebate Program (MDRP).
- The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.
- The HCPCS Code effective date represents the date the HCPCS code was established
- Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacture as they are not classified as covered outpatient drugs.
- Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/ncci-medicaid/medicaid-ncci-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Biologics	J9119	Injection, cemiplimab-rwlc, 1 mg	1 mg	10/1/2019	Libtayo®	cemiplimab-rwlc injection, for intravenous use	<p>Indicated</p> <ul style="list-style-type: none"> • for the treatment of patients with metastatic cutaneous squamous cell carcinoma (CSCC) or locally advanced CSCC who are not candidates for curative surgery or curative radiation. • for the treatment of patients with locally advanced BCC (laBCC) previously treated with a hedgehog pathway inhibitor or for whom a hedgehog pathway inhibitor is not appropriate. • for the treatment of patients with metastatic BCC (mBCC) previously treated with a hedgehog pathway inhibitor or for whom a hedgehog pathway inhibitor is not appropriate. • for the first-line treatment of patients with non-small cell lung cancer (NSCLC) whose tumors have high PD-L1 expression [Tumor Proportion Score (TPS) ≥ 50%] as determined by an FDA-approved test, with no EGFR, ALK or ROS1 aberrations, and is: <ul style="list-style-type: none"> - locally advanced where patients are not candidates for surgical resection or definitive chemoradiation OR - metastatic. • in combination with platinum-based chemotherapy for the first-line treatment of adult patients with non-small cell lung cancer (NSCLC) with no EGFR, ALK or ROS1 aberrations and is: <ul style="list-style-type: none"> - locally advanced where patients are not candidates for surgical resection or definitive chemoradiation OR - metastatic. 	700	18 years	N/A	N/A	Y	Y		12/20/2022
Drugs	J9120	Injection, dactinomycin, 0.5 mg	0.5 mg	1/1/2000	Cosmegen®	dactinomycin for injection, for intravenous use	<p>Indicated for the treatment of:</p> <ul style="list-style-type: none"> • adult and pediatric patients with Wilms tumor, as part of a multi-phase, combination chemotherapy regimen • adult and pediatric patients with rhabdomyosarcoma, as part of a multi-phase, combination chemotherapy regimen • adult and pediatric patients with Ewing sarcoma, as part of a multi-phase, combination chemotherapy regimen • adult and pediatric patients with metastatic, nonseminomatous testicular cancer, as part of a multi-phase, combination chemotherapy regimen • post-menarchal patients with gestational trophoblastic neoplasia, as a single agent or as part of a combination chemotherapy regimen • adult patients with locally recurrent or locoregional solid malignancies, as a component of palliative or adjunctive regional perfusion 	42	N/A	N/A	N/A	Y	Y		9/25/2018
Drugs	J9130	Dacarbazine, 100 mg	100 mg	1/1/2000	N/A	dacarbazine for injection	Indicated for the treatment of metastatic malignant melanoma and as secondary-line therapy when used in combination with other effective agents for Hodgkin's disease.	91	N/A	N/A	N/A	Y	Y		6/10/2019
Biologics	J9144	Injection, daratumumab, 10 mg and hyaluronidase-fihj	10 mg	1/1/2021	Darzalex Faspro™	daratumumab and hyaluronidase-fihj injection, for subcutaneous use	<p>Indicated for the treatment of adult patients with:</p> <ul style="list-style-type: none"> • multiple myeloma in combination with bortezomib, melphalan and prednisone in newly diagnosed patients who are ineligible for autologous stem cell transplant • multiple myeloma in combination with lenalidomide and dexamethasone in newly diagnosed patients who are ineligible for autologous stem cell transplant and in patients with relapsed or refractory multiple myeloma 	900	18 years	N/A	N/A	Y	Y		12/16/2021
Biologics	J9145	Injection, daratumumab, 10 mg	10 mg	1/1/2017	Darzalex®	daratumumab injection, for intravenous use	<p>Indicated for the treatment of adult patients with multiple myeloma:</p> <ul style="list-style-type: none"> • in combination with lenalidomide and dexamethasone in patients with relapsed or refractory multiple myeloma who have received at least one prior therapy. 	1,120	18 years	N/A	N/A	Y	Y		9/21/2020
Drugs	J9150	Injection, daunorubicin, 10 mg	10 mg	1/1/2000	N/A	daunorubicin hydrochloride injection	In combination with other approved anticancer drugs, daunorubicin is indicated for remission induction in acute nonlymphocytic leukemia (myelogenous, monocytic, erythroid) of adults and for remission induction in acute lymphocytic leukemia of children and adults.	60	N/A	N/A	N/A	Y	Y		6/10/2019

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

•Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.

•11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacturer participating in the Medicaid Drug Rebate Program (MDRP).

•The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.

•The HCPCS Code effective date represents the date the HCPCS code was established

•Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacture as they are not classified as covered outpatient drugs.

•Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/nccli-medicaid/medicaid-nczi-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Drugs	J9153	Injection, liposomal, 1 mg daunorubicin and 2.27 mg cytarabine	1 mg/2.27 mg	1/1/2019	Vyxeos™	daunorubicin and cytarabine liposome injection, for intravenous use	Indicated for: • the treatment of adults with newly-diagnosed therapy-related acute myeloid leukemia (t-AML) or AML with myelodysplasia-related changes (AML-MRC). • the treatment of newly-diagnosed therapy-related acute myeloid leukemia (t-AML) or AML with myelodysplasia-related changes (AML-MRC) in pediatric patients 1 year and older.	660	1 year	N/A	N/A	Y	Y		4/26/2021
Drugs	J9155	Injection, degarelix, 1 mg	1 mg	1/1/2010	Firmagon®	degarelix for injection for subcutaneous administration	Indicated for the treatment of patients with advanced prostate cancer.	320	18 years	N/A	Males Only	Y	Y		10/4/2018
Drugs	J9171	Injection, docetaxel, 1 mg	1 mg	1/1/2010	Docetrez®, Taxotere®	docetaxel injection concentrate, intravenous infusion	Indicated for: • Breast Cancer (BC): single agent for locally advanced or metastatic BC after chemotherapy failure; and with doxorubicin and cyclophosphamide as adjuvant treatment of operable node-positive BC. • Non-Small Cell Lung Cancer (NSCLC): single agent for locally advanced or metastatic NSCLC after platinum therapy failure; and with cisplatin for unresectable, locally advanced or metastatic untreated NSCLC. • Hormone Refractory Prostate Cancer (HRPC): with prednisone in androgen independent (hormone refractory) metastatic prostate cancer. • Gastric Adenocarcinoma (GC): with cisplatin and fluorouracil for untreated, advanced GC, including the	500	N/A	N/A	N/A	Y	Y		6/8/2019
Biologicals	J9173	Injection, durvalumab, 10 mg	10 mg	1/1/2019	Imfinzi®	durvalumab injection, for intravenous use	Imfinzi is a programmed death-ligand 1 (PD-L1) blocking antibody indicated for the treatment of patients with: Indicated in: • combination with lenalidomide and dexamethasone for the treatment of adult patients with multiple myeloma who have received one to three prior therapies. • combination with pomalidomide and dexamethasone for the treatment of adult patients with multiple myeloma who have received at least two prior therapies including lenalidomide and a proteasome inhibitor.	420	18 years	N/A	N/A	Y	Y		12/20/2022
Biologicals	J9176	Injection, elotuzumab, 1 mg	1 mg	1/1/2017	Empliciti®	elotuzumab for injection, for intravenous use	Indicated: - as a single agent for the treatment of adult patients with locally advanced or metastatic urothelial cancer who:	5,600	18 years	N/A	N/A	Y	Y		5/20/2019
Biologicals	J9177	Injection, enfortumab vedotin-ejfv, 0.25 mg	0.25 mg	7/1/2020	Padcev®	enfortumab vedotin-ejfv for injection, for intravenous use	Indicated: - as a single agent for the treatment of adult patients with locally advanced or metastatic urothelial cancer who:	2,080	18 years	N/A	N/A	Y	Y		2/16/2024
Drugs	J9178	Injection, epirubicin HCl, 2 mg	2 mg	1/1/2004	Ellence®	epirubicin hydrochloride injection	Indicated as a component of adjuvant therapy in patients with evidence of axillary node tumor involvement following resection of primary breast cancer.	300	18 years	N/A	N/A	Y	Y		10/10/2018
Drugs	J9179	Injection, eribulin mesylate, 0.1 mg	0.1 mg	1/1/2012	Halaven®	eribulin mesylate injection, for intravenous use	Indicated for the treatment of patients with: • Metastatic breast cancer who have previously received at least two chemotherapeutic regimens for the treatment of metastatic disease. Prior therapy should have included an anthracycline and a taxane in either the adjuvant or metastatic setting. • Unresectable or metastatic liposarcoma who have received a prior anthracycline-containing regimen.	160	18 years	N/A	N/A	Y	Y		6/4/2019
Drugs	J9181	Injection, etoposide, 10 mg	10 mg	1/1/2000	Etopophos®, Toposar™	etoposide phosphate for injection, for intravenous use	Indicated for the treatment of patients with: • Refractory testicular tumors, in combination with other chemotherapeutic drugs. • Small cell lung cancer, in combination with cisplatin, as first-line treatment.	300	18 years	N/A	N/A	Y	Y		6/10/2019
Drugs	J9185	Injection, fludarabine phosphate, 50 mg	50 mg	1/1/2000	N/A	fludarabine phosphate for injection for intravenous use	Indicated for the treatment of adult patients with B-cell chronic lymphocytic leukemia (CLL) who have not responded to or whose disease has progressed during treatment with at least 1 standard alkylating-agent containing regimen. The safety and effectiveness of fludarabine in previously untreated or non-refractory patient with CLL have not been established.	16	18 years	N/A	N/A	Y	Y		10/10/2018
Drugs	J9190	Injection, fluorouracil, 500 mg	500 mg	1/1/2000	Adrucil®	fluorouracil injection for intravenous use	Indicated for the treatment of patients with: • Adenocarcinoma of the colon and rectum • Adenocarcinoma of the breast • Gastric adenocarcinoma • Pancreatic adenocarcinoma	45	18 years	N/A	N/A	Y	Y		4/10/2019
Drugs	J9196	Injection, gemcitabine hydrochloride (accord), not therapeutically equivalent to J9201, 200 mg	200 mg	4/1/2023	N/A	gemcitabine injection, for intravenous use (Accord)	Indicated: • in combination with carboplatin, for the treatment of advanced ovarian cancer that has relapsed at least 6 months after completion of platinum-based therapy. • in combination with paclitaxel, for first-line treatment of metastatic breast cancer after failure of prior anthracycline-containing adjuvant chemotherapy, unless anthracyclines were clinically contraindicated. • in combination with cisplatin for the treatment of non-small cell lung cancer. • as a single agent for the treatment of pancreatic cancer.	64	18 years	N/A	N/A	Y	Y		3/16/2023

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

•Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.

•11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacturer participating in the Medicaid Drug Rebate Program (MDRP).

•The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.

•The HCPCS Code effective date represents the date the HCPCS code was established

•Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacturer as they are not classified as covered outpatient drugs.

•Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/nccli-medicaid/medicaid-nccli-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Drugs	J9198	Injection, gemcitabine hydrochloride, (infugem), 100 mg	100 mg	7/1/2020	Infugem™	gemcitabine in sodium chloride injection, for intravenous use	Indicated: • in combination with carboplatin, for the treatment of advanced ovarian cancer that has relapsed at least 6 months after completion of platinum-based therapy. • in combination with paclitaxel, for first-line treatment of metastatic breast cancer after failure of prior anthracycline-containing adjuvant chemotherapy, unless anthracyclines were clinically contraindicated. • in combination with cisplatin for the treatment of non-small cell lung cancer. • as a single agent for the treatment of pancreatic cancer.	128	18 years	N/A	N/A	Y	Y		6/17/2020
Drugs	J9200	Injection, floxuridine, 500 mg	500 mg	1/1/2000	N/A	floxuridine for injection, for intra-arterial infusion	Effective in the palliative management of gastrointestinal adenocarcinoma metastatic to the liver, when given by continuous regional intra-arterial infusion in carefully selected patients who are considered incurable by surgery or other means. Patients with known disease extending beyond an area capable of	5	18 years	N/A	N/A	Y	Y		10/26/2018
Drugs	J9201	Injection, gemcitabine hydrochloride, not otherwise specified, 200 mg	200 mg	1/1/2000	Gemzar®	gemcitabine for injection, for intravenous use	Indicated: • In combination with carboplatin, for the treatment of advanced ovarian cancer that has relapsed at least 6 months after completion of platinum-based therapy. • In combination with paclitaxel, for first-line treatment of metastatic breast cancer after failure of prior anthracycline-containing adjuvant chemotherapy, unless anthracyclines were clinically contraindicated. • In combination with cisplatin for the treatment of non-small cell lung cancer. • As a single agent for the treatment of pancreatic cancer.	64	18 years	N/A	N/A	Y	Y		1/9/2020
Biologicals	J9203	Injection, gemtuzumab ozogamicin, 0.1 mg	0.1 mg	1/1/2018	Mylotarg™	gemtuzumab ozogamicin injection, for intravenous use	Indicated for: • the treatment of newly-diagnosed CD33-positive acute myeloid leukemia (AML) in adults. • the treatment of newly-diagnosed CD33-positive acute myeloid leukemia (AML) in pediatric patients 1 month and older. • the treatment of relapsed or refractory CD33-positive AML in adults and in pediatric patients 2 years	275	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	Indication specific age restrictions: • Newly-diagnosed CD33-positive acute myeloid leukemia: 1 month of age and	7/28/2020
Biologicals	J9204	Injection, mogamulizumab-kpkc, 1 mg	1 mg	10/1/2019	Poteligeo®	mogamulizumab-kpkc injection, for intravenous use	Indicated for the treatment of adult patients with relapsed or refractory mycosis fungoides or Sézary syndrome after at least one prior systemic therapy.	700	18 years	N/A	N/A	Y	Y		9/27/2019
Drugs	J9205	Injection, irinotecan liposome, 1 mg	1 mg	1/1/2017	Onivyde™	irinotecan liposome injection, for intravenous use	Indicated: - in combination with fluorouracil and leucovorin, for the treatment of patients with metastatic adenocarcinoma of the pancreas after disease progression following gemcitabine-based therapy.	645	18 years	N/A	N/A	Y	Y		3/22/2024
Drugs	J9206	Injection, irinotecan, 20 mg	20 mg	1/1/2000	Camptosar®	irinotecan injection, intravenous infusion	Indicated for: • First-line therapy in combination with 5-fluorouracil and leucovorin for patients with metastatic	88	18 years	N/A	N/A	Y	Y		4/10/2019
Drugs	J9207	Injection, ixabepilone, 1 mg	1 mg	1/1/2009	Ixempra®	ixabepilone for injection, for intravenous use	Indicated for the treatment • In combination with capecitabine for patients with metastatic or locally advanced breast cancer resistant to treatment with an anthracycline and a taxane, or whose cancer is taxane resistant and for	180	18 years	N/A	N/A	Y	Y		2/23/2023
Drugs	J9208	Injection, ifosfamide, 1 gram	1 g	1/1/2000	Ifex®	ifosfamide for injection, intravenous use	Indicated for use in combination with certain other approved antineoplastic agents for third-line chemotherapy of germ cell testicular cancer. It should be used in combination with mesna for prophylaxis of hemorrhagic cystitis.	30	18 years	N/A	N/A	Y	Y		6/4/2019
Drugs	J9209	Injection, mesna, 200 mg	200 mg	1/1/2000	Mesnex®	mesna injection solution	Indicated as a prophylactic agent in reducing the incidence of ifosfamide-induced hemorrhagic cystitis.	90	18 years	N/A	N/A	Y	Y		8/5/2021
Biologicals	J9210	Injection, emapalumab-lzsg, 1 mg	1 mg	10/1/2019	Gamifant™	emapalumab-lzsg injection, for intravenous use	Indicated for the treatment of adult and pediatric (newborn and older) patients with primary hemophagocytic lymphohistiocytosis (HLH) with refractory, recurrent or progressive disease or intolerance with conventional HLH therapy.	14,000	N/A	N/A	N/A	Y	Y		5/27/2020
Drugs	J9211	Injection, idarubicin hydrochloride, 5 mg	5 mg	1/1/2000	Idamycin®	idarubicin hydrochloride for injection	Indicated in combination with other approved antileukemic drugs for the treatment of acute myeloid leukemia in adults. This includes French-American-British (FAB) classifications M1 through M7.	36	18 years	N/A	N/A	Y	Y		10/31/2018
Biologicals	J9214	Injection, interferon, alfa-2b, recombinant, 1 million units	1 million units	1/1/2000	Intron® A	interferon alfa-2b recombinant for injection	Indicated for: hairy cell leukemia, malignant melanoma, follicular lymphoma, condylomata acuminata, AIDS-related Kaposi's sarcoma, chronic hepatitis C and chronic hepatitis B. Please see package insert for additional information on each indication.	1,050	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	Indication specific: 18 years and older for all indications except chronic Hepatitis B and C. Hepatitis B - 1 year of age and older	6/4/2019
Biologicals	J9215	Injection, interferon, alfa-n3, (human leukocyte derived), 250,000 IU	250,000 IU	1/1/2000	Alferon® N	interferon alfa-n3 injection	Indicated for condyloma acuminata.	100	18 years	N/A	N/A	Y	Y		10/4/2018
Biologicals	J9216	Injection, interferon, gamma-1b, 3 million units	3 million units	1/1/2000	Actimmune®	interferon gamma-1b injection, for subcutaneous use	Indicated for: • Reducing the frequency and severity of serious infections associated with Chronic Granulomatous Disease (CGD)	18.67	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	Indication specific age restrictions: CGD: 1 year and older	5/6/2019
Drugs	J9217	Leuprolide acetate (for depot suspension), 7.5 mg	7.5 mg	1/1/2000	Eligard®, Lupron Depot®	leuprolide acetate for injectable suspension, for doses 7.5 mg and greater	Eligard: Indicated for the treatment of advanced prostate cancer. Lupron Depot: Indicated for the treatment of advanced prostatic cancer.	6	18 years	N/A	Males Only	Y	Y		2/19/2024

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

- Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.
- 11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacture participating in the Medicaid Drug Rebate Program (MDRP).
- The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.
- The HCPCS Code effective date represents the date the HCPCS code was established
- Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacture as they are not classified as covered outpatient drugs.
- Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/ncci-medicaid/medicaid-ncci-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Drugs	J9218	Leuprolide acetate, per 1 mg	per 1 mg	1/1/2000	N/A	leuprolide acetate injection	Indicated in the palliative treatment of advanced prostatic cancer.	31	N/A	N/A	Males Only	Y	Y		2/19/2024
Drugs	J9223	Injection, lurbinectedin, 0.1 mg	0.1 mg	1/1/2021	Zepzelca™	lurbinectedin for injection, for intravenous use	Indicated for the treatment of adult patients with metastatic small cell lung cancer (SCLC) with disease progression on or after platinum-based chemotherapy.	160	18 years	N/A	N/A	Y	Y		12/28/2020
Drugs	J9225	Histrelin implant (Vantas), 50 mg	50 mg	1/1/2006	Vantas®	histrelin acetate subcutaneous implant	Indicated for the palliative treatment of advanced prostate cancer.	1	18 years	N/A	Males Only	Y	Y		2/19/2024
Drugs	J9226	Histrelin implant (Supprelin LA), 50 mg	50 mg	1/1/2008	Supprelin® LA	histrelin acetate subcutaneous implant	Indicated for the treatment of children with central precocious puberty (CPP).	1	2 years	N/A	N/A	Y	Y		2/19/2024
Biologicals	J9227	Injection, isatuximab-irfc, 10 mg	10 mg	10/1/2020	Sarclisa®	isatuximab-irfc injection, for intravenous use	Indicated • in combination with pomalidomide and dexamethasone, for the treatment of adult patients with multiple myeloma who have received at least two prior therapies including lenalidomide and a Indicated for:	700	18 years	N/A	N/A	Y	Y		4/26/2021
Biologicals	J9228	Injection, ipilimumab, 1 mg	1 mg	1/1/2012	Yervoy®	ipilimumab injection, for intravenous use	• Adjuvant treatment of patients with cutaneous melanoma with pathologic involvement of regional lymph nodes of more than 1 mm who have undergone complete resection, including total	2,800	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	Indication specific age restrictions: • Melanoma as a single agent	3/21/2023
Biologicals	J9229	Injection, inotuzumab ozogamicin, 0.1 mg	0.1 mg	1/1/2019	Besponsa™	inotuzumab ozogamicin injection, for intravenous use	Indicated for the treatment of relapsed or refractory CD22-positive B-cell precursor acute lymphoblastic leukemia (ALL) in adult and pediatric patients 1 year and older.	108	1 year	N/A	N/A	Y	Y		5/3/2024

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

•Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.

•11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacture participating in the Medicaid Drug Rebate Program (MDRP).

•The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.

•The HCPCS Code effective date represents the date the HCPCS code was established

•Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacture as they are not classified as covered outpatient drugs.

•Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/ncci-medicaid/medicaid-ncci-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Drugs	J9245	Injection, melphalan hydrochloride, not otherwise specified, 50 mg	50 mg	1/1/2000	Alkeran*	melphalan hydrochloride for injection	Indicated for the palliative treatment of patients with multiple myeloma for whom oral therapy is not appropriate.	3	18 years	N/A	N/A	Y	Y		6/17/2020
Drugs	J9246	Injection, melphalan (evomela), 1 mg	1 mg	7/1/2020	Evomela*	melphalan for injection, for intravenous use	Indicated for: • use as a high-dose conditioning treatment prior to hematopoietic progenitor (stem) cell transplantation in patients with multiple myeloma.	500	18 years	N/A	N/A	Y	Y		9/28/2021
Drugs	J9259	Injection, paclitaxel protein-bound particles (american regent), not therapeutically equivalent to J9264, 1 mg	1 mg	7/1/2023	N/A	paclitaxel protein-bound particles for injectable suspension, (albumin-bound), for intravenous use	Indicated for the treatment of: • Metastatic breast cancer, after failure of combination chemotherapy for metastatic disease or relapse within 6 months of adjuvant chemotherapy. Prior therapy should have included an anthracycline unless clinically contraindicated.	1,600	18 years	N/A	N/A	Y	Y		6/22/2023
Drugs	J9260	Methotrexate sodium, 50 mg	50 mg	1/1/2000	N/A	methotrexate sodium injection, 50 mg	• Methotrexate is indicated in the treatment of gestational choriocarcinoma, chorioadenoma destruens and hydatidiform mole. • In acute lymphocytic leukemia, methotrexate is indicated in the prophylaxis of meningeal leukemia and is used in maintenance therapy in combination with other chemotherapeutic agents. Methotrexate is also	3,000	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	Indication specific age restrictions: • Cancer chemotherapy: None • Polyarticular-course juvenile	1/26/2024
Drugs	J9261	Injection, nelarabine, 50 mg	50 mg	1/1/2007	Arranon*	nelarabine injection, for intravenous use	Indicated for the treatment of patients with T-cell acute lymphoblastic leukemia and T-cell lymphoblastic lymphoma in adult and pediatric patients age 1 year and older whose disease has not responded to or has relapsed following treatment with at least two chemotherapy regimens.	450	1 year	N/A	N/A	Y	Y		12/16/2021
Drugs	J9262	Injection, omacetaxine mepesuccinate, 0.01 mg	0.01 mg	1/1/2014	Synribo*	omacetaxine mepesuccinate for injection, for subcutaneous use	Indicated for the treatment of adult patients with chronic or accelerated phase chronic myeloid leukemia (CML) with resistance and/or intolerance to two or more tyrosine kinase inhibitors.	10,625	18 years	N/A	N/A	Y	Y		9/21/2018
Drugs	J9263	Injection, oxaliplatin, 0.5 mg	0.5 mg	1/1/2004	Eloxatin*	oxaliplatin injection for intravenous use	Indicated for: • Adjuvant treatment of stage III colon cancer in patients who have undergone complete resection of the	1,500	18 years	N/A	N/A	Y	Y		6/4/2019
Drugs	J9264	Injection, paclitaxel protein-bound particles, 1 mg	1 mg	1/1/2006	Abraxane*	paclitaxel protein-bound particles for injectable suspension, (albumin-bound)	Indicated for the treatment: • Metastatic breast cancer, after failure of combination chemotherapy for metastatic disease or relapse within six months of adjuvant chemotherapy. Prior therapy should have included an anthracycline unless	1,600	18 years	N/A	N/A	Y	Y		5/25/2023
Biologicals	J9266	Injection, pegaspargase, per single dose vial (3,750 IU)	per single dose vial (3,750 IU)	1/1/2000	Oncaspar*	pegaspargase injection, for intramuscular or intravenous use	Indicated as a component of a multi-agent chemotherapeutic regimen for treatment of patients with: • First line acute lymphoblastic leukemia • Acute lymphoblastic leukemia and hypersensitivity to asparaginase	6	1 year	N/A	N/A	Y	Y		8/24/2018

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

•Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.

•11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacture participating in the Medicaid Drug Rebate Program (MDRP).

•The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.

•The HCPCS Code effective date represents the date the HCPCS code was established

•Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacture as they are not classified as covered outpatient drugs.

•Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/nccli-medicaid/medicaid-nccli-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Drugs	J9267	Injection, paclitaxel, 1 mg	1 mg	1/1/2015	Taxol®	paclitaxel injection	Indicated for breast cancer, ovarian cancer, non-small cell lung cancer, and AIDS-related kaposi sarcoma. See package insert for full details of each indication.	875	18 years	N/A	N/A	Y	Y		9/27/2018
Drugs	J9268	Injection, pentostatin, per 10 mg	10 mg	7/15/2001	Nipent®	pentostatin for injection	Indicated as single-agent treatment for both untreated and alpha-interferon-refractory hairy cell leukemia patients with active disease as defined by clinically significant anemia, neutropenia, thrombocytopenia, or disease-related symptoms.	3	18 years	N/A	N/A	Y	Y		9/21/2018
Biologics	J9269	Injection, tagraxofusp-erzs, 10 micrograms	10 mcg	10/1/2019	Elzonris™	tagraxofusp-erzs injection, for intravenous use	Indicated for the treatment of blastic plasmacytoid dendritic cell neoplasm (BPDCN) in adults and in pediatric patients 2 years and older.	2,000	2 years	N/A	N/A	Y	Y		10/3/2019
Biologics	J9271	Injection, pembrolizumab, 1 mg	1 mg	1/1/2016	Keytruda®	pembrolizumab injection, for intravenous use	Melanoma: 1. Indicated for the treatment of patients with unresectable or metastatic melanoma. 2. Indicated for the adjuvant treatment of adult and pediatric (12 years and older) patients with Stage IIB, Endometrial Cancer (EC)	400	The safety and effectiveness of Keytruda as a single	N/A	N/A	Y	Y		2/27/2024
Biologics	J9272	Injection, dostarlimab-gxly, 10 mg	10 mg	1/1/2022	Jemperli	dostarlimab-gxly injection, for intravenous use	• Indicated for the treatment of adult patients with mismatch repair deficient (dMMR) recurrent or advanced endometrial cancer, as determined by an FDA-approved test, that has progressed on or following prior treatment with a platinum-containing regimen in any setting and are not candidates for curative surgery or radiation.	150	18 years	N/A	Endometrial Cancer: Females only Solid Tumors: None	Y	Y		9/13/2023
Biologics	J9273	Injection, tisotumab vedotin-tftv, 1 mg	1 mg	4/1/2022	Tivdak™	tisotumab vedotin-tftv for injection, for intravenous use	Indicated for the treatment of adult patients with recurrent or metastatic cervical cancer with disease progression on or after chemotherapy.	400	18 years	N/A	N/A	Y	Y		3/21/2022
Biologics	J9274	Injection, tebentafusp-tebn, 1 microgram	1 mcg	10/1/2022	Kimtrak®	tebentafusp-tebn injection, for intravenous use	Indicated for the treatment of HLA-A*02:01-positive adult patients with unresectable or metastatic uveal melanoma.	500	18 years	N/A	N/A	Y	Y		9/15/2022
Drugs	J9280	Injection, mitomycin, 5 mg	5 mg	1/1/2000	Mutamycin®	mitomycin for injection, 5 mg	Mitomycin is not recommended as single-agent, primary therapy. It has been shown to be useful in the therapy of disseminated adenocarcinoma of the stomach or pancreas in proven combinations with other approved chemotherapeutic agents and as palliative treatment when other modalities have failed. Mitomycin is not recommended to replace appropriate surgery and/or radiotherapy.	10	18 years	N/A	N/A	Y	Y		6/7/2019

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

•Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.

•11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacture participating in the Medicaid Drug Rebate Program (MDRP).

•The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.

•The HCPCS Code effective date represents the date the HCPCS code was established

•Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacture as they are not classified as covered outpatient drugs.

•Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/ncci-medicaid/medicaid-ncci-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Drugs	J9281	Mitomycin pyelocalyceal instillation, 1 mg	1 mg	1/1/2021	Jelmyto™	mitomycin for pyelocalyceal solution	Indicated for the treatment of adult patients with low-grade Upper Tract Urothelial Cancer (LG-UTUC).	400	18 years	N/A	N/A	Y	Y		12/28/2020
Biologicals	J9286	Injection, glofitamab-gxbm, 2.5 mg	2.5 mg	1/1/2024	Columvi™	glofitamab-gxbm injection, for intravenous use	Indicated for the treatment of adult patients with relapsed or refractory diffuse large B-cell lymphoma, not otherwise specified (DLBCL, NOS) or large B-cell lymphoma (LBC) arising from follicular lymphoma, after two or more lines of systemic therapy.	24	18 years	N/A	N/A	Y	Y		12/22/2023
Drugs	J9293	Injection, mitoxantrone hydrochloride, per 5 mg	5 mg	1/1/2000	N/A	mitoxantrone hydrochloride injection, solution	Indicated: • For reducing neurologic disability and/or the frequency of clinical relapses in patients with secondary (chronic) progressive, progressive relapsing, or worsening relapsing-remitting multiple sclerosis (i.e., patients whose neurologic status is significantly abnormal between relapses). Mitoxantrone is not indicated in the treatment of patients with primary progressive multiple sclerosis. • In combination with corticosteroids is indicated as initial chemotherapy for the treatment of patients with pain related to advanced hormone-refractory prostate cancer. • In combination with other approved drug(s) is indicated in the initial therapy of acute nonlymphocytic leukemia (ANLL) in adults. This category includes myelogenous, promyelocytic, monocytic, and erythroid acute leukemias.	30	18 years	N/A	N/A	Y	Y	Lifetime Maximum Dose: 70 units	10/31/2018
Drugs	J9294	Injection, pemetrexed (hospira), not therapeutically equivalent to J9305, 10 mg	10 mg	4/1/2023	N/A	pemetrexed for injection, for intravenous use (Hospira)	Indicated: • In combination with cisplatin for the initial treatment of patients with locally advanced or metastatic, non-squamous, non-small cell lung cancer (NSCLC). • As a single agent for the maintenance treatment of patients with locally advanced or metastatic, non-squamous NSCLC whose disease has not progressed after four cycles of platinum-based first-line chemotherapy. • As a single agent for the treatment of patients with recurrent, metastatic non-squamous, NSCLC after prior chemotherapy. Limitations of Use: Pemetrexed for Injection is not indicated for the treatment of patients with squamous cell, non-small cell lung cancer. • Initial treatment, in combination with cisplatin, of patients with malignant pleural mesothelioma whose disease is unresectable or who are otherwise not candidates for curative surgery.	300	18 years	N/A	N/A	Y	Y		3/16/2023
Biologicals	J9295	Injection, necitumumab, 1 mg	1 mg	1/1/2017	Portrazza™	necitumumab injection, for intravenous use	Indicated, in combination with gemcitabine and cisplatin, for first-line treatment of patients with metastatic squamous non-small cell lung cancer. Limitation of Use: Portrazza is not indicated for treatment of non-squamous non-small cell lung cancer.	3,200	18 years	N/A	N/A	Y	Y		7/2/2018
Drugs	J9296	Injection, pemetrexed (accord), not therapeutically equivalent to J9305, 10 mg	10 mg	4/1/2023	N/A	pemetrexed injection, for intravenous use (Accord)	Indicated: • in combination with pembrolizumab and platinum chemotherapy, for the initial treatment of patients with metastatic non-squamous non-small cell lung cancer (NSCLC), with no EGFR or ALK genomic tumor aberrations. • in combination with cisplatin for the initial treatment of patients with locally advanced or metastatic, non-squamous, NSCLC. • as a single agent for the maintenance treatment of patients with locally advanced or metastatic, non-squamous NSCLC whose disease has not progressed after four cycles of platinum-based first-line chemotherapy. • as a single agent for the treatment of patients with recurrent, metastatic non-squamous, NSCLC after prior chemotherapy. Limitations of Use: Pemetrexed Injection is not indicated for the treatment of patients with squamous cell, non-small cell lung cancer. • Initial treatment, in combination with cisplatin, of patients with malignant pleural mesothelioma whose disease is unresectable or who are otherwise not candidates for curative surgery.	300	18 years	N/A	N/A	Y	Y		3/16/2023

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

•Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.

•11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacture participating in the Medicaid Drug Rebate Program (MDRP).

•The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.

•The HCPCS Code effective date represents the date the HCPCS code was established

•Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacture as they are not classified as covered outpatient drugs.

•Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/ncci-medicaid/medicaid-ncci-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Drugs	J9297	Injection, pemtrexed (sandoz), not therapeutically equivalent to J9305, 10 mg	10 mg	4/1/2023	N/A	pemtrexed injection, for intravenous use (Sandoz)	Indicated: • in combination with pembrolizumab and platinum chemotherapy, for the initial treatment of patients with metastatic non-squamous non-small cell lung cancer (NSCLC), with no EGFR or ALK genomic tumor aberrations. • in combination with cisplatin for the initial treatment of patients with locally advanced or metastatic, non-squamous, NSCLC. • as a single agent for the maintenance treatment of patients with locally advanced or metastatic, non-squamous NSCLC whose disease has not progressed after four cycles of platinum-based first-line chemotherapy.	300	18 years	N/A	N/A	Y	Y		3/16/2023
Biologicals	J9298	Injection, nivolumab and relatimab-rmbw, 3 mg/1 mg	3 mg/1 mg	10/1/2022	Opduaig™	nivolumab and relatimab-rmbw injection, for intravenous use	Indicated for the treatment of adult and pediatric patients 12 years of age or older with unresectable or metastatic melanoma.	320	12 years	N/A	N/A	Y	Y		9/15/2022
Biologicals	J9299	Injection, nivolumab, 1 mg	1 mg	1/1/2016	Opdivo®	nivolumab injection, for intravenous use	Indicated for: Melanoma: • adult and pediatric (12 years and older) patients with unresectable or metastatic melanoma, as a single agent or in combination with ipilimumab. • for the adjuvant treatment of adult and pediatric patients 12 years and older with completely resected Stage IIB, Stage IIC, Stage III, or Stage IV melanoma. NSCLC: • the treatment of patients with metastatic non-small cell lung cancer (NSCLC) and progression on or after	1,260	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	Indication specific age restrictions: • MSI-H or dMMR mCRC - 12 years of age and older • Melanoma, as a single agent, in combination with ipilimumab, or in the adjuvant setting - 12 years and older	5/3/2024
Biologicals	J9301	Injection, obinutuzumab, 10 mg	10 mg	1/1/2015	Gazyva®	obinutuzumab Injection, for intravenous use	Indicated: • In combination with chlorambucil, for the treatment of patients with previously untreated chronic lymphocytic leukemia. • In combination with bendamustine followed by Gazyva monotherapy, for the treatment of patients with follicular lymphoma who relapsed after, or are refractory to, a rituximab-containing regimen. • In combination with chemotherapy followed by Gazyva monotherapy in patients achieving at least a partial remission, for the treatment of adult patients with previously untreated stage II bulky, III or IV follicular lymphoma.	400	18 years	N/A	N/A	Y	Y		7/16/2018
Biologicals	J9302	Injection, ofatumumab, 10 mg	10 mg	1/1/2011	Arzerra®	ofatumumab injection, for intravenous use	Indicated for the treatment of chronic lymphocytic leukemia (CLL): • in combination with chlorambucil, for the treatment of previously untreated patients with CLL for whom fludarabine-based therapy is considered inappropriate. • in combination with fludarabine and cyclophosphamide for the treatment of patients with relapsed CLL • for extended treatment of patients who are in complete or partial response after at least two lines of therapy for recurrent or progressive CLL. • for the treatment of patients with CLL refractory to fludarabine and alemtuzumab.	1,000	18 years	N/A	N/A	Y	Y	Pregnancy: May cause fetal B-cell depletion.	7/16/2018
Biologicals	J9303	Injection, panitumumab, 10 mg	10 mg	1/1/2008	Vectibix®	panitumumab injection, for intravenous use	Indicated for the treatment of wild-type RAS (defined as wild-type in both KRAS and NRAS as determined by an FDA-approved test for this use) metastatic colorectal cancer (mCRC): - In combination with Folfox for first-line treatment. - As monotherapy following disease progression after prior treatment with fluoropyrimidine, oxaliplatin, and irinotecan-containing chemotherapy. Limitation of Use: Vectibix is not indicated for the treatment of patients with RAS-mutant mCRC or for whom RAS mutation status is unknown.	270	18 years	N/A	N/A	Y	Y		6/4/2019
Drugs	J9304	Injection, pemtrexed (pemfexy), 10 mg	10 mg	10/1/2020	Pemfexy™	pemtrexed injection, for intravenous use	Indicated: • in combination with cisplatin for the initial treatment of patients with locally advanced or metastatic	300	18 years	N/A	N/A	Y	Y		1/23/2023

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

•Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.

•11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacture participating in the Medicaid Drug Rebate Program (MDRP).

•The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.

•The HCPCS Code effective date represents the date the HCPCS code was established

•Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacture as they are not classified as covered outpatient drugs.

•Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/ncci-medicaid/medicaid-ncci-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Drugs	J9305	Injection, pemetrexed, not otherwise specified, 10 mg	10 mg	10/1/2020	Alimta*	pemetrexed for injection, for intravenous use	Indicated: <ul style="list-style-type: none"> • In combination with cisplatin for the initial treatment of patients with locally advanced or metastatic, non-squamous, non-small cell lung cancer (NSCLC). • As a single agent for the maintenance treatment of patients with locally advanced or metastatic, non-squamous NSCLC whose disease has not progressed after four cycles of platinum-based first-line chemotherapy. • As a single agent for the treatment of patients with recurrent metastatic non-squamous, NSCLC after prior chemotherapy. • Initial treatment, in combination with cisplatin, of patients with malignant pleural mesothelioma whose disease is unresectable or who are otherwise not candidates for curative surgery. • In combination with carboplatin and pembrolizumab for the initial treatment of patients with metastatic, non-squamous NSCLC. Limitations of Use: Not indicated for the treatment of patients with squamous cell, non-small cell lung cancer.	300	18 years	N/A	N/A	Y	Y		12/12/2022
Biologics	J9306	Injection, pertuzumab, 1 mg	1 mg	1/1/2014	Perjeta*	pertuzumab injection, for intravenous use	Indicated for: <ul style="list-style-type: none"> • Use in combination with trastuzumab and docetaxel for treatment of patients with HER2-positive metastatic breast cancer (MBC) who have not received prior anti-HER2 therapy or chemotherapy for metastatic disease. • Use in combination with trastuzumab and chemotherapy as o Neoadjuvant treatment of patients with HER2-positive, locally advanced, inflammatory, or early stage breast cancer (either greater than 2 cm in diameter or node positive) as part of a complete treatment regimen for early breast cancer. 	1,260	18 years	N/A	N/A	Y	Y		7/2/2018
Drugs	J9307	Injection, pralatrexate, 1 mg	1 mg	1/1/2011	Folotyn*	pralatrexate injection, for intravenous use	Indicated for the treatment of patients with relapsed or refractory peripheral T-cell lymphoma.	400	18 years	N/A	N/A	Y	Y		8/24/2018
Biologics	J9308	Injection, ramucirumab, 5 mg	5 mg	1/1/2016	Cyramza*	ramucirumab injection, for intravenous use	Indicated: <ul style="list-style-type: none"> • As a single agent or in combination with paclitaxel, for treatment of advanced gastric or gastro-esophageal junction adenocarcinoma, with disease progression on or after prior fluoropyrimidine- or platinum-containing chemotherapy. • In combination with docetaxel, for treatment of metastatic non-small cell lung cancer with disease progression on or after platinum-based chemotherapy. Patients with EGFR or ALK genomic tumor aberrations should have disease progression on FDA-approved therapy for these aberrations prior to receiving Cyramza. • In combination with erlotinib, for first-line treatment of metastatic non-small cell lung cancer with epidermal growth factor receptor (EGFR) exon 19 deletions or exon 21 (L858R) mutations. • In combination with Folfiri, for the treatment of metastatic colorectal cancer with disease progression on or after prior therapy with bevacizumab, oxaliplatin, and a fluoropyrimidine. • As a single agent, for the treatment of hepatocellular carcinoma in patients who have an alpha fetoprotein of >400 ng/mL and have been treated with sorafenib. 	900	18 years	N/A	N/A	Y	Y		6/17/2020

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

•Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.

•11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacture participating in the Medicaid Drug Rebate Program (MDRP).

•The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.

•The HCPCS Code effective date represents the date the HCPCS code was established

•Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacture as they are not classified as covered outpatient drugs.

•Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/nccli-medicaid/medicaid-nccli-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Biologics	J9309	Injection, polatuzumab vedotin-piiq, 1 mg	1 mg	1/1/2020	Polivy*	polatuzumab vedotin-piiq for injection, for intravenous use	Indicated: • in combination with bendamustine and a rituximab product for the treatment of adult patients with relapsed or refractory diffuse large B-cell lymphoma, not otherwise specified, after at least two prior therapies. • in combination with a rituximab product, cyclophosphamide, doxorubicin, and prednisone (R-CHP) for the treatment of adult patients who have previously untreated diffuse large B-cell lymphoma (DLBCL), not otherwise specified (NOS) or high-grade B-cell lymphoma (HGBL) and who have an International Prognostic Index score of 2 or greater.	560	18 years	N/A	N/A	Y	Y		5/25/2023
Biologics	J9311	Injection, rituximab 10 mg and hyaluronidase	10 mg	1/1/2019	Rituxan Hycela*	rituximab and hyaluronidase human injection, for subcutaneous use	Indicated for the treatment of adult patients with: • Follicular Lymphoma (FL): o Relapsed or refractory, follicular lymphoma as a single agent o Previously untreated follicular lymphoma in combination with first line chemotherapy and, in patients achieving a complete or partial response to rituximab in combination with chemotherapy, as single-agent maintenance therapy o Non-progressing (including stable disease), follicular lymphoma as a single agent after first-line cyclophosphamide, vincristine, and prednisone (CVP) chemotherapy • Diffuse Large B-cell Lymphoma (DLBCL): o Previously untreated diffuse large B-cell lymphoma in combination with cyclophosphamide, doxorubicin,	700	18 years	N/A	N/A	Y	Y		4/19/2019
Biologics	J9312	Injection, rituximab, 10 mg	10 mg	1/1/2019	Rituxan*	rituximab injection, for intravenous use	Indicated for the treatment of adult patients with: • Non-Hodgkin's Lymphoma (NHL) - Relapsed or refractory, low grade or follicular, CD20-positive B-cell NHL as a single agent. - Previously untreated follicular, CD20-positive, B-cell NHL in combination with first line chemotherapy and, in patients achieving a complete or partial response to Rituxan in combination with chemotherapy, as single-agent maintenance therapy. - Non-progressing (including stable disease), low-grade, CD20-positive, B-cell NHL as a single agent after first-line cyclophosphamide, vincristine, and prednisone (CVP) chemotherapy. - Previously untreated diffuse large B-cell, CD20-positive NHL in combination with (cyclophosphamide, doxorubicin, vincristine, and prednisone) (CHOP) or other anthracycline-based chemotherapy regimens. Indicated for the treatment of pediatric patients aged 6 months and older with mature B-cell NHL and	500	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	Indication Specific: • CLL, RA, PV: 18 years of age and older • GPA and MPA: 2 years of age and older • NHL and B-AL: 6 months of age and older	1/13/2022
Drugs	J9314	Injection, pemetrexed (teva), not therapeutically equivalent to J9305, 10 mg	10 mg	1/1/2023	N/A	pemetrexed for injection, for intravenous use (Teva)	Indicated: • in combination with pembrolizumab and platinum chemotherapy, for the initial treatment of patients with metastatic non-squamous non-small cell lung cancer (NSCLC), with no epidermal growth factor	300	18 years	N/A	N/A	Y	Y		12/12/2022
Biologics	J9316	Injection, pertuzumab, trastuzumab, and hyaluronidase-zzxf, per 10 mg	10 mg	1/1/2021	Phesgo™	pertuzumab, trastuzumab, and hyaluronidase-zzxf injection, for subcutaneous	Indicated for: • Use in combination with chemotherapy as: o neoadjuvant treatment of patients with HER2-positive, locally advanced, inflammatory, or early stage	300	18 years	N/A	N/A	Y	Y		12/28/2020
Biologics	J9317	Injection, sacituzumab govitecan-hziy, 2.5 mg	2.5 mg	1/1/2021	Trodelvy™	sacituzumab govitecan-hziy for injection, for intravenous	Indicated for the treatment of adult patients with: • Unresectable locally advanced or metastatic triple-negative breast cancer (mTNBC) who have received	2,304	18 years	N/A	N/A	Y	Y		3/16/2023
Drugs	J9318	Injection, romidepsin, non-lyophilized, 0.1 mg	0.1 mg	10/1/2021	N/A	romidepsin for injection, for intravenous use (non-lyophilized)	Indicated for: • The treatment of cutaneous T-cell lymphoma (CTCL) in adult patients who have received at least one prior systemic therapy.	2,200	18 years	N/A	N/A	Y	Y		1/13/2022
Drugs	J9319	Injection, romidepsin, lyophilized, 0.1 mg	0.1 mg	10/1/2021	Istodax*	romidepsin for injection, for intravenous use (lyophilized)	Indicated for: • Treatment of cutaneous T-cell lymphoma (CTCL) in patients who have received at least one prior systemic therapy.	1600	18 years	N/A	N/A	Y	Y		9/29/2021
Drugs	J9320	Injection, streptozocin, 1 gram	1 g	1/1/2000	Zanosar*	streptozocin powder, for solution	Indicated in the treatment of metastatic islet cell cancer of pancreas.	20	N/A	N/A	N/A	Y	Y		6/7/2019
Biologics	J9321	Injection, epcoritamab-bysp, 0.16 mg	0.16 mg	1/1/2024	Epkinly™	epcoritamab-bysp injection, for subcutaneous use	Indicated for the treatment of adult patients with relapsed or refractory diffuse large B-cell lymphoma (DLBCL), not otherwise specified, including DLBCL arising from indolent lymphoma, and high-grade B-cell lymphoma after two or more lines of systemic therapy.	1,500	18 years	N/A	N/A	Y	Y		12/22/2023
Drugs	J9323	Injection, pemetrexed ditromethamine, 10 mg	10 mg	7/1/2023	N/A	pemetrexed ditromethamine for injection, for intravenous	Indicated: • in combination with cisplatin for the initial treatment of patients with locally advanced or metastatic,	300	18 years	N/A	N/A	Y	Y		6/22/2023

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

•Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.

•11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacturer participating in the Medicaid Drug Rebate Program (MDRP).

•The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.

•The HCPCS Code effective date represents the date the HCPCS code was established

•Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacturer as they are not classified as covered outpatient drugs.

•Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/nccli-medicaid/medicaid-nccli-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Drugs	J9324	Injection, pemtrexed (pemrydi rti), 10 mg	10 mg	1/1/2024	Pemrydi RTU*	pemtrexed injection, for intravenous use (Shipka)	Pemtrexed injection is indicated: - in combination with pembrolizumab and platinum chemotherapy, for the initial treatment of patients with metastatic non-squamous non-small cell lung cancer (NSCLC), with no EGFR or ALK genomic tumor aberrations. - in combination with cisplatin for the initial treatment of patients with locally advanced or metastatic, indicated for the local treatment of unresectable cutaneous, subcutaneous, and nodal lesions in patients with melanoma recurrent after initial surgery.	300	18 years	N/A	N/A	Y	Y		5/3/2024
Biologics	J9325	Injection, talimogene laherparepvec, per 1 million plaque forming units	1 million PFU	1/1/2017	Imlygic*	talimogene laherparepvec suspension for intralesional injection		800	18 years	N/A	N/A	Y	Y		7/16/2018
Drugs	J9328	Injection, temozolomide, 1 mg	1 mg	1/1/2010	Temodar*	temozolomide for injection, for intravenous use	Indicated in adult patients for: • Treatment of newly diagnosed glioblastoma multiforme (GBM) concomitantly with radiotherapy and then as maintenance treatment. • Treatment of refractory anaplastic astrocytoma. • Adjuvant treatment of newly diagnosed anaplastic astrocytoma. (Recommended dosing is for oral	6,200	18 years	N/A	N/A	Y	Y		10/26/2023
Drugs	J9330	Injection, temsirolimus, 1 mg	1 mg	1/1/2009	Torisel*	temsirolimus injection, for intravenous use	Indicated for the treatment of advanced renal cell carcinoma.	125	N/A	N/A	N/A	Y	Y		9/25/2018
Drugs	J9331	Injection, sirolimus protein-bound particles, 1 mg	1 mg	1/1/2000	Fyarro™	sirolimus protein-bound particles for injectable suspension (albumin-bound), for intravenous use	Indicated for the treatment of adult patients with locally advanced unresectable or metastatic malignant perivascular epithelioid cell tumor (PEComa).	1,200	18 years	N/A	N/A	Y	Y		6/6/2022
Biologics	J9332	Injection, efgartigimod alfa-fcab, 2mg	2 mg	1/1/2002	Vyvgart™	efgartigimod alfa-fcab injection, for intravenous use	Indicated for the treatment of generalized myasthenia gravis (gMG) in adult patients who are anti-acetylcholine receptor (AChR) antibody positive.	2,400	18 years	N/A	N/A	Y	Y		6/6/2022
Biologics	J9333	Injection, rozanolixizumab-noli, 1 mg	1 mg	1/1/2024	Rystiggo*	rozanolixizumab-noli injection, for subcutaneous use	Indicated for the treatment of generalized myasthenia gravis (gMG) in adult patients who are anti-acetylcholine receptor (AChR) or antimuscle-specific tyrosine kinase (MuSK) antibody positive.	4,200	18 years	N/A	N/A	Y	Y		12/22/2023
Biologics	J9334	Injection, efgartigimod alfa, 2 mg and hyaluronidase-qvfc	2 mg	1/1/2024	Vyvgart* Hytulo	efgartigimod alfa and hyaluronidase-qvfc injection, for subcutaneous use	Indicated for the treatment of generalized myasthenia gravis (gMG) in adult patients who are anti-acetylcholine receptor (AChR) antibody positive.	2,016	18 years	N/A	N/A	Y	Y		12/22/2023
Drugs	J9340	Injection, thiotepa, 15 mg	15 mg	1/1/2000	N/A	thiotepa injection, powder, lyophilized, for solution	Thiotepa has been tried with varying results in the palliation of a wide variety of neoplastic diseases. However, the most consistent results have been seen in the following tumors: adenocarcinoma of the breast; adenocarcinoma of the ovary; for controlling intracavitary effusions secondary to diffuse or localized neoplastic diseases of various serosal cavities; for the treatment of superficial papillary	20	18 years	N/A	N/A	Y	Y		9/21/2018
Biologics	J9345	Injection, retifanlimab-dlwr, 1 mg	1 mg	10/1/2023	Zynyz™	retifanlimab-dlwr injection, for intravenous use	Indicated for the treatment of adult patients with metastatic or recurrent locally advanced Merkel cell carcinoma.	1,000	18 years	N/A	N/A	Y	Y	9/2023: NC Suggested Max Monthly Units updated from 500 units to 1,000 units	9/28/2023
Biologics	J9347	Injection, tremelimumab-actl, 1 mg	1 mg	7/1/2023	Imjudo*	tremelimumab-actl injection, for intravenous use	Indicated: • in combination with durvalumab, for the treatment of adult patients with unresectable hepatocellular carcinoma (uHCC). • in combination with durvalumab and platinum-based chemotherapy for the treatment of adult patients with metastatic non-small cell lung cancer (NSCLC) with no sensitizing epidermal growth factor receptor (EGFR) mutation or anaplastic lymphoma kinase (ALK) genomic tumor aberrations.	300	18 years	N/A	N/A	Y	Y		6/22/2023
Biologics	J9348	Injection, naxitamab-gagk, 1 mg	1 mg	7/1/2021	Danyelza*	naxitamab-gagk injection, for intravenous use	Indicated, in combination with granulocyte-macrophage colony-stimulating factor (GM-CSF), for the treatment of pediatric patients 1 year of age and older and adult patients with relapsed or refractory high-	800	1 year	N/A	N/A	Y	Y		6/28/2021
Biologics	J9349	Injection, tafasitamab-cxix, 2 mg	2 mg	4/1/2021	Monjuvi*	tafasitamab-cxix for injection, for intravenous use	Indicated in combination with lenalidomide for the treatment of adult patients with relapsed or refractory diffuse large B-cell lymphoma (DLBCL) not otherwise specified, including DLBCL arising from low grade lymphoma, and who are not eligible for autologous stem cell transplant (ASCT).	5,400	18 years	N/A	N/A	Y	Y		3/25/2021
Biologics	J9350	Injection, mosunetuzumab-axgb, 1 mg	1 mg	7/1/2023	Lunsumio™	mosunetuzumab-axgb injection, for intravenous use	Indicated for the treatment of adult patients with relapsed or refractory follicular lymphoma after two or more lines of systemic therapy.	123	18 years	N/A	N/A	Y	Y		6/22/2023
Drugs	J9351	Injection, topotecan, 0.1 mg	0.1 mg	1/1/2011	Hycamtin*	topotecan for injection	Indicated for: • Metastatic carcinoma of the ovary after disease progression on or after initial or subsequent chemotherapy. • Small cell lung cancer platinum-sensitive disease in patients who progressed after first-line chemotherapy. • Combination therapy with cisplatin for Stage IV-B, recurrent, or persistent carcinoma of the cervix which	400	18 years	N/A	N/A	Y	Y		9/12/2018
Drugs	J9352	Injection, trabectedin, 0.1 mg	0.1 mg	1/1/2017	Yondelis*	trabectedin for injection, for intravenous use	Indicated for the treatment of patients with unresectable or metastatic liposarcoma or leiomyosarcoma who received a prior anthracycline-containing regimen.	80	18 years	N/A	N/A	Y	Y		9/12/2018
Biologics	J9353	Injection, margetuximab-cmkb, 5 mg	5 mg	7/1/2021	Margenza™	margetuximab-cmkb injection, for intravenous use	Indicated, in combination with chemotherapy, for the treatment of adult patients with metastatic HER2-positive breast cancer who have received two or more prior anti-HER2 regimens, at least one of which was for metastatic disease.	900	18 years	N/A	N/A	Y	Y		6/28/2021

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

•Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.

•11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacture participating in the Medicaid Drug Rebate Program (MDRP).

•The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.

•The HCPCS Code effective date represents the date the HCPCS code was established

•Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacture as they are not classified as covered outpatient drugs.

•Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/ncci-medicaid/medicaid-ncci-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Biologics	J9354	Injection, ado-trastuzumab emtansine, 1 mg	1 mg	1/1/2014	Kadcyla®	ado-trastuzumab emtansine for injection, for intravenous use	Indicated, as a single agent, for the treatment of patients with HER2-positive, metastatic breast cancer who previously received trastuzumab and a taxane, separately or in combination. Patients should have either: • received prior therapy for metastatic disease, or • developed disease recurrence during or within six months of completing adjuvant therapy. • The adjuvant treatment of patients with HER2-positive early breast cancer who have residual invasive disease after neoadjuvant taxane and trastuzumab-based treatment.	1,160	18 years	N/A	N/A	Y	Y		6/4/2019
Biologics	J9355	Injection, trastuzumab, excludes biosimilar, 10 mg	10 mg	1/1/2000	Herceptin®	trastuzumab for injection, for intravenous use	Indicated for: • The treatment of HER2-overexpressing breast cancer. • The treatment of HER2-overexpressing metastatic gastric or gastroesophageal junction adenocarcinoma.	196	18 years	N/A	N/A	Y	Y		9/12/2018
Biologics	J9356	Injection, trastuzumab, 10 mg and Hyaluronidase-oysk	10 mg	7/1/2019	Herceptin Hylecta™	trastuzumab and hyaluronidase-oysk injection, for subcutaneous use	Indicated in adults for the treatment of HER2-overexpressing breast cancer. Select patients for therapy based on an FDA-approved companion diagnostic for trastuzumab.	120	18 years	N/A	N/A	Y	Y		6/3/2019
Drugs	J9357	Injection, valrubicin, intravesical, 200 mg	200 mg	1/1/2000	Valstar®	valrubicin solution, concentrate, for intravesical use	Indicated for intravesical therapy of Bacillus Calmette-Guérin (BCG) -refractory carcinoma in situ (CIS) of the urinary bladder in patients for whom immediate cystectomy would be associated with unacceptable morbidity or mortality.	20	18 years	N/A	N/A	Y	Y		9/12/2018
Biologics	J9358	Injection, fam-trastuzumab deruxtecan-nxki, 1 mg	1 mg	7/1/2020	Enhertu®	fam-trastuzumab deruxtecan-nxki for injection, for intravenous use	Indicated for the treatment of: • adult patients with unresectable or metastatic HER2-positive breast cancer who have received a prior anti-HER2-based regimen either: - in the metastatic setting, OR - in the neoadjuvant or adjuvant setting and have developed disease recurrence during or within six	1,800	18 years	N/A	N/A	Y	Y		12/20/2022
Biologics	J9359	Injection, loncastuximab tesirine-lpyl, 0.075 mg	0.075 mg	4/1/2022	Zynlonta™	loncastuximab tesirine-lpyl for injection, for intravenous use	Indicated for the treatment of adult patients with relapsed or refractory large B-cell lymphoma after two or more lines of systemic therapy, including diffuse large B-cell lymphoma (DLBCL) not otherwise specified, DLBCL arising from low grade lymphoma, and high-grade B-cell lymphoma.	800	18 years	N/A	N/A	Y	Y		3/17/2022
Drugs	J9360	Injection, vinblastine sulfate, 1 mg	1 mg	1/1/2009	N/A	vinblastine sulfate injection	Indicated in the palliative treatment of the following: Frequently Responsive Malignancies - • Generalized Hodgkin's disease (Stages III and IV, Ann Arbor modification of Rye staging system)	250	N/A	N/A	N/A	Y	Y		9/12/2018
Drugs	J9370	Vincristine sulfate, 1 mg	1 mg	1/1/2000	Vincasar PFS®	vincristine sulfate injection solution	Indicated in acute leukemia. Vincasar PFS has also been shown to be useful in combination with other oncolytic agents in Hodgkin's disease, non Hodgkin's malignant lymphomas, rhabdomyosarcoma, neuroblastoma, and Wilms' tumor.	20	N/A	N/A	N/A	Y	Y		9/12/2018
Drugs	J9371	Injection, vincristine sulfate liposome, 1 mg	1 mg	1/1/2014	Marqibo®	vincristine sulfate liposome injection, for intravenous infusion	Indicated for the treatment of adult patients with Philadelphia chromosome-negative (Ph-) acute lymphoblastic leukemia (ALL) in second or greater relapse or whose disease has progressed following two or more anti-leukemia therapies. This indication is based on overall response rate. Clinical benefit such as improvement in overall survival has not been verified.	30	18 years	N/A	N/A	Y	Y		8/5/2021
Biologics	J9376	Injection, pozelimab-bbfg, 1 mg	1 mg	4/1/2024	Veopoz™	pozelimab-bbfg injection, for intravenous or subcutaneous use	Indicated for the treatment of adult and pediatric patients 1 year of age and older with CD55-deficient protein-losing enteropathy (PLE), also known as CHAPLE disease.	4,000	1 year	N/A	N/A	Y	Y		4/12/2024

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

•Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.

•11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacture participating in the Medicaid Drug Rebate Program (MDRP).

•The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.

•The HCPCS Code effective date represents the date the HCPCS code was established

•Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacture as they are not classified as covered outpatient drugs.

•Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/ncci-medicaid/medicaid-ncci-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Biologics	J9381	Injection, teplizumab-mzwv, 5 mcg	5 mcg	7/1/2023	Tzield™	teplizumab-mzwv injection, for intravenous use	Indicated to delay the onset of Stage 3 type 1 diabetes (T1D) in adults and pediatric patients aged 8 years and older with Stage 2 T1D.	9,600	8 years	N/A	N/A	Y	Y		6/22/2023
Drugs	J9390	Injection, vinorelbine tartrate, per 10 mg	10 mg	1/1/2000	Navelbine®	vinorelbine tartrate injection, for intravenous use	Indicated: • In combination with cisplatin for first-line treatment of patients with locally advanced or metastatic non-small cell lung cancer (NSCLC).	40	18 years	N/A	N/A	Y	Y		9/27/2018
Drugs	J9393	Injection, fulvestrant (teva), not therapeutically equivalent to J9395, 25 mg	25 mg	1/1/2023	N/A	fulvestrant injection, for intramuscular use (Teva)	Indicated for the treatment of: • Hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative advanced breast cancer in postmenopausal women not previously treated with endocrine therapy. • HR-positive advanced breast cancer in postmenopausal women with disease progression following endocrine therapy. • HR-positive, HER2-negative advanced or metastatic breast cancer in postmenopausal women in combination with ribociclib, as initial endocrine based therapy or following disease progression on endocrine therapy. • HR-positive, HER2-negative advanced or metastatic breast cancer in combination with palbociclib or abemaciclib in women with disease progression after endocrine therapy.	60	18 years	N/A	Females Only	Y	Y		12/6/2022
Drugs	J9394	Injection, fulvestrant (fresenius kabi) not therapeutically equivalent to J9395, 25 mg	25 mg	1/1/2023	N/A	fulvestrant injection, for intramuscular use (Fresenius Kabi)	Monotherapy Fulvestrant injection is indicated for the treatment of: • Hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative advanced breast cancer in postmenopausal women not previously treated with endocrine therapy, or • HR-positive advanced breast cancer in postmenopausal women with disease progression following endocrine therapy.	60	18 years	N/A	Females Only	Y	Y		12/6/2022
Drugs	J9395	Injection, fulvestrant, 25 mg	25 mg	1/1/2004	Faslodex®	fulvestrant injection, for intramuscular use	Indicated for the treatment of HR-positive advanced breast cancer in postmenopausal women with disease progression following endocrine therapy.	60	18 years	N/A	Females only	Y	Y		10/10/2018
Biologics	J9400	Injection, ziv-aflibercept, 1 mg	1 mg	1/1/2014	Zaltrap®	ziv-aflibercept injection for intravenous infusion	Indicated in combination with 5-fluorouracil, leucovorin, irinotecan-(FOLFIRI), for the treatment of patients with metastatic colorectal cancer (mCRC) that is resistant to or has progressed following an oxaliplatin-containing regimen.	1,800	18 years	N/A	N/A	Y	Y		6/7/2019
Drugs	J9600	Injection, porfimer sodium, 75 mg	75 mg	1/1/2000	Photofrin®	porfimer sodium injection	Indicated for: Esophageal Cancer • Palliation of patients with completely obstructing esophageal cancer, or of patients with partially obstructing esophageal cancer who, in the opinion of their physician, cannot be satisfactorily treated with Nd:YAG laser therapy Endobronchial Cancer • Treatment of microinvasive endobronchial non-small cell lung cancer (NSCLC) in patients for whom surgery and radiotherapy are not indicated • Reduction of obstruction and palliation of symptoms in patients with completely or partially obstructing endobronchial NSCLC High-Grade Dysplasia in Barrett's Esophagus • Ablation of high-grade dysplasia (HGD) in Barrett's esophagus (BE) patients who do not undergo esophagectomy	8	18 years	N/A	N/A	Y	Y		6/6/2019

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

•Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.

•11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacturer participating in the Medicaid Drug Rebate Program (MDRP).

•The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.

•The HCPCS Code effective date represents the date the HCPCS code was established

•Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacture as they are not classified as covered outpatient drugs.

•Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/ncci-medicaid/medicaid-ncci-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Biologicals	J9999	Not otherwise classified, antineoplastic drugs	1 mg	1/1/2000	Loqtorzi™	toripalimab-tpzi injection, for intravenous use	<p>Toripalimab-tpzi injection is indicated:</p> <ul style="list-style-type: none"> • in combination with cisplatin and gemcitabine, for first-line treatment of adults with metastatic or with recurrent locally advanced nasopharyngeal carcinoma (NPC). • as a single agent for the treatment of adults with recurrent unresectable or metastatic NPC with disease progression on or after a platinum-containing chemotherapy. 	1,440	18 years	N/A	N/A	Y	Y		12/22/2023
Biologicals	J9999	Not otherwise classified, antineoplastic drugs	1 mcg	1/1/2000	Besremi®	ropeginterferon alfa-2b-njft injection, for subcutaneous use	Indicated for the treatment of adults with polycythemia vera.	1,500	18 years	N/A	N/A	Y	Y	1/2024: Procedure code updated from J3590 to J9999 to align with product's FDA-approved indication effective 2/1/2024.	1/26/2024
Biologicals	P9041	Infusion, albumin (human), 5%, 50 mL	50 mL	1/1/2001	Albutein®, Plasbumin®	albumin (human), 5%	<p>Plasbumin: Indicated for:</p> <ul style="list-style-type: none"> • Emergency treatment of hypovolemic shock • Burn therapy • Cardiopulmonary bypass 	1,550	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	Product specific age restrictions: • Plasbumin: 18 years of age and older	9/25/2018
Biologicals	P9045	Infusion, albumin (human), 5%, 250 mL	250 mL	1/1/2002	Albuked™-5, Albuminex®, AlbuRx®, Albutein®,	albumin (human) U.S.P., 5% solution for injection - 250 mL	<p>Albuked-5: Albuked-5 is indicated for:</p> <ul style="list-style-type: none"> • Emergency treatment of hypovolemic shock • Burn therapy 	620	Pediatric Use: Ensure dose is appropriate for body weight. The safety of albumin	N/A	N/A	Y	Y		4/23/2024
Biologicals	P9046	Infusion, albumin (human), 25%, 20 mL	20 mL	1/1/2002	Albutein®	albumin (human) U.S.P., 25% solution for injection - 20 mL	<p>Albutein 25% is indicated for:</p> <ul style="list-style-type: none"> • Hypovolemia • Cardiopulmonary bypass procedures • Acute nephrosis • Hypoalbuminemia • Ovarian hyperstimulation syndrome • Neonatal hyperbilirubinemia • Adult respiratory distress syndrome (ARDS) • Prevention of central volume depletion after paracentesis due to cirrhotic ascites 	775	Pediatric Use: No human or animal data. Use only if clearly needed.	N/A	N/A	Y	Y		4/23/2024

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

•Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.

•11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacture participating in the Medicaid Drug Rebate Program (MDRP).

•The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.

•The HCPCS Code effective date represents the date the HCPCS code was established

•Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacture as they are not classified as covered outpatient drugs.

•Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/nccli-medicaid/medicaid-nccli-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Biologics	P9047	Infusion, albumin (human), 25%, 50 mL	50 mL	1/1/2002	Albuked, Albuminar®, Albutein®, Flexbumin,	albumin (human), 25%	Plasbumin and Albuked: Indicated for: • Emergency treatment of hypovolemic shock • Burn therapy • Hypoproteinemia with or without edema	310	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	Product specific age restrictions: • Kedbumin: 12 years of age and older	9/25/2018
Drugs	Q0138	Injection, ferumoxylol, for treatment of iron deficiency anemia, 1 mg (non-ESRD use)	1 mg	1/1/2010	Feraheme®	ferumoxylol injection, for intravenous use (non-ESRD use)	• Indicated for the treatment of iron deficiency anemia in adult patients with chronic kidney disease (CKD). • Treatment of iron deficiency anemia in adult patients who have intolerance to oral iron or have had unsatisfactory response to oral iron.	1,020	18 years	N/A	N/A	Y	Y		10/26/2018
Drugs	Q0139	Injection, ferumoxylol, for treatment of iron deficiency anemia, 1 mg (for ESRD on dialysis)	1 mg	1/1/2010	Feraheme®	ferumoxylol injection, for intravenous use (ESRD use)	Indicated for the treatment of iron deficiency anemia in adult patients • With chronic kidney disease (CKD) or • Who have intolerance to oral iron or have had unsatisfactory response to oral iron.	1,020	18 years	N/A	N/A	Y	Y		10/26/2018
Drugs	Q0144	Azithromycin dihydrate, oral, capsule/powder, 1 g	1 g	1/1/2000	Zithromax®	azithromycin, oral	Approved indication for use in the PADP: • Sexually Transmitted Diseases Other FDA approved indications: Indicated for the treatment of mild to moderate infections caused by designated, susceptible bacteria: • Acute bacterial exacerbations of chronic bronchitis in adults • Acute bacterial sinusitis in adults • Uncomplicated skin and skin structure infections in adults • Urethritis and cervicitis in adults • Genital ulcer disease in men • Acute otitis media in pediatric patients • Community-acquired pneumonia in adults and pediatric patients • Pharyngitis/tonsillitis in adults and pediatric patients • Mycobacterial Infections Limitations of Use: • Azithromycin should not be used in patients with pneumonia who are judged to be inappropriate for oral therapy because of moderate to severe illness or risk factors. • To reduce the development of drug-resistant bacteria and maintain the effectiveness of azithromycin and other antibacterial drugs, azithromycin should be used only to treat infections that are proven or strongly suspected to be caused by susceptible bacteria.	2	N/A	N/A	N/A	Y	Y		6/7/2019
Biologics	Q0224	Injection, pemivibart, for the pre-exposure prophylaxis only, for certain adults and adolescents (12 years of age and older weighing at least 40 kg) with no known SARS-CoV-2 exposure, and who either have moderate-to-severe immune compromise due to a medical condition or receipt of immunosuppressive medications or treatments, and are unlikely to mount an adequate immune response to	4500 mg (1 dose)	3/22/2024	Pemgarda	pemivibart injection, for intravenous use	The U.S. FDA has issued an EUA for the emergency use of the unapproved product Pemgarda (pemivibart), a SARS-CoV-2 spike protein-directed attachment inhibitor, for the pre-exposure prophylaxis of coronavirus disease 2019 (COVID-19) in adults and adolescents (12 years of age and older weighing at least 40 kg): • who are not currently infected with SARS-CoV-2 and who have not had a known recent exposure to an individual infected with SARS-CoV-2 and: • who have moderate-to-severe immune compromise due to a medical condition or receipt of immunosuppressive medications or treatments and are unlikely to mount an adequate immune response to COVID-19 vaccination. Pemgarda has been authorized by FDA for the emergency use described above. Pemgarda is not FDA-approved for any use, including use for pre-exposure prophylaxis of COVID-19.	1	12 years	N/A	N/A	Y	N		5/3/2024

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

- Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.
- 11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacturer participating in the Medicaid Drug Rebate Program (MDRP).
- The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.
- The HCPCS Code effective date represents the date the HCPCS code was established
- Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacture as they are not classified as covered outpatient drugs.
- Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/ncci-medicaid/medicaid-ncci-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Drugs	Q2009	Injection, fosphenytoin, 50 mg phenytoin equivalent	50 mg	1/1/2001	Cerebyx*	fosphenytoin sodium injection, for intravenous or intramuscular use	Indicated for the treatment of generalized tonic-clonic status epilepticus and prevention and treatment of seizures occurring during neurosurgery. Cerebyx can also be substituted, as short-term use, for oral phenytoin. Cerebyx should be used only when oral phenytoin administration is not possible.	164	N/A	N/A	N/A	Y	Y		3/21/2022
Biologics	Q2043	Sipuleucel-T, minimum of 50 million autologous CD54+ cells activated with PAP-GM-CSF, including leukapheresis and all other preparatory procedures, per infusion	250 mL	7/1/2011	Provenge*	sipuleucel-T, suspension for intravenous infusion	Indicated for the treatment of asymptomatic or minimally symptomatic metastatic castrate-resistant (hormone refractory) prostate cancer.	3	N/A	N/A	Males Only	Y	Y		7/16/2018
Drugs	Q2050	Injection, doxorubicin hydrochloride, liposomal, not otherwise specified, 10 mg	10 mg	7/1/2013	Doxil*	doxorubicin hydrochloride liposome injection, for intravenous use	Indicated for: <ul style="list-style-type: none"> • Ovarian cancer after failure of platinum-based chemotherapy. • AIDS-related Kaposi's Sarcoma after failure of prior systemic chemotherapy or intolerance to such therapy. • Multiple Myeloma in combination with bortezomib in patients who have not previously received bortezomib and have received at least one prior therapy. 	30	18 years	N/A	N/A	Y	Y		6/10/2019

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

- Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.
- 11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacture participating in the Medicaid Drug Rebate Program (MDRP).
- The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.
- The HCPCS Code effective date represents the date the HCPCS code was established
- Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacture as they are not classified as covered outpatient drugs.
- Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/ncci-medicare/medicaid-ncci-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Biologicals	Q4081	Injection, epoetin alfa, 100 units (for ESRD on dialysis) (for renal dialysis facilities and hospital use)	100 units	1/1/2007	Epogen®, Procrit®	epoetin alfa injection, for intravenous or subcutaneous use (for ESRD on dialysis)	Indicated for treatment of anemia due to - Chronic Kidney Disease (CKD) in patients on dialysis and not on dialysis. - Zidovudine in patients with HIV-infection. - The effects of concomitant myelosuppressive chemotherapy, and upon initiation, there is a minimum of two additional months of planned chemotherapy. • Reduction of allogeneic RBC transfusions in patients undergoing elective, noncardiac, nonvascular surgery. Limitations of Use: Epoetin alfa has not been shown to improve quality of life, fatigue, or patient well-being. Not indicated for use: • In patients with cancer receiving hormonal agents, biologic products, or radiotherapy, unless also receiving concomitant myelosuppressive chemotherapy. • In patients with cancer receiving myelosuppressive chemotherapy when the anticipated outcome is cure. • In patients with cancer receiving myelosuppressive chemotherapy in whom the anemia can be managed by transfusion. • In patients scheduled for surgery who are willing to donate autologous blood. • In patients undergoing cardiac or vascular surgery. • As a substitute for RBC transfusions in patients who require immediate correction of anemia.	1,960	1 month	N/A	N/A	Y	Y		1/12/2022
Biologicals	Q5101	Injection, filgrastim-sndz, biosimilar, (Zarxio), 1 microgram	1 mcg	4/1/2018	Zarxio®	filgrastim-sndz injection, for subcutaneous or intravenous use	Indicated to: • Decrease the incidence of infection, as manifested by febrile neutropenia, in patients with nonmyeloid malignancies receiving myelosuppressive anticancer drugs associated with a significant incidence of severe neutropenia with fever. • Reduce the time to neutrophil recovery and the duration of fever, following induction or consolidation chemotherapy treatment of patients with acute myeloid leukemia (AML). • Reduce the duration of neutropenia and neutropenia-related clinical sequelae, e.g., febrile neutropenia, in patients with nonmyeloid malignancies undergoing myeloablative chemotherapy followed by bone marrow transplantation (BMT). • Mobilize autologous hematopoietic progenitor cells into the peripheral blood for collection by leukapheresis. • Reduce the incidence and duration of sequelae of severe neutropenia (e.g., fever, infections, oropharyngeal ulcers) in symptomatic patients with congenital neutropenia, cyclic neutropenia, or idiopathic neutropenia.	59,520	N/A	N/A	N/A	Y	Y		6/6/2019
Biologicals	Q5103	Injection, infliximab-dyyb, biosimilar, (Inflectra), 10 mg	10 mg	4/1/2018	Inflectra®	infliximab-dyyb lyophilized concentrate for injection, for intravenous use	Indicated for: Crohn's Disease: • reducing signs and symptoms and inducing and maintaining clinical remission in adult patients with moderately to severely active disease who have had an inadequate response to conventional therapy. • reducing the number of draining enterocutaneous and rectovaginal fistulas and maintaining fistula closure in adult patients with fistulizing disease. Pediatric Crohn's Disease: Indicated for: Crohn's Disease: • Reducing signs and symptoms and inducing and maintaining clinical remission in adult patients with moderately to severely active disease who have had an inadequate response to conventional therapy. • Reducing the number of draining enterocutaneous and rectovaginal fistulas and maintaining fistula closure in adult patients with fistulizing disease. Pediatric Crohn's Disease: • Reducing signs and symptoms and inducing and maintaining clinical remission in pediatric patients with moderately to severely active disease who have had an inadequate response to conventional therapy. Ulcerative Colitis: • Reducing signs and symptoms, inducing and maintaining clinical remission and mucosal healing, and eliminating corticosteroid use in adult patients with moderately to severely active disease who have had an inadequate response to conventional therapy. Pediatric Ulcerative Colitis: • Indicated for the treatment of anemia due to: o Chronic kidney disease (CKD) in patients on dialysis and not on dialysis. o Zidovudine in patients with HIV-infection. o The effects of concomitant myelosuppressive chemotherapy, and upon initiation, there is a minimum of two additional months of planned chemotherapy. • Indicated for the reduction of allogeneic RBC transfusions in patients undergoing elective, noncardiac, nonvascular surgery. Limitations of Use: Retacrit has not been shown to improve quality of life, fatigue, or patient well-being. Not indicated for use in:	140	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	Crohn's Disease and Ulcerative Colitis: 6 years of age and older Plaque Psoriasis, Psoriatic Arthritis, Ankylosing Spondylitis: 18 years of age and older	7/26/2019
Biologicals	Q5104	Injection, infliximab-abda, biosimilar, (Renflexis), 10 mg	10 mg	4/1/2018	Renflexis®	infliximab-abda for injection, for intravenous use	Indicated for: Crohn's Disease: • Reducing signs and symptoms and inducing and maintaining clinical remission in adult patients with moderately to severely active disease who have had an inadequate response to conventional therapy. • Reducing the number of draining enterocutaneous and rectovaginal fistulas and maintaining fistula closure in adult patients with fistulizing disease. Pediatric Crohn's Disease: • Reducing signs and symptoms and inducing and maintaining clinical remission in pediatric patients with moderately to severely active disease who have had an inadequate response to conventional therapy. Ulcerative Colitis: • Reducing signs and symptoms, inducing and maintaining clinical remission and mucosal healing, and eliminating corticosteroid use in adult patients with moderately to severely active disease who have had an inadequate response to conventional therapy. Pediatric Ulcerative Colitis: • Indicated for the treatment of anemia due to: o Chronic kidney disease (CKD) in patients on dialysis and not on dialysis. o Zidovudine in patients with HIV-infection. o The effects of concomitant myelosuppressive chemotherapy, and upon initiation, there is a minimum of two additional months of planned chemotherapy. • Indicated for the reduction of allogeneic RBC transfusions in patients undergoing elective, noncardiac, nonvascular surgery. Limitations of Use: Retacrit has not been shown to improve quality of life, fatigue, or patient well-being. Not indicated for use in:	140	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	Indication specific. • Crohn's Disease: 6 years and older • Ulcerative Colitis: 6 years and older • Rheumatoid Arthritis in combination with methotrexate: 18 years and older • Ankylosing Spondylitis: 18 years and older • Psoriatic Arthritis: 18 years and older • Plaque Psoriasis: 18 years	7/26/2019
Biologicals	Q5105	Injection, epoetin alfa-epbx, biosimilar, (Retacrit) (for esrd on dialysis), 100 units	100 units	7/1/2018	Retacrit™	epoetin alfa-epbx injection, for intravenous or subcutaneous use (for ESRD on dialysis)	Indicated for the treatment of anemia due to: o Chronic kidney disease (CKD) in patients on dialysis and not on dialysis. o Zidovudine in patients with HIV-infection. o The effects of concomitant myelosuppressive chemotherapy, and upon initiation, there is a minimum of two additional months of planned chemotherapy. • Indicated for the reduction of allogeneic RBC transfusions in patients undergoing elective, noncardiac, nonvascular surgery. Limitations of Use: Retacrit has not been shown to improve quality of life, fatigue, or patient well-being. Not indicated for use in:	1,960	1 month	N/A	N/A	Y	Y		1/12/2022

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

•Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.

•11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacture participating in the Medicaid Drug Rebate Program (MDRP).

•The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.

•The HCPCS Code effective date represents the date the HCPCS code was established

•Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacture as they are not classified as covered outpatient drugs.

•Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/ncci-medicaid/medicaid-ncci-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Biologics	Q5106	Injection, epoetin alfa-epbx, biosimilar, (retacrit) (for non-esrd use), 1000 units	1,000 units	7/1/2018	Retacrit™	epoetin alfa-epbx injection, for intravenous or subcutaneous use (for non-ESRD use)	<ul style="list-style-type: none"> Indicated for the treatment of anemia due to: <ul style="list-style-type: none"> ○ Chronic kidney disease (CKD) in patients on dialysis and not on dialysis. ○ Zidovudine in patients with HIV-infection. ○ The effects of concomitant myelosuppressive chemotherapy, and upon initiation, there is a minimum of two additional months of planned chemotherapy. Indicated for the reduction of allogeneic RBC transfusions in patients undergoing elective, noncardiac, nonvascular surgery. 	630	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	<ul style="list-style-type: none"> Indication specific age restrictions: <ul style="list-style-type: none"> • CKD not on dialysis: 1 month of age and older • Anemia due to concomitant myelosuppressive chemotherapy: 5 years of age 	1/12/2022
Biologics	Q5107	Injection, bevacizumab, (mvasi), 10 mg	10 mg	1/1/2019	Mvasi™	bevacizumab-awwb injection, for intravenous use	<ul style="list-style-type: none"> Indicated for the treatment of: <ul style="list-style-type: none"> • Metastatic colorectal cancer, in combination with intravenous fluorouracil-based chemotherapy for first- or second-line treatment. • Metastatic colorectal cancer, in combination with fluoropyrimidine-irinotecan- or fluoropyrimidine-oxaliplatin-based chemotherapy for second-line treatment in patients who have progressed on a first-line bevacizumab product-containing regimen. Limitations of Use: Mvasi is not indicated for adjuvant treatment of colon cancer. 	420	18 years	N/A	N/A	Y	Y		7/20/2022
Biologics	Q5108	Injection, pegfilgrastim-jmdb (fulphila), biosimilar, 0.5 mg	0.5 mg	10/1/2018	Fulphila™	pegfilgrastim-jmdb injection, for subcutaneous use	<ul style="list-style-type: none"> Indicated to decrease the incidence of infection, as manifested by febrile neutropenia, in patients with non-myeloid malignancies receiving myelosuppressive anti-cancer drugs associated with a clinically significant incidence of febrile neutropenia. Limitations of Use: Fulphila is not indicated for the mobilization of peripheral blood progenitor cells for hematopoietic stem cell transplantation. 	36	N/A	N/A	N/A	Y	Y		3/21/2023
Biologics	Q5110	Injection, filgrastim-aafi, biosimilar, (Nivestym), 1 microgram	1 mcg	10/1/2018	Nivestym™	filgrastim-aafi injection, for subcutaneous or intravenous use	<ul style="list-style-type: none"> Indicated to: <ul style="list-style-type: none"> • Decrease the incidence of infection, as manifested by febrile neutropenia, in patients with nonmyeloid malignancies receiving myelosuppressive anti-cancer drugs associated with a significant incidence of severe neutropenia with fever. • Reduce the time to neutrophil recovery and the duration of fever, following induction or consolidation chemotherapy treatment of patients with acute myeloid leukemia (AML). • Reduce the duration of neutropenia and neutropenia-related clinical sequelae, e.g., febrile neutropenia, in patients with nonmyeloid malignancies undergoing myeloablative chemotherapy followed by bone marrow transplantation (BMT). • Mobilize autologous hematopoietic progenitor cells into the peripheral blood for collection by leukapheresis. • Reduce the incidence and duration of sequelae of severe neutropenia (e.g., fever, infections, oropharyngeal ulcers) in symptomatic patients with congenital neutropenia, cyclic neutropenia, or idiopathic neutropenia. 	59,520	N/A	N/A	N/A	Y	Y		12/28/2018
Biologics	Q5111	Injection, pegfilgrastim-cbqv (udenyca), biosimilar, 0.5 mg	0.5 mg	1/1/2019	Udenyca®	pegfilgrastim-cbqv injection, for subcutaneous use	<ul style="list-style-type: none"> Indicated to decrease the incidence of infection, as manifested by febrile neutropenia, in patients with non-myeloid malignancies receiving myelosuppressive anti-cancer drugs associated with a clinically significant incidence of febrile neutropenia. • Increase survival in patients acutely exposed to myelosuppressive doses of radiation (Hematopoietic) 	36	N/A	N/A	N/A	Y	Y		2/27/2024
Biologics	Q5112	Injection, trastuzumab-dttb, biosimilar, (Ontruzant), 10 mg	10 mg	7/1/2019	Ontruzant®	trastuzumab-dttb for injection, for intravenous use	<ul style="list-style-type: none"> Indicated for: <ul style="list-style-type: none"> • The treatment of HER2-overexpressing breast cancer. • The treatment of HER2-overexpressing metastatic gastric or gastroesophageal junction adenocarcinoma. Select patients for therapy based on an FDA-approved companion diagnostic for a trastuzumab product. 	196	18 years	N/A	N/A	Y	Y		5/25/2020
Biologics	Q5113	Injection, trastuzumab-pkrb, biosimilar, (Herzuma), 10 mg	10 mg	7/1/2019	Herzuma®	trastuzumab-pkrb for injection, for intravenous use	<ul style="list-style-type: none"> Indicated for: <ul style="list-style-type: none"> • the treatment of HER2-overexpressing breast cancer. • the treatment of HER2-overexpressing metastatic gastric or gastroesophageal junction adenocarcinoma. Select patients for therapy based on an FDA-approved companion diagnostic for a trastuzumab product. 	196	18 years	N/A	N/A	Y	Y		4/29/2020

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

•Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.

•11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacturer participating in the Medicaid Drug Rebate Program (MDRP).

•The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.

•The HCPCS Code effective date represents the date the HCPCS code was established

•Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacture as they are not classified as covered outpatient drugs.

•Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/ncci-medicaid/medicaid-ncci-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Biologics	Q5114	Injection, trastuzumab-dkst, biosimilar, (Ogivri), 10 mg	10 mg	7/1/2019	Ogivri™	trastuzumab-dkst for injection, for intravenous use	Indicated for: <ul style="list-style-type: none"> • The treatment of HER2-overexpressing breast cancer. • The treatment of HER2-overexpressing metastatic gastric or gastroesophageal junction adenocarcinoma. Select patients for therapy based on an FDA-approved companion diagnostic for a trastuzumab product.	196	18 years	N/A	N/A	Y	Y		12/4/2019
Biologics	Q5115	Injection, rituximab-abbs, biosimilar, (Truxima), 10 mg	10 mg	7/1/2019	Truxima®	rituximab-abbs injection, for intravenous use	Indicated for the treatment of adult patients with: <ul style="list-style-type: none"> • Non-Hodgkin's Lymphoma (NHL) - Relapsed or refractory, low grade or follicular, CD20-positive B-cell NHL as a single agent. - Previously untreated follicular, CD20-positive, B-cell NHL in combination with first line chemotherapy and, in patients achieving a complete or partial response to a rituximab product in combination with chemotherapy, as single-agent maintenance therapy. - Non-progressing (including stable disease), low-grade, CD20-positive, B-cell NHL as a single agent after first-line cyclophosphamide, vincristine, and prednisone (CVP) chemotherapy. - Previously untreated diffuse large B-cell, CD20-positive NHL in combination with (cyclophosphamide, doxorubicin, vincristine, and prednisone) (CHOP) or other anthracycline-based chemotherapy regimens. 	500	18 years	N/A	N/A	Y	Y		12/4/2019
Biologics	Q5116	Injection, trastuzumab-qyyp, biosimilar, (trazimera), 10 mg	10 mg	10/1/2019	Trazimera™	trastuzumab-qyyp for injection, for intravenous use	Indicated for: <ul style="list-style-type: none"> • The treatment of HER2-overexpressing breast cancer. • The treatment of HER2-overexpressing metastatic gastric or gastroesophageal junction adenocarcinoma. 	196	18 years	N/A	N/A	Y	Y		3/26/2020
Biologics	Q5117	Injection, trastuzumab-anns, biosimilar, (kanjinti), 10 mg	10 mg	10/1/2019	Kanjinti™	trastuzumab-anns for injection, for intravenous use	Indicated for: <ul style="list-style-type: none"> • The treatment of HER2 overexpressing breast cancer. • The treatment of HER2 overexpressing metastatic gastric or gastroesophageal junction adenocarcinoma. Select patients for therapy based on an FDA-approved companion diagnostic for a trastuzumab product.	196	18 years	N/A	N/A	Y	Y		12/14/2021

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

•Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.

•11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacture participating in the Medicaid Drug Rebate Program (MDRP).

•The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.

•The HCPCS Code effective date represents the date the HCPCS code was established

•Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacture as they are not classified as covered outpatient drugs.

•Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/ncci-medicaid/medicaid-nczi-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Biologics	Q5118	Injection, bevacizumab-bvzr, biosimilar, (Zirabev), 10 mg	10 mg	10/1/2019	Zirabev™	bevacizumab-bvzr injection, for intravenous use	Indicated for the treatment of: <ul style="list-style-type: none"> Metastatic colorectal cancer, in combination with intravenous fluorouracil-based chemotherapy for first- or second-line treatment. Metastatic colorectal cancer, in combination with fluoropyrimidine-irinotecan- or fluoropyrimidine-oxaliplatin-based chemotherapy for second-line treatment in patients who have progressed on a first-line bevacizumab product-containing regimen. Unresectable, locally advanced, recurrent or metastatic non-squamous non-small cell lung cancer, in combination with carboplatin and paclitaxel for first-line treatment. Recurrent glioblastoma in adults. Metastatic renal cell carcinoma in combination with interferon alfa. Persistent, recurrent, or metastatic cervical cancer, in combination with paclitaxel and cisplatin or paclitaxel and topotecan. Epithelial ovarian, fallopian tube, or primary peritoneal cancer: <ul style="list-style-type: none"> in combination with carboplatin and paclitaxel, followed by Zirabev as a single agent, for stage III or IV disease following initial surgical resection. in combination with paclitaxel, pegylated liposomal doxorubicin, or topotecan for platinum-resistant recurrent disease who received no more than 2 prior chemotherapy regimens. in combination with carboplatin and paclitaxel or carboplatin and gemcitabine, followed by Zirabev as a single agent, for platinum-sensitive recurrent disease. Added at Request of the State Per NCCN Guidelines:	420	18 years	N/A	N/A	Y	Y		7/20/2022
Biologics	Q5119	Injection, rituximab-pvvr, biosimilar, (Ruxience), 10 mg	10 mg	7/1/2020	Ruxience™	rituximab-pvvr injection, for intravenous use	Indicated for the treatment of adult patients with: <ul style="list-style-type: none"> Non-Hodgkin's Lymphoma (NHL): <ul style="list-style-type: none"> Relapsed or refractory, low grade or follicular, CD20-positive B-cell NHL as a single agent. Previously untreated follicular, CD20-positive, B-cell NHL in combination with first line chemotherapy 	500	18 years	N/A	N/A	Y	Y		12/16/2021
Biologics	Q5120	Injection, pegfilgrastim-bmez (ziextenzo), biosimilar, 0.5 mg	0.5 mg	7/1/2020	Ziextenzo™	pegfilgrastim-bmez injection, for subcutaneous use	Indicated to: <ul style="list-style-type: none"> decrease the incidence of infection, as manifested by febrile neutropenia, in patients with nonmyeloid malignancies receiving myelosuppressive anti-cancer drugs associated with a clinically significant incidence of febrile neutropenia. increase survival in patients acutely exposed to myelosuppressive doses of radiation (Hematopoietic Syndrome of Acute Radiation Syndrome). Limitations of Use: Ziextenzo is not indicated for the mobilization of peripheral blood progenitor cells for hematopoietic stem cell transplantation.	36	N/A	N/A	N/A	Y	Y		3/22/2024
Biologics	Q5121	Injection, infliximab-axxq, biosimilar, (avsoola), 10 mg	10 mg	7/1/2020	Avsoola™	infliximab-axxq for injection, for intravenous use	Indicated for: Crohn's Disease: <ul style="list-style-type: none"> reducing signs and symptoms and inducing and maintaining clinical remission in adult patients with moderately to severely active disease who have had an inadequate response to conventional therapy. reducing the number of draining enterocutaneous and rectovaginal fistulas and maintaining fistula closure in adult patients with fistulizing disease. Pediatric Crohn's Disease: <ul style="list-style-type: none"> reducing signs and symptoms and inducing and maintaining clinical remission in pediatric patients with moderately to severely active disease who have had an inadequate response to conventional therapy. Ulcerative Colitis: <ul style="list-style-type: none"> reducing signs and symptoms, inducing and maintaining clinical remission and mucosal healing, and eliminating corticosteroid use in adult patients with moderately to severely active disease who have had an inadequate response to conventional therapy. Pediatric Ulcerative Colitis: <ul style="list-style-type: none"> reducing signs and symptoms and inducing and maintaining clinical remission in pediatric patients with moderately to severely active disease who have had an inadequate response to conventional therapy. Rheumatoid Arthritis in combination with methotrexate: <ul style="list-style-type: none"> reducing signs and symptoms, inhibiting the progression of structural damage, and improving physical function in patients with moderately to severely active disease. Ankylosing Spondylitis:	140	Indication Specific Age Restrictions (see comments)	N/A	N/A	Y	Y	Indication specific age restrictions: Crohn's disease and ulcerative colitis: 6 years of age and older RA, ankylosing spondylitis, psoriasis: 18 years of age and older	9/21/2020
Biologics	Q5122	Injection, pegfilgrastim-apgf (nyvepria), biosimilar, 0.5 mg	0.5 mg	1/1/2021	Nyvepria™	pegfilgrastim-apgf injection, for subcutaneous use	Indicated to decrease the incidence of infection, as manifested by febrile neutropenia, in patients with non-myeloid malignancies receiving myelosuppressive anti-cancer drugs associated with a clinically significant incidence of febrile neutropenia. Limitations of Use: Nyvepria is not indicated for the mobilization of peripheral blood progenitor cells for hematopoietic stem cell transplantation.	36	N/A	N/A	N/A	Y	Y		3/21/2023
Biologics	Q5123	Injection, rituximab-arrx, biosimilar, (riabni), 10 mg	10 mg	7/1/2021	Riabni™	rituximab-arrx injection, for intravenous use	Indicated for the treatment of: <ul style="list-style-type: none"> Adult patients with non-Hodgkin's Lymphoma (NHL): <ul style="list-style-type: none"> Relapsed or refractory, low grade or follicular, CD20-positive B-cell NHL as a single agent. Previously untreated follicular, CD20-positive, B-cell NHL in combination with first line chemotherapy and, in patients achieving a complete or partial response to a rituximab product in combination with chemotherapy, as single-agent maintenance therapy. 	500	18 years	N/A	N/A	Y	Y		7/20/2022

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

•Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.

•11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacture participating in the Medicaid Drug Rebate Program (MDRP).

•The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.

•The HCPCS Code effective date represents the date the HCPCS code was established

•Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacture as they are not classified as covered outpatient drugs.

•Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/ncci-medicare/medicaid-nci-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Biologics	Q5124	Injection, ranibizumab-nuna, biosimilar, (byooviz), 0.1 mg	0.1 mg	4/1/2022	Byooviz™	ranibizumab-nuna injection, for intravitreal use	Indicated for the treatment of patients with: - Neovascular (Wet) Age-Related Macular Degeneration (AMD) - Macular Edema Following Retinal Vein Occlusion (RVO) - Myopic Choroidal Neovascularization (mCNV)	20	18 years	N/A	N/A	Y	Y		6/20/2022
Biologics	Q5125	Injection, filgrastim-ayow, biosimilar, (releuko), 1 microgram	1 mcg	10/1/2022	Releuko®	filgrastim-ayow injection, for subcutaneous or intravenous use	Indicated to: • Decrease the incidence of infection, as manifested by febrile neutropenia, in patients with nonmyeloid malignancies receiving myelosuppressive anti-cancer drugs associated with a significant incidence of severe neutropenia with fever. • Reduce the time to neutrophil recovery and the duration of fever, following induction or consolidation chemotherapy treatment of patients with acute myeloid leukemia (AML). • Reduce the duration of neutropenia and neutropenia-related clinical sequelae, e.g., febrile neutropenia.	59,520	N/A	N/A	N/A	Y	Y		9/15/2022
Biologics	Q5126	Injection, bevacizumab-maly, biosimilar, (alymys), 10 mg	10 mg	1/1/2023	Alymys®	bevacizumab-maly injection, for intravenous use	Indicated for the treatment of: • Metastatic colorectal cancer, in combination with intravenous fluorouracil-based chemotherapy for first- or second-line treatment. • Metastatic colorectal cancer, in combination with fluoropyrimidine-irinotecan- or fluoropyrimidine-	420	18 years	N/A	N/A	Y	Y		12/12/2022
Biologics	Q5127	Injection, pegfilgrastim-fpgk (stimufend), biosimilar, 0.5 mg	0.5 mg	4/1/2023	Stimufend®	pegfilgrastim-fpgk injection, for subcutaneous use	Indicated to: • Decrease the incidence of infection, as manifested by febrile neutropenia, in patients with non-myeloid malignancies receiving myelosuppressive anti-cancer drugs associated with a clinically significant incidence of febrile neutropenia.	36	N/A	N/A	N/A	Y	Y		10/26/2023
Biologics	Q5128	Injection, ranibizumab-eqrn (cimerli), biosimilar, 0.1 mg	0.1 mg	4/1/2023	Cimerli™	ranibizumab-eqrn injection, for intravitreal use	Indicated for the treatment of patients with: - Neovascular (Wet) Age-Related Macular Degeneration (AMD) - Macular Edema Following Retinal Vein Occlusion (RVO) - Diabetic Macular Edema (DME) - Diabetic Retinopathy (DR) - Myopic Choroidal Neovascularization (mCNV)	20	18 years	N/A	N/A	Y	Y		3/16/2023
Biologics	Q5129	Injection, bevacizumab-adcd (vegzelma), biosimilar, 10 mg	10 mg	4/1/2023	Vegzelma®	bevacizumab-adcd injection, for intravenous use	Indicated for the treatment of: • Metastatic colorectal cancer, in combination with intravenous fluorouracil-based chemotherapy for first- or second-line treatment. • Metastatic colorectal cancer, in combination with fluoropyrimidine-irinotecan- or fluoropyrimidine-oxaliplatin-based chemotherapy for second-line treatment in patients who have progressed on a first-line bevacizumab product-containing regimen. - Limitations of Use: Vegzelma is not indicated for adjuvant treatment of colon cancer. • Unresectable, locally advanced, recurrent or metastatic non-squamous non-small cell lung cancer, in combination with carboplatin and paclitaxel for first-line treatment. • Recurrent glioblastoma in adults. • Metastatic renal cell carcinoma in combination with interferon alfa. • Persistent, recurrent, or metastatic cervical cancer, in combination with paclitaxel and cisplatin, or paclitaxel and topotecan. • Epithelial ovarian, fallopian tube, or primary peritoneal cancer: o in combination with carboplatin and paclitaxel, followed by Vegzelma as a single agent, for stage III or IV disease following initial surgical resection o in combination with paclitaxel, pegylated liposomal doxorubicin, or topotecan for platinum-resistant recurrent disease who received no more than 2 prior chemotherapy regimens	420	18 years	N/A	N/A	Y	Y		5/25/2023
Biologics	Q5130	Injection, pegfilgrastim-pbbk (fynetra), biosimilar, 0.5 mg	0.5 mg	4/1/2023	Fynetra®	pegfilgrastim-pbbk injection, for subcutaneous use	Indicated to decrease the incidence of infection, as manifested by febrile neutropenia, in patients with nonmyeloid malignancies receiving myelosuppressive anti-cancer drugs associated with a clinically	36	N/A	N/A	N/A	Y	Y		5/25/2023

**North Carolina Division of Health Benefits
Physician Administered Drug Program Catalog**

- Unless otherwise indicated, the catalog contains procedure codes representing drugs, biologics, devices and vaccines which are only covered for FDA approved indications. Covered indications that are not FDA approved are identified with **.
- 11 digit National Drug Codes (NDCs) are required to be billed along with their corresponding procedure code. Drugs and biologics must be classified as CMS covered outpatient drugs from a labeler/manufacturer participating in the Medicaid Drug Rebate Program (MDRP).
- The Max Daily Units for radiopharmaceuticals represents one therapeutic dose or diagnostic dose.
- The HCPCS Code effective date represents the date the HCPCS code was established
- Procedure codes for covered devices and vaccines are not required to be from a rebating labeler/manufacturer as they are not classified as covered outpatient drugs.
- Medically Unlikely Edits (MUEs) are used by NC Medicaid to reduce the improper payment for medical drug claims. A MUE for a HCPCS/CPT code is the maximum units of service that a provider would report under most circumstances for a single beneficiary on a single date of service. Not all HCPCS/CPT codes have a MUE. CMS publishes MUE values on its website: <https://www.cms.gov/medicare/coding-billing/ncci-medicare/medicaid-ncci-edit-files>

Category	HCPCS Code	HCPCS Description	HCPCS Code Billing Unit	HCPCS Effective Date	Brand Name	Generic Name	FDA Approved Indications (See Package Insert for full FDA approved indication descriptions)	NC Suggested Max Monthly Units	Minimum Age	Maximum Age	Gender Restrictions	NDC Required	Rebating Labeler Required	Comments	Last Modified Date
Drugs	Q9991	Injection, buprenorphine extended-release (Sublocade), less than or equal to 100 mg	less than or equal to 100 mg	7/1/2018	Sublocade™	buprenorphine extended-release injection, for subcutaneous use, less than or equal to 100 mg	Indicated for the treatment of moderate to severe opioid use disorder in patients who have initiated treatment with a transmucosal buprenorphine-containing product, followed by dose adjustment for a minimum of 7 days.	2	18 years	N/A	N/A	Y	Y		9/27/2018
Drugs	Q9992	Injection, buprenorphine extended-release (Sublocade), greater than 100 mg	greater than 100 mg	7/1/2018	Sublocade™	buprenorphine extended-release injection, for subcutaneous use, greater	Indicated for the treatment of moderate to severe opioid use disorder in patients who have initiated treatment with a transmucosal buprenorphine-containing product, followed by dose adjustment for a minimum of 7 days.	2	18 years	N/A	N/A	Y	Y		9/27/2018
Drugs	S0D13	Esketamine, nasal spray, 1 mg	1 mg	1/1/2021	Spravato™	esketamine nasal spray	<ul style="list-style-type: none"> • Indicated, in conjunction with an oral antidepressant, for the treatment of treatment-resistant depression (TRD) in adults. • Indicated for depressive symptoms in adults with major depressive disorder (MDD) with acute suicidal ideation or behavior. Limitations of Use: Spravato is not approved as an anesthetic agent. The safety and effectiveness of	728	18 years	N/A	N/A	Y	Y		12/28/2020
Drugs	S0O28	Injection, famotidine, 20 mg	20 mg	1/1/2000	Pepcid®	famotidine injection	Indicated in some hospitalized patients with pathological hypersecretory conditions or intractable ulcers, or as an alternative to the oral dosage forms for short term use in patients who are unable to take oral	62	1 year	N/A	N/A	Y	Y	11/2020 Coverage effective 1/1/2019 per DHB request	11/10/2023
Drugs	S0O80	Injection, pentamidine isethionate, 300 mg	300 mg	1/1/2000	Pentam® 300	pentamidine isethionate for injection	Indicated for the treatment and prevention of pneumonia caused by Pneumocystis carinii.	42	4 months	N/A	N/A	Y	Y		8/24/2018
Biologicals	S0145	Injection, pegylated interferon alfa-2a, 180 mcg per mL	180 mcg	7/1/2005	Pegasys®	peginterferon alfa-2a injection, for subcutaneous use	Chronic Hepatitis C (CHC): •Adult Patients: In combination therapy with other hepatitis C virus drugs for adults with compensated	5	Indication Specific Age Restrictions	N/A	N/A	Y	Y	Indication specific age restrictions:	7/2/2018
Biologicals	S0148	Injection, pegylated interferon alfa-2b, 10 mcg	10 mcg	10/1/2010	PegIntron®	peginterferon alfa-2b injection, for subcutaneous use	Indicated for treatment of Chronic Hepatitis C (CHC) in patients with compensated liver disease.	105	3 years	N/A	N/A	Y	Y		6/7/2019
Drugs	S0189	Testosterone pellet, 75 mg	75 mg	1/1/2002	Testopel®	testosterone pellets for subcutaneous implantation	Indicated for replacement therapy in conditions associated with a deficiency or absence of endogenous testosterone: • Primary hypogonadism (congenital or acquired) - testicular failure due to cryptorchidism, bilateral torsion, orchitis, vanishing testes syndrome; or orchiectomy. • Hypogonadotropic hypogonadism (congenital or acquired) - gonadotropic LHRH deficiency, or pituitary - hypothalamic injury from tumors, trauma or radiation.	6	N/A	N/A	Males Only	Y	Y		9/21/2018
Drugs	S0190	Mifepristone, oral, 200 mg	200 mg	1/1/2000	Mifeprex®	mifepristone tablets, for oral use	Indicated, in a regimen with misoprostol, for the medical termination of intrauterine pregnancy through 70 days gestation.	1	N/A	N/A	Females Only	Y	Y		3/15/2019
Drugs	S0191	Misoprostol, oral, 200 mcg	200 mcg	1/1/2000	Cytotec®	misoprostol tablets, for oral use	Indicated, in a regimen with mifepristone, for the medical termination of intrauterine pregnancy through 70 days gestation.	4	N/A	N/A	Females Only	Y	Y	Only covered for non-FDA approved indication in the PADP program	11/30/2021
Drugs	S4993	Contraceptive pills for birth control	1 pack (1 pack = 21- or 28-tablet pack; 3 packs = 91-tablet pack)	4/1/2002	N/A	contraceptive pills for birth control	Indicated as birth control.	14 in a 12-month interval	8 years	55 years	Females Only	Y	Y	3/2024: Effective 2/1/2024, HCPCS billing unit of 1 pack clarified to be defined as 1 pack = 21- or 28-tablet pack and 3 packs = 91-tablet pack. Suggested max monthly updated to match NCTracks 14 packs per year, effective 7/1/2019. Use of code limited to LHDS.	5/21/2024